

Transcriptional Profiles of the Human Pathogenic Fungus *Paracoccidioides brasiliensis* in Mycelium and Yeast Cells*[§]

Received for publication, January 18, 2005, and in revised form, March 15, 2005
Published, JBC Papers in Press, April 22, 2005, DOI 10.1074/jbc.M500625200

Maria Sueli S. Felipe,^{a,b} Rosângela V. Andrade,^{a,c} Fabrício B. M. Arraes,^{a,c} André M. Nicola,^{a,c} Andréa Q. Maranhão,^a Fernando A. G. Torres,^a Ildinete Silva-Pereira,^a Márcio J. Poças-Fonseca,^a Élide G. Campos,^a Lídia M. P. Moraes,^a Patrícia A. Andrade,^a Aldo H. F. P. Tavares,^a Simoneide S. Silva,^a Cynthia M. Kyaw,^a Diorge P. Souza,^a PbGenome Network,^d Maristela Pereira,^a Rosália S. A. Jesuino,^e Edmar V. Andrade,^e Juliana A. Parente,^e Gisele S. Oliveira,^e Mônica S. Barbosa,^e Natália F. Martins,^f Ana L. Fachin,^g Renato S. Cardoso,^g Geraldo A. S. Passos,^{g,h} Nalvo F. Almeida,ⁱ Maria Emília M. T. Walter,^j Célia M. A. Soares,^e Maria José A. Carvalho,^{a,c} and Marcelo M. Brígido^{a,c}

From the ^aDepartamento de Biologia Celular, Universidade de Brasília, 70910-900, Brasília, DF, Brazil, ^jDepartamento de Ciência da Computação, Universidade de Brasília, 70910-900, Brasília, DF, Brazil, ^fEmbrapa-Recursos Genéticos e Biotecnologia, W5 Norte, 70770-900, Brasília, DF, Brazil, ^eDepartamento de Bioquímica, Universidade Federal de Goiás, 74001-970, Goiânia, GO, Brazil, ⁱDepartamento de Computação e Estatística, Universidade Federal de Mato Grosso do Sul, 79070-900, Campo Grande, Mississippi, Brazil, ^gDepartamento de Genética, Universidade de São Paulo, 14040-900, Ribeirão Preto, SP, Brazil, and ^hFaculdade de Odontologia, Universidade de São Paulo, 14040-900, Ribeirão Preto, SP, Brazil

***Paracoccidioides brasiliensis* is the causative agent of paracoccidioidomycosis, a disease that affects 10 million individuals in Latin America. This report depicts the results of the analysis of 6,022 assembled groups from mycelium and yeast phase expressed sequence tags, covering about 80% of the estimated genome of this dimorphic, thermo-regulated fungus. The data provide a comprehensive view of the fungal metabolism, including overexpressed transcripts, stage-specific genes, and also those that are up- or down-regulated as assessed by *in silico* electronic subtraction and cDNA microarrays. Also, a significant differential expression pattern in mycelium and yeast cells was detected, which was confirmed by Northern blot analysis, providing insights into differential metabolic adaptations. The overall transcriptome analysis provided information about sequences related to the cell cycle, stress response, drug resistance, and signal transduction pathways of the**

pathogen. Novel *P. brasiliensis* genes have been identified, probably corresponding to proteins that should be addressed as virulence factor candidates and potential new drug targets.

The dimorphic human pathogenic fungus *Paracoccidioides brasiliensis* is the etiological agent of paracoccidioidomycosis (PCM)¹ (1), a major health problem in Latin America. High positive skin tests (75%) in the adult population reinforce the importance of the mycosis in endemic rural areas, where it has been estimated to affect around 10 million individuals, 2% of whom will develop the fatal acute or chronic disease (2). The acute form of PCM chiefly compromises the reticuloendothelial system; the chronic form mainly affects adult males with a high frequency of pulmonary and/or mucocutaneous involvement (1). Chronic severe multifocal PCM may also cause granulomatous lesions in the central nervous system (3). Regardless of the affected organ, PCM usually evolves to the formation of fibrotic sequelae, permanently hindering the patient's health.

P. brasiliensis Undergoes a Dimorphic Process *in Vivo*—It is assumed that the fungus exists as a soil saprophyte, producing propagules that can infect humans and produce disease after transition to the pathogenic yeast form (4). Pathogenicity has been intimately associated with this process, since *P. brasiliensis* strains unable to differentiate into the yeast form are avirulent (5). Mammalian estrogens inhibit dimorphism, explaining the lower incidence of disease in females (6). The mycelium-to-yeast transition in *P. brasiliensis* is governed by the rise in temperature that occurs upon contact of mycelia or conidia with the human host. *In vitro*, it can be reversibly reproduced by shifting the growth temperature between 22 and 36 °C. Molecular events related to genes that control signal transduction, cell wall synthesis, and integrity are likely to be involved in this dimorphic transition.

¹ The abbreviations used are: PCM, paracoccidioidomycosis; contig, group of overlapping clones; EST, expressed sequence tag; PbAEST, *P. brasiliensis* assembled EST sequence; MAPK, mitogen-activated protein kinase.

* This work was supported by MCT, CNPq, CAPES, FUB, UFG, and FUNDECT-MS.

[§] The on-line version of this article (available at <http://www.jbc.org>) contains nine additional tables.

The nucleotide sequence(s) reported in this paper has been submitted to the GenBank™/EBI Data Bank with accession number(s) CA580326-CA584263, CN238087-CN253933, and CN373644-CN373755.

Minimal information about cDNA microarray experiments was deposited in the MIAMEExpress databank (EMBL) under the accession numbers E-MEXP-103 and A-MEXP-71. The sequences are also available at <https://www.biomol.unb.br/Pb>.

^b To whom correspondence should be addressed. Tel.: 55-307-2423; Fax: 55-61-3498411; E-mail: msueli@unb.br.

^c These authors contributed equally to this work.

^d PbGenome Network: Alda Maria T. Ferreira, Alessandra Dantas, Alessandra J. Baptista, Alexandre M. Bailão, Ana Lídia Bonato, André C. Amaral, Bruno S. Daher, Camila M. Silva, Christiane S. Costa, Clayton L. Borges, Cléber O. Soares, Cristina M. Junta, Daniel A. S. Anjos, Edans F. O. Sandes, Eduardo A. Donadi, Elza T. Sakamoto-Hojo, Flávio R. Araújo, Flávia C. Albuquerque, Gina C. Oliveira, João Ricardo M. Almeida, Juliana C. Oliveira, Kláudia G. Jorge, Larissa Fernandes, Lorena S. Derengowski, Luís Artur M. Bataus, Marcus A. M. Araújo, Marcus K. Inoue, Marlene T. De-Souza, Mauro F. Almeida, Nádia S. Parachin, Nadya S. Castro, Odair P. Martins, Patrícia L. N. Costa, Paula Sandrin-Garcia, Renata B. A. Soares, Stephano S. Mello, and Viviane C. B. Reis.

P. brasiliensis genome size was estimated to be ~30 Mb (7). A study of *P. brasiliensis* gene density suggests that this fungus contains between 7,500 and 9,000 genes,² which is in agreement with the estimated gene number for ascomycete fungi genomes.

Here are presented the results of an effort to achieve a comprehensive metabolic view of the *P. brasiliensis* dimorphic life cycle based on analysis of 6,022 groups generated from both mycelium and yeast phases. This view arises from both a general metabolism perspective and the identification of the precise metabolic points that distinguish both morphological phases. Overexpressed genes and those that are up- or down-regulated in both stages were identified. Expression levels were assessed by cDNA microarrays and some were confirmed by Northern blot. Drug targets and genes related to virulence were also detected in several metabolic pathways. Finally, the majority of genes involved in signal transduction pathways (cAMP/protein kinase A, Ca²⁺/calmodulin, and MAPKs) possibly participating in cell differentiation and infection were annotated, and now we are able to describe the corresponding signaling systems in *P. brasiliensis*.

MATERIALS AND METHODS

Fungus—*P. brasiliensis* isolate Pb01 (ATCC MYA-826) was grown at either 22 °C in the mycelium form (14 days) or 36 °C as yeast (7 days) in semisolid Fava Neto's medium. Following incubation, cells were collected for immediate RNA extraction with Trizol reagent (Invitrogen).

Construction of cDNA Libraries and Sequencing—Poly(A)⁺ mRNA was isolated from total mycelium and yeast RNA through oligo(dT)-cellulose columns (Stratagene). Unidirectional cDNA libraries were constructed in λZAPII following supplier's instructions (Stratagene). Phagemids containing fungal cDNA were then mass-excised and replicated in XL-1 Blue MRF' cells. In order to generate ESTs, single pass 5'-end sequencing of cDNAs was performed by standard fluorescence labeling dye terminator protocols with T7 flanking vector primer. Samples were loaded onto a MegaBACE 1000 DNA sequencer (Amersham Biosciences) for automated sequence analysis.

EST Processing Pipeline and Annotation—PHRED quality assessment and computational analysis were carried out as previously described (8). EST assembly was performed using the software package CAP3 (9) plus a homemade scaffolding program. Sequences of at least 100 nucleotides, with PHRED ≥20, were considered for clustering. A total of 20,271 ESTs were selected by these exclusion criteria. Contaminant and rRNA sequences were then removed to generate a set of 19,718 ESTs, which was submitted to CAP3 clustering, generating 2,655 contigs and leaving 3,367 ESTs as singlets. Contigs plus singlets comprise the base set of 6,022 *P. brasiliensis* assembled EST sequences (PbAESTs) that underwent further analysis. Annotation was carried out using a system that essentially compared these assemblies with sequences available in public databases. The BLASTX program (10) was used for annotation along with GenBankTM nonredundant (nr), cluster of orthologous groups (COG), and gene ontology (GO) data bases. The GO data base was also used to assign EC numbers to assemblies. Additionally, we used the FASTA program (11) to compare assemblies with *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* predicted polypeptides. The INTERPROSCAN program (12) was used to obtain domain and family classification of the assemblies. Metabolic pathways were analyzed using maps obtained in the KEGG Web site (13) with annotated EC numbers, and this information was used to help in assigning function to PbAESTs.

Differential Expression Analysis in Silico by Electronic Subtraction—To assign a differential expression character, the contigs formed with mycelium and yeast ESTs were statistically evaluated using a test previously described (14) with a confidence of 95%.

cDNA Microarrays and Data Analysis—A set of two microarrays containing a total of 1,152 clones in the form of PCR products was spotted in duplicate on 2.5 × 7.5-cm Hybond N⁺ nylon membranes (Amersham Biosciences). Arrays were prepared using a Generation III Array Spotter (Amersham Biosciences). Complementary DNA inserts of both *P. brasiliensis* libraries were amplified in 96-well plates using

vector-PCR amplification with T3 forward and T7 reverse universal primers. Membranes were first hybridized against the T3 [α -³³P]dCTP-labeled oligonucleotide. The amount of DNA deposited in each spot was estimated by the quantification of the obtained signals. After stripping, membranes were used for hybridization against α -³³P-labeled cDNA complex probes. The latter were prepared by reverse transcription of 10 μ g of filamentous or yeast *P. brasiliensis* total RNA using oligo(dT)₁₂₋₁₈ primer. One hundred microliters of [α -³³P]cDNA complex probe (30–50 million cpm) was hybridized against nylon microarrays. Imaging plates were scanned by a phosphor imager (Cyclone; Packard Instruments) to capture the hybridization signals. BZScan software was employed to quantify the signals with background subtraction. Spots were matched with a template grid. The ratio between vector and cDNA complex probe hybridization values for each spot was used as the reference normalization value. Total intensity normalization using the median expression value was adopted as previously described (15). Gene expression data analyzed here were obtained from three independent determinations for each phase (filamentous or yeast). We used the significance analysis of microarrays method (16) to assess the significant variations in gene expression between both mycelium and yeast. Briefly, this method is based on *t* test statistics, specially modified to high throughput analysis. A global error chance, the false discovery rate, and a gene error chance (*q* value) are calculated by the software.

Northern Blot Analysis—Total RNA (15 μ g) was separated in a 1.5% denaturing formaldehyde agarose gel and transferred to a Hybond-N nylon membrane (GE Healthcare). Probes were radiolabeled with the random primers DNA labeling system (Invitrogen) using [α -³²P]dATP. Membranes were incubated with the probes in hybridization buffer (50% formamide, 4× SSPE, 5× Denhardt's solution, 0.1% SDS, 100 μ g/ml herring sperm DNA) at 42 °C overnight and then washed twice (2× SSC, 1% SDS) at 65 °C for 1 h. Signal bands were visualized using a Typhoon 9210 phosphor imager (GE Healthcare).

URLs—Details of the results and raw data are available for download from the World Wide Web: Pbgeneome project Web site (www.biomol.unb.br/Pb); Gene Ontology Consortium (www.geneontology.org); Cluster of Orthologous Genes (www.ncbi.nlm.nih.gov/COG); INTERPROSCAN (www.ebi.ac.uk/interpro/); National Center for Biotechnology Information (www.ncbi.nlm.nih.gov/); Kyoto Encyclopedia of Genes and Genomes (www.genome.ad.jp/kegg/); BZScan Software (tagc.univ-mrs.fr); Audic and Claverie statistical test (teleton.bio.unipd.it/bioinfo/IDEG6_form/); Significance Analysis of Microarrays method (www-stat.stanford.edu/~tibs/SAM/); *Candida albicans* data base (genolist.pasteur.fr/CandidaDB/); genomes from *Aspergillus nidulans* and *Neurospora crassa* (www.broad.mit.edu/annotation/fungi/aspergillus/).

RESULTS

Transcriptome Features—In sequencing the *P. brasiliensis* transcriptome, EST data were generated from nonnormalized cDNA libraries of mycelium and yeast cells. The size range of the cDNA inserts ranged from 0.5 to 2.5 kb. Single pass 5' sequencing was performed on 25,598 cDNA clones, randomly selected from both libraries. Upon removal of bacterial and rRNA contaminant sequences, a total of 19,718 high quality ESTs underwent CAP3 assembly, yielding 2,655 contigs and 3,367 singlets, which constitute the so-called 6,022 *P. brasiliensis* Assembled EST (PbAEST) data base. Contigs presented an average size of 901 bp, and the number of ESTs assembled into contigs varied from 2 to 657 in the largest one (PbAEST 1068), which corresponds to M51, a previously reported *P. brasiliensis* mycelium-specific transcript (17). Of the 6,022 PbAESTs, 4,198 (69.4%) showed a probable homologue in GenBankTM, and 4,130 (68.3%) showed a fungus homologue (Fig. 1A and Supplemental Table I). We had used MIPS functional categories to classify 2,931 PbAESTs into 12 major groups. *P. brasiliensis* showed a slightly higher percentage of PbAESTs (4%) related to cellular communication and signal transduction (Fig. 1B) compared with *S. cerevisiae* functional categorization (3.4%).

Highly and Differentially Expressed Genes—The 27 highly transcribed genes found in the *P. brasiliensis* transcriptome, using a cut-off of 50 reads, are shown in Supplemental Table II. Some of them were previously reported (8). Also, up- and down-regulated genes in mycelium and yeast cells were detected by statistical comparison of the number of sequences in corre-

² C. Reinoso, G. Niño-Vega, G. San-Blas, and A. Dominguez (2003) IV Congreso Virtual de Micología, personal communication.




FIG. 1. *P. brasiliensis* transcriptome characterization. A, distribution of blast best hit among organisms. Each PbAEST was tested against the GenBank™ nr data base, and the best hit organism was computed. A PbAEST was considered as not assigned when the best hit exceeded an *E* value of 10^{-10} . B, functional categorization of the PbAESTs using MIPS classification. We included 2931 curator-reviewed annotations in this analysis.

sponding PbAESTs (Table I). In order to support the electronic subtraction data, cDNAs from each phase were used to probe cDNA microarrays membranes containing 1,152 clones, which were selected based on the following criteria: (i) ESTs exclusive for a particular morphotype; (ii) ESTs corresponding to genes more expressed in mycelium or yeast cells; and (iii) some ESTs equally expressed in both cell types. From the 1,152 clones, 328 genes were up-regulated during the dimorphic transition: 58 in mycelium and 270 in yeast (data not shown).

The cDNA microarray experiment confirmed most of the electronic subtraction data and also points out to new differentially expressed genes. Among them, a subclass of about 40 up-regulated genes in mycelium and yeast are described in Table I, which includes M51, M32, hydrophobin 1/2, the highly expressed yeast PbY20 protein, and some other genes that have previously been described as differentially expressed in *P. brasiliensis* by different approaches (17–20). Other key up-regulated genes related to the metabolism of *P. brasiliensis* (Table I) are described and discussed elsewhere in this work. Interestingly, we have found a yeast phase preferentially expressed gene that possibly encodes a previously characterized *P. brasiliensis* estradiol-binding protein (21), also described in *C. albicans* and in other fungi (22). It is speculated that the interaction of the 17- β -estradiol hormone with a cytoplasmic protein inhibits the mycelium-to-yeast transition, explaining the lower incidence of PCM in females.

Metabolic Overview—*P. brasiliensis* seems to be capable of producing ATP from the classical pathways of glycolysis, alcohol fermentation, and oxidative phosphorylation, since alcohol dehydrogenase, cytochrome genes, ATP synthase subunits, and pyrophosphatase genes were annotated. All genes encoding glycolytic enzymes were identified in both mycelium and yeast. Genes corresponding to the citrate cycle enzymes and to the components of complexes I, II, III, and IV were found, reflecting the ability of the fungus to perform complete aerobic pyruvate degradation and oxidative phosphorylation. Its putative capacity to also grow in anaerobiosis was evidenced by the alternative conversion of pyruvate to ethanol. Last, it may be able to utilize two-carbon sources in the form of acetate and ethanol through the glyoxylate cycle and obtain sulfite and nitrite from the environment.

In order to validate the carbon source utilization profile predicted by the transcriptome data, two *P. brasiliensis* isolates (Pb01 and Pb18) were grown in McVeigh-Morton minimum

medium supplemented with different carbon sources and growth patterns were qualitatively evaluated (Supplemental Table III). We observed that, in accordance to the transcriptome analysis prediction, several mono- and disaccharides, such as D-glucose, D-fructose, D-galactose, D-mannose, D-sorbitol, α -trehalose, maltose, and sucrose were indeed utilized. On the other hand, the predicted assimilation of D-inositol was not confirmed. Transcripts related to the consumption of L-sorbose and L-lactose were not detected; in fact, *P. brasiliensis* was unable to grow in L-sorbose as the sole carbon source. We consider that the unpredicted fungal growth in L-lactose can be explained by the fact that the *P. brasiliensis* cDNA libraries were not constructed under induction conditions. The observation that fructose, galactose, and glycerol were only utilized by Pb01 and not by Pb18 isolate may simply reflect strain biological variability as previously observed (7). A detailed description of *P. brasiliensis* metabolism, including a list of PbAESTs, is shown in Supplemental Table IV.

Differential Metabolism between Mycelium and Yeast—The up-regulated genes encoding enzymes in mycelium and yeast cells listed in Table I are highlighted in Fig. 2. The differential expression pattern of these genes (with the exception of glucokinase from mycelium cells) was confirmed by Northern blot analysis (Fig. 3). In general, the gene overexpression pattern suggests that mycelium saprophytic cells possess an aerobic metabolism, in contrast with yeast cells. Actually, mycelium up-regulated genes correspond to the main regulatory points of the citrate cycle, such as the genes coding for isocitrate dehydrogenase and succinyl-CoA synthetase; this strongly suggests a metabolic shunt to oxidative phosphorylation. Also, glucokinase is induced, producing glucose 6-phosphate, which is possibly converted through the oxidative pentose phosphate pathway to ribose 5-phosphate, and then to salvage pathways of purine and pyrimidine biosynthesis. In fact, this correlates well with the overexpression of adenylate kinase and uridine kinase genes. The excess of ribose 5-phosphate is probably converted to fructose 6-phosphate and glyceraldehyde 3-phosphate by the nonoxidative pentose phosphate pathway catalyzed by the overexpressed transaldolase. Those sugars are converted to pyruvate and acetyl-CoA for the citrate cycle in aerobic conditions.

In contrast, *P. brasiliensis* yeast cells overexpress the genes encoding alcohol dehydrogenase I and pyruvate dehydrogenase E1 subunit (Table I and Fig. 3); the latter can be detected in high levels in cultures of *S. cerevisiae* grown both anaerobically

TABLE I
Differentially expressed genes in mycelium and yeast cells detected by electronic subtraction and cDNA microarray analysis

The PbAESTs were analyzed as to their differential expression by two methods: a statistical analysis of the number of mycelium and yeast ESTs clustered in each PbAEST (14) and a cDNA microarray analysis of 1,152 PbAESTs, chosen according to the electronic subtraction criteria. A differential pattern of genes encoding enzymes was used in the analysis of the differential metabolism.

PbAEST	EC number	Annotated function	Number of reads ^a		<i>p</i> value ^b	-Fold change ^c	Accession number/Best hit organism/ <i>E</i> value
			M	Y			
Mycelium up-regulated genes							
1068		M51 ^{d,e}	653	4	0.000000	41666.0	BE758605/ <i>P. brasiliensis</i> /0.0
2274	4.4.1.5	Lactoylglutathione lyase ^e	75	0	0.000000	7.0	NP_105614.1/ <i>Mesorhizobium loti</i> /1e-11
2521		Hydrophobin 1 ^{d,f}	56	0	0.000000		AAM88289.1/ <i>P. brasiliensis</i> /2e-51
1789		HSP90 co-chaperone ^f	19	10	0.018169		CAD21185.1/ <i>N. crassa</i> /4e-48
2509	1.15.1.1	Copper-zinc superoxide dismutase ^f	14	5	0.010801		Q9Y8D9/ <i>A. fumigatus</i> /1e-68
2458		Unknown ^f	13	6	0.025336		
2478		Hydrophobin 2 ^{d,f}	9	0	0.000951		AAR11449.1/ <i>P. brasiliensis</i> /2e-70
1287	1.13.11.32	2-nitropropane dioxygenase ^f	8	1	0.008606		CAB91335.2/ <i>N. crassa</i> /e-133
1318		Amino acid permease ^e	8	0	0.001907	50.4	CAD21063.1/ <i>N. crassa</i> /0.0
1470		Unknown ^e	8	2	0.021572	20.1	
2269	2.7.4.3	Adenylate kinase ^f	5	1	0.046263		NP_011097.1/ <i>S. cerevisiae</i> /1e-42
2364		Unknown ^e	5	1	0.046263	3.6	
379		Unknown ^e	5	1	0.046263	4.9	
1092	4.2.1.22	Cystathionine β-synthase ^f	4	0	0.030842		AAL09565.1/ <i>Pichia pastoris</i> /4e-96
2356	2.2.1.2	Transaldolase ^f	4	0	0.030842		NP_013458.1/ <i>S. cerevisiae</i> /e-108
2476	3.1.2.22	Palmitoyl-protein thioesterase ^f	4	0	0.030842		I58097/ <i>H. sapiens</i> /8e-42
4135	1.1.1.41	Isocitrate dehydrogenase ^g	1	0	0.248690	3.1	O13302/ <i>Acetobacter capsulatum</i> /6e-31
5530	6.2.1.5	β-Succinyl CoA synthetase ^g	1	0	0.248690	2.7	T49777/ <i>N. crassa</i> /9e-73
4749	2.7.1.2	Glucokinase ^g	1	0	0.248690	1.7	Q92407/ <i>Aspergillus niger</i> /2e-50
4246	2.7.1.48	Uridine-kinase ^g	1	0	0.248690	2.7	T41020/ <i>S. pombe</i> /3e-28
Yeast up-regulated genes							
2536		Y20 protein ^{e,d}	27	88	0.000000	8.7	AAL50803.1/ <i>P. brasiliensis</i> /e-106
2431	1.1.1.1	Alcohol dehydrogenase I ^f	2	45	0.000000		P41747/ <i>Aspergillus flavus</i> /e-129
737	3.5.1.41	Xylanase/chitin deacetylase ^e	8	33	0.000023	2.8	NP_223015.1/ <i>Helicobacter pylori</i> /e-113
201		Putative membrane protein Nce2 ^e	0	27	0.000000	25.2	NP_015475.1/ <i>S. cerevisiae</i> /5e-08
797	3.1.6.6	Choline sulfatase ^e	3	15	0.001602	4.8	NP_248721.1/ <i>P. aeruginosa</i> /e-104
814		Glyoxylate pathway regulator ^e	0	15	0.000016	17.7	NP_009936.1/ <i>S. cerevisiae</i> /4e-37
1704		60S ribosomal protein L19 ^f	0	14	0.000032		NP_596715.1/ <i>S. pombe</i> /6e-49
1585	1.8.4.8	PAPS reductase ^e	1	12	0.000815	5.1	AAG24520.1/ <i>Penicillium chrysogenum</i> /e-121
63		Putative methyltransferase ^e	3	11	0.011314	2.5	CAD21381.1/ <i>N. crassa</i> /2e-46
778		Putative estradiol-binding protein ^e	3	11	0.011314	29.5	NP_012049.1/ <i>S. cerevisiae</i> /1e-31
136		Unknown ^{d,f}	4	10	0.030950	3.9	
767		Unknown ^e	3	10	0.017732	2.2	
701	1.2.4.1	Pyruvate dehydrogenase ^f	1	9	0.004973		Q10489/ <i>S. pombe</i> /1e-72
1724		Putative sterol transporter ^e	0	6	0.007915	29.3	NP_013748.1/ <i>S. cerevisiae</i> /4e-12
171	2.6.1.42	Branched-chain aminotransferase ^f	0	5	0.015790		NP_012078.1/ <i>S. cerevisiae</i> /7e-87
1983	1.6.5.3	NADH dehydrogenase (ubiquinone reductase) ^f	0	4	0.031496		S47150/ <i>N. crassa</i> /1e-19
244	1.1.1.69	Gluconate dehydrogenase ^f	0	4	0.031496		NP_471610.1/ <i>Listeria innocua</i> /1e-09
258	3.3.2.1	Isochorismatase ^f	0	4	0.031496		NP_436193.1/ <i>Sinorhizobium meliloti</i> /1e-20
279	2.5.1.15	Dihydropteroate synthase ^f	0	4	0.031496		T49535/ <i>N. crassa</i> /1e-38
314	2.6.1.1	Aspartate aminotransferase ^f	0	4	0.031496		NP_509047.1/ <i>Caenorhabditis elegans</i> /4e-96
555	6.2.1.3	Acyl-CoA synthetase ^f	0	4	0.031496		NP_275799.1/ <i>Methanothermobacter thermautotrophicus</i> /9e-89
756	6.3.5.7	Glutamyl-tRNA amidotransferase ^f	0	4	0.031496		Q33446/ <i>A. nidulans</i> /1e-15
865	4.1.3.1	Isocitrate lyase ^f	0	4	0.031496		AAK72548.2/ <i>Coccidioides immitis</i> /e-119
963	2.6.1.9	Histidinol-phosphate aminotransferase ^f	0	4	0.031496		P36605/ <i>S. pombe</i> /4e-87
980	3.5.1.4	Acetamidase ^f	0	4	0.031496		AAK31195.1/ <i>Aspergillus terreus</i> /2e-09
3073	1.14.13.3	Phenylacetate hydroxylase ^g	0	1	0.249998	2.3	AAF21760.1/ <i>P. chrysogenum</i> /2e-48

^a Number of mycelium (M)- and yeast (Y)-derived ESTs in the PbAEST.

^b *p* value for the Audic and Claverie test.

^c -Fold change found for the microarray experiments.

^d Previously shown to be differential by Northern blot or proteome analysis.

^e Electronic subtraction and cDNA microarray analysis; differential pattern in both analyses.

^f Electronic subtraction differential pattern and not assayed in cDNA microarray analysis.

^g Singlets that are differential in cDNA microarray analysis.

and aerobically in the presence of ethanol (23). The carbohydrate metabolism is probably shifted toward ethanol production, reflecting the anaerobic behavior of the yeast form as previously reported (24). Several pathways that provide sub-

strates for the glyoxylate cycle are up-regulated in the yeast cells (Table I and Fig. 3). First, isocitrate lyase redirects the metabolic flow using ethanol and acetate as two-carbon sources and generating oxaloacetate, which can be reconverted to glu-

involved in cell cycle and in the basic genetic information flow machinery (DNA replication, repair, recombination, transcription, RNA processing, translation, and post-translational modifications) are well conserved in comparison with their counterparts from *S. cerevisiae*. Also, sequences related to mitochondrial replication, budding, sporulation, and mating were also annotated (Supplemental Table V).

From the cell cycle-related orthologues identified in *P. brasiliensis*, those related to the structure and assembly of the cytoskeleton, chromatin structure, chromosome segregation, cyclins, and cell cycle control genes were highlighted. Genes related to the major DNA repair mechanisms found in yeast (mismatch, base excision, and recombination systems) were identified in *P. brasiliensis*, although not every component was represented, since cells were not subjected to DNA-damaging conditions. The *RAD52* gene, which plays an essential role in *S. cerevisiae* recombination, is also present in the *P. brasiliensis* transcriptome.

Among the identified transcription factors, the orthologues for *MAT*, *MCM1*, and *NsdD* are of relevance, since they are implicated in ascomycete sexual reproduction. These genes represent a strong evidence for mating in *P. brasiliensis*, so far not yet described, which is reinforced by the detection of six transcripts involved in meiotic recombination.

Stress Responses—Cell differentiation in *P. brasiliensis* requires a temperature shift, which might be associated with a stress response. We have found 48 sequences encoding molecular chaperones and their associated co-chaperones in *P. brasiliensis* transcriptome (Supplemental Table VI). These sequences were divided into nine groups: small chaperones (four genes), HSP40 (9), HSP60 (10), HSP70 (7), HSP90 (4), HSP100 (4), 14-3-3 (2), calnexin (1), and immunophilins (7). Eight of these are differentially expressed: calnexin, *cct7* (cytoplasmic hsp60) and *sba1* (HSP90/70 co-chaperone) for the mycelium form and *cpr1* (HSP90/70 co-chaperone), *hsp42*, *hsp60*, *ssc1* (HSP70), and *hsp90* for the yeast form. From these, *hsp60* and *hsp70* had been previously characterized as differentially expressed in yeast (25, 26). cDNA microarray analysis confirmed the differential expression pattern of *sba1*. Furthermore, the number of chaperone and co-chaperone ESTs is 38% larger in the yeast cDNA library than in the mycelium library. These data represent an evidence of an altogether higher expression of HSPs in yeast cells, compatible with growth at 37 °C.

Oxidative agents may cause stress and damage to *P. brasiliensis* cells. They may originate from the activity of host macrophages or from intracellular oxidative species. *P. brasiliensis* contains several genes encoding enzymes with known or putative antioxidant properties, such as superoxide dismutases, catalases (two isoenzymes), peroxiredoxins, and a novel cytochrome *c* peroxidase (Supplemental Table VII). Homologues to genes encoding secondary antioxidant enzymes belonging to the glutathione *S*-transferase family were also found. Several transcription factors may be involved in the induction of antioxidant defenses in *P. brasiliensis*. Homologues to *YAP1*, *HAP3*, and *SKN7* from *S. cerevisiae* (27) were discovered in the transcriptome, showing that the oxidative stress regulators from *P. brasiliensis* and baker's yeast might be conserved.

Signal Transduction Pathways—Transcriptome analysis and reverse annotation revealed several putative components of the biosignaling pathways in *P. brasiliensis* (Supplemental Table VIII), such as (i) MAPK signaling for cell integrity, cell wall construction, pheromone/mating, and osmotic regulation; (ii) cAMP/protein kinase A, regulating fungal development and virulence, and (iii) calcium-calmodulin-calcineurin, controlling growth at high temperature. Furthermore, a *ras* homologue sequence was detected raising the possibility of cross-talk

among the distinct signal transduction pathways (Fig. 4).

In budding yeast, the MAPK cascade responsible for cell integrity mediates cell cycle regulation and cell wall synthesis, responding to different signals including temperature, changes in external osmolarity, and mating pheromone. Components of this pathway identified in *P. brasiliensis* encompass the most classical steps, with the exception of a cell surface tyrosine kinase-like receptor that was not found in the transcriptome so far analyzed. Rho1p is a small GTP-binding protein of the *Rho* subfamily required for cell growth and coordinated regulation of cell wall construction (28) through the synthesis of β -1,3-glucan. It also activates Pkc1p, which in turn regulates the MAPK pathway.

Transcripts related to the pathway for activation by mating pheromone were identified in the *P. brasiliensis* transcriptome. The intermediary components appear to be constitutively expressed in both mycelium and yeast forms. Intriguingly, mating has not yet been described in *P. brasiliensis*. Conversely, the Hog1 MAPK cascade is activated when there is an increase in the environment osmolarity. One of its targets, Glo1p, which controls genes required for cell adaptation and survival upon osmotic stress in *S. cerevisiae* (29), was also detected in *P. brasiliensis*.

The cAMP/protein kinase A is a cascade known to regulate fungal differentiation and virulence. From the genes identified in *P. brasiliensis*, we highlight a homologue to several fungal adenylate cyclases; the low affinity cAMP phosphodiesterase, encoded by the gene *Pde1*; homologues to both the regulatory and the catalytic subunits of protein kinase A, which is involved in the regulation of the cell surface flocculin Flo11p/Muc1p (30). In *P. brasiliensis* exogenous cAMP is known to inhibit the process of filamentation (31). Both the catalytic (CnaA) and the Ca⁺²-binding regulatory B (CnaB) subunits of calcineurin were found in *P. brasiliensis*. In dimorphic fungi, cAMP- and calcineurin-dependent pathways seem to be involved in differentiation. As in the pathogenic fungus *Cryptococcus neoformans* (32), calcineurin might also play a role in mating of *P. brasiliensis*. In several pathogenic and nonpathogenic fungi, *RAS* is involved in filamentation, pseudohyphal/hyphal growth, and mating (33). A *RAS*-related transcript was identified in *P. brasiliensis*, but further studies are required to elucidate its function in mycelium-to-yeast transition and in the mechanism of pathogenicity.

Virulence Genes, Drug Targets and Resistance—In order to identify genes that could be related to *P. brasiliensis* virulence, its transcriptome has been searched for orthologues assigned as virulence factors in human pathogenic fungi, as defined by Falkow's postulate (34). Table II lists 28 *P. brasiliensis* sequences, which were previously experimentally established as virulence or essential genes in *C. albicans*, *C. neoformans*, and *Aspergillus fumigatus*. They were subdivided into four classes: metabolism-, cell wall-, and signal transduction-related and others. Some of these genes has been considered for antifungal therapy and are also listed in Table III as potential drug targets.

MAPK-related sequences, whose orthologues in *C. albicans* were experimentally correlated to hyphal formation and virulence, were also detected. The extrapolation to the *P. brasiliensis* model is not direct, since yeast, not hyphae, is the pathogenic cell type, but several MAPK homologues are found in species exhibiting diverse morphology and infection habits (35). A *cavps34* orthologue, identified in *P. brasiliensis* transcriptome (*vps34*), is implicated in the protein/lipid transport from the Golgi apparatus/endosome to the vacuole and has been proved to be important to *C. albicans* virulence (36).

Noteworthy is the finding of glyoxylate cycle genes in




FIG. 4. **Signaling pathways in *P. brasiliensis*.** Shown are cAMP/protein kinase A regulating fungal development and virulence; MAPK signaling for cell integrity, cell wall construction, pheromone/mating, and osmoregulation; calcium-calmodulin-calcineurin controlling cell survival under stress conditions; and Ras allowing cross-talk of extracellular signals. For abbreviations of gene names see Supplemental Table VIII.

P. brasiliensis, since its activity has been reported as a fungal virulence requirement (37). The activity of the key enzymes malate synthase and isocitrate lyase was reported to be up-regulated in *C. albicans* upon phagocytosis (38). Both enzymes were detected in the *P. brasiliensis* transcriptome, with isocitrate lyase being overexpressed in the yeast phase, as confirmed by Northern blot analysis (Fig. 3).

The cell wall, as the most obvious difference between human and fungal cells, represents a prime target for antifungals. Genes involved in its biogenesis and assembly can act as virulence factors and therefore are putative drug targets. We have identified orthologues to chitin synthase 3 (*chs3*), glucosamine-6-phosphate acetyltransferase (*gna1*), mannosyltransferase (*pmt1*), and α 1,2-mannosyltransferase (*mnt1*) genes and glycosidases Phr1p and Phr2p. The expression of the two last genes in *C. albicans* is responsive to the product of *prp2* (39), a pH-related transcription factor also present in the *P. brasiliensis* transcriptome. The detection of chitin deacetylase, as an overexpressed yeast gene confirmed by cDNA microarray and Northern blot (data not shown), points out to a novel target for drug research in *P. brasiliensis*.

Microbe resistance to reactive oxygen and nitrogen intermediates plays an important role in virulence (40). We were able to identify sequences that are oxidative stress response orthologues, including an alternative oxidase (*aox1*), a copper/zinc superoxide dismutase (*sod1*), and two different catalase orthologues, one of them a peroxisomal *cat1*, as recently described (41).

The urate oxidase gene detected in the *P. brasiliensis* transcriptome, but not in *S. cerevisiae*, *C. albicans*, and *Homo sapiens* genomes, suggests that uric acid could be degraded to allantoin. In addition, the presence of a *C. neoformans* urease

orthologue also probably reflects the degradation of urea to ammonia and carbamate. A role in virulence and sporulation has been assigned for both genes (42). The production of urea has been involved in an improved *in vitro* survival for those microorganisms exposed to an acidic environment. In this view, it could be related to the survival of the fungus in the host cells.

The development of new drugs is crucial, considering the problem of emerging drug resistance and toxicity (37). Novel drug targets have been found through the analysis of genome sequences. The genes listed in Table III have no homologues in the human genome and therefore could be considered for the development of new antifungal drugs. Most therapies designed to treat fungal infections target the ergosterol biosynthetic pathway (43). The orthologue of C-24 sterol methyltransferase (*ERG6*) is present in *P. brasiliensis*. In addition, modulation of sphingolipid metabolism exerts a deep impact on cell viability. The synthesis of inositol-phosphoryl-ceramide from phytoceramide catalyzed by the product of the *aur1* gene, present in *P. brasiliensis*, corresponds to the first specific step of this pathway (44). Translation elongation factors have also been pointed out as drug targets (37). In the *P. brasiliensis* transcriptome, we have found an elongation factor-3 sequence that is absent in human genome (45) and thus can be addressed for pharmaceutical purposes.

Twenty PbAESTs annotated as related to multiple drug resistance genes were identified (Supplemental Table IX). They include 12 *S. cerevisiae* orthologues, 10 of which are related to the ABC transporter and two to major facilitator superfamilies (46). One of them corresponds to Pfr1, a gene recently described in *P. brasiliensis* (47), and another is related to the *CDR1* gene from *C. albicans*, which is up-regulated in the presence of human steroid hormones (48). It has been speculated that

TABLE II
Putative virulence or essential genes found in *P. brasiliensis* transcriptome related to the experimentally confirmed orthologues of *C. albicans*, *C. neoformans*, and/or *A. fumigatus*

PBAEST	Orthologue name	AC number/Organism	E value ^a	Remarks
Metabolic genes				
2403	<i>ura3^b</i>	DCCKA (<i>C. albicans</i>) O13410 (<i>A. fumigatus</i>)	3e-41 2e-83	
0670	<i>nmt^b</i>	AAA34351 (<i>C. albicans</i>) AAA17547 (<i>C. neoformans</i>)	8e-60 1e-60	Lipid synthesis
3750	<i>fas2^b</i>	JC4086 (<i>C. albicans</i>)	7e-33	
1224	<i>hem3</i>	094048 (<i>C. albicans</i>)	1e-58	Hemosynthesis
3819	<i>tps1^b</i>	CAA69223 (<i>C. albicans</i>)	1e-36	Glucose metabolism
1693	<i>icl1</i>	AAF34690 (<i>C. albicans</i>)	1e-112	Glyoxylate cycle
0831	<i>mls1</i>	AAF34695 (<i>C. albicans</i>)	1e-122	Glyoxylate cycle
1735	<i>pabaA^b</i>	AAD31929 (<i>A. fumigatus</i>)	1e-12	Purine synthesis
Cell wall genes				
4346	<i>chs3</i>	P30573 (<i>C. albicans</i>)	7e-22	Potential drug targets
4968	<i>gna1^b</i>	BAA36496 (<i>C. albicans</i>)	4e-16	
1067	<i>mnt1</i>	CAA67930 (<i>C. albicans</i>)	9e-49	
2980	<i>pmt1</i>	AAC31119 (<i>C. albicans</i>)	4e-46	
2382	<i>phr1</i>	AAF73430 (<i>C. albicans</i>)	2e-40*	
1375	<i>phr2</i>	AAB80716 (<i>C. albicans</i>)	1e-114	
Signal transduction				
4452	<i>cek1</i>	A47211 (<i>C. albicans</i>)	3e-30	Hyphal formation
1110	<i>cpp1</i>	P43078 (<i>C. albicans</i>)	6e-16	
267	<i>cst20</i>	AAB38875 (<i>C. albicans</i>)	6e-48	
358	<i>hog1^b</i>	Q92207 (<i>C. albicans</i>)	2e-59	Osmoregulation
988	<i>nik1^b</i>	AAC72284 (<i>C. albicans</i>)	7e-37	Hyphal development
Other fungal virulence determinant genes				
623	<i>cat1^b</i>	CAA07164 (<i>C. albicans</i>)	1e-172	Peroxisomal catalase
3553	<i>mdr1^b</i>	CAA76194 (<i>C. albicans</i>)	2e-27	
3306	<i>plb1^b</i>	AAF08980 (<i>C. albicans</i>)	2e-38	Important in host cell penetration
4267	<i>top1^b</i>	Q00313 (<i>C. albicans</i>)	4e-56	
5012	<i>vps34^b</i>	CAA70254 (<i>C. albicans</i>)	2e-29	Vesicle trafficking
2516	<i>sod1^b</i>	AAK01665 (<i>C. neoformans</i>)	4e-51	Nitric oxide detoxification
2463	<i>ure1^b</i>	AAC62257 (<i>C. neoformans</i>)	6e-76	
1102	<i>aox1^b</i>	AAM22475 (<i>C. neoformans</i>)	2e-48	Resistance to oxidative stress

^a All *P. brasiliensis* assembled ESTs are BBH with *C. albicans* orthologues, except *phr1* (marked with an asterisk).

^b Putatively novel *P. brasiliensis* virulence genes.

TABLE III
Potential drug targets genes found in *P. brasiliensis* transcriptome with no homologues in the human genome

PbAEST	Annotated function	Orthologue accession numbers	E-value	Remarks
Cell wall				
5198	β -1,3-glucan synthase	AAD37783	2e-108	Preferentially expressed in mycelium
4988	α -1,3-glucan synthase	AAL18964	2e-70	Preferentially expressed in yeast
0265	Rho	AAK08118	2e-92	Signal transduction
1147	Chitin synthase I	AAF82801	2e-81	
1927	Chitin synthase II	Q92444	3e-66	
4346	Chitin synthase IV	AF107624	2e-65	
3958	Chitin synthase asmA	JC5546	1e-64	
0737	Xylanase/Chitin deacetylase	ZP_00126582	1e-12	Up-regulated in <i>P. brasiliensis</i> yeast cells
5473	Bud neck involved	NP_014166	1e-12	Required to link CHS3p and CHS4p to the septins
1063	α -1,2-Mannosyltransferase	NP_009764	1e-20	Involved in protein glycosylation
Glyoxylate cycle				
2402	Malate synthase	P28344	1e-37	
1688	Isocitrate lyase	AAK72548.2	1e-144	Up-regulated in <i>P. brasiliensis</i> yeast cells
Other targets				
1959	Δ (24)-Sterol C-methyltransferase	T50969	4e-44	Ergosterol biosynthesis
0200	Aureobasidin resistance protein	AAD22750	1e-43	Sphingolipid synthesis
0845	Elongation factor 3	BAA33893	1e-142	Unique and essentially required for fungal translational machinery
4129	Urate oxidase	P33282	6e-77	Sporulation and pathogenesis
2456	Urease	AAC49868	3e-94	Sporulation and pathogenesis

steroid hormones are involved in morphological changes as well as in pathogenicity in *P. brasiliensis* and also in drug resistance in *C. albicans*. Interestingly, the process of infection of *P. brasiliensis* is strongly biased toward males, albeit the role of steroid hormones in the expression of ABC transporters in this organism remains to be investigated.

DISCUSSION

The *P. brasiliensis* transcriptome described here is represented by 6,022 EST clusters that may cover about 80% of the

fungal total genome, whose gene number has been estimated to be ~8,000 genes.³ This number greatly exceeds the previous EST studies in this fungus (8, 49). The analysis compares the two fungal cell types as well as their metabolic behavior. The results obtained probably reflect the adaptations associated with the mycelium (soil) and yeast (human host) environments. Most importantly, they provide new insights with respect to

³ G. San-Blas, personal communication.

signal transduction pathways, virulence genes, and drug targets for this pathogen.

The transcription profile of the mycelium infective phase suggests the shunting of pyruvate into aerobic metabolism, since the expression of the ESTs encoding enzymes of the trichloroacetic acid cycle are up-regulated in this fungal phase. In contrast, the yeast transcription profile evidenced the deviation of pyruvate from the glycolytic pathway into anaerobic metabolism; this observation is consistent with a lower oxygen level in infected tissues. Its putative ability to produce ethanol suggests a potential anaerobic pathway for *P. brasiliensis*, which is dependent on the metabolic state of the cell. It seems that the main regulatory effector on the shunting of the end product of glycolysis into aerobic or anaerobic metabolism is temperature; therefore, it can be hypothesized that this physical factor is the central trigger of all of these molecular events, since it was the only parameter changed in the *in vitro* cultivation of yeast and mycelium of *P. brasiliensis*. Experiments are currently being carried out in order to confirm the *in vivo* expression profile of the differentially expressed genes in macrophages and human pulmonary epithelial cells infected by *P. brasiliensis*.

Since *P. brasiliensis* is a medical problem in Latin America, the prediction of new drug targets from sequence information is of great importance. Chitin deacetylase, which is absent in humans and highly expressed in the parasitic yeast, could be a specific drug target for PCM therapy if it is shown to play a key role in the fungal metabolism during human infection. Functional analysis of the *P. brasiliensis* genes described in this work will lead to important information on cellular differentiation, pathogenicity, and/or virulence. These issues can only be addressed when molecular tools are developed for this organism. In conclusion, the knowledge of the transcribed sequences of *P. brasiliensis* will most likely facilitate the development of new therapeutics to PCM and other medically relevant mycosis.

Acknowledgments—We are grateful to Hugo Costa Paes and Robert Miller for English text revision.

REFERENCES

1. Franco, M. (1987) *J. Med. Vet. Mycol.* **25**, 5–18
2. Restrepo, A., McEwen, J. G. & Castaneda, E. (2001) *Med. Mycol.* **39**, 233–241
3. de Almeida, S. M., Queiroz-Telles, F., Teive, H. A., Ribeiro, C. E. & Werneck, L. C. (2004) *J. Infect.* **48**, 193–198
4. San-Blas, G., Nino-Vega, G. & Iturriaga, T. (2002) *Med. Mycol.* **40**, 225–242
5. San-Blas, G. & Nino-Vega, G. (2001) in *Fungal Pathogenesis: Principles and Clinical Applications*, pp. 205–226, Marcel Dekker, New York
6. Salazar, M. E., Restrepo, A. & Stevens, D. A. (1988) *Infect. Immun.* **56**, 711–713
7. Cano, M. I., Cispalino, P. S., Galindo, I., Ramirez, J. L., Mortara, R. A. & da Silveira, J. F. (1998) *J. Clin. Microbiol.* **36**, 742–747
8. Felipe, M. S., Andrade, R. V., Petrofeza, S. S., Maranhão, A. Q., Torres, F. A., Albuquerque, P., Arraes, F. B., Arruda, M., Azevedo, M. O., Baptista, A. J., Bataus, L. A., Borges, C. L., Campos, E. G., Cruz, M. R., Daher, B. S., Dantas, A., Ferreira, M. A., Ghil, G. V., Jesuino, R. S., Kyaw, C. M., Leitao, L., Martins, C. R., Moraes, L. M., Neves, E. O., Nicola, A. M. Alves, E. S., Parente, J. A., Pereira, M., Pocas-Fonseca, M. J., Resende, R., Ribeiro, B. M., Saldanha, R. R., Santos, S. C., Silva-Pereira, I., Silva, M. A., Silveira, E., Simoes, I. C., Soares, R. B., Souza, D. P., De-Souza, M. T., Andrade, E. V., Xavier, M. A., Veiga, H. P., Venancio, E. J., Carvalho, M. J., Oliveira, A. G., Inoue, M. K., Almeida, N. F., Walter, M. E., Soares, C. M. & Brigido, M. M. (2003) *Yeast* **20**, 263–271
9. Huang, X. & Madan, A. (1999) *Genome Res.* **9**, 868–877
10. Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D. J. (1997) *Nucleic Acids Res.* **25**, 3389–3402
11. Pearson, W. R. & Lipman, D. J. (1988) *Proc. Natl. Acad. Sci. U. S. A.* **85**, 2444–2448
12. Apweiler, R., Biswas, M., Fleischmann, W., Kanapin, A., Karavidopoulou, Y., Kersey, P., Kriventseva, E. V., Mittard, V., Mulder, N., Phan, I. & Zdobnov, E. (2001) *Nucleic Acids Res.* **29**, 44–48
13. Kanehisa, M. & Goto, S. (2000) *Nucleic Acids Res.* **28**, 27–30
14. Audic, S. & Claverie, J. M. (1997) *Genome Res.* **7**, 986–995
15. Quackenbush, J. (2002) *Nat. Genet.* **32**, (suppl.) 496–501
16. Tusher, V. G., Tibshirani, R. & Chu, G. (2001) *Proc. Natl. Acad. Sci. U. S. A.* **98**, 5116–5121
17. Venancio, E. J., Kyaw, C. M., Mello, C. V., Silva, S. P., Soares, C. M., Felipe, M. S. & Silva-Pereira, I. (2002) *Med. Mycol.* **40**, 45–51
18. Albuquerque, P., Kyaw, C. M., Saldanha, R. R., Brigido, M. M., Felipe, M. S. & Silva-Pereira, I. (2004) *Fungal Genet. Biol.* **41**, 510–520
19. Cunha, A. F., Sousa, M. V., Silva, S. P., Jesuino, R. S., Soares, C. M. & Felipe, M. S. (1999) *Med. Mycol.* **37**, 115–121
20. Marques, E. R., Ferreira, M. E., Drummond, R. D., Felix, J. M., Menossi, M., Savoldi, M., Travassos, L. R., Puccia, R., Batista, W. L., Carvalho, K. C., Goldman, M. H. & Goldman, G. H. (2004) *Mol. Genet. Genomics* **271**, 667–677
21. Loose, D. S., Stover, E. P., Restrepo, A., Stevens, D. A. & Feldman, D. (1983) *Proc. Natl. Acad. Sci. U. S. A.* **80**, 7659–7663
22. Madani, N. D., Malloy, P. J., Rodriguez-Pombo, P., Krishnan, A. V. & Feldman, D. (1994) *Proc. Natl. Acad. Sci. U. S. A.* **91**, 922–926
23. Pronk, J. T., Yde Steensma, H. & Van Dijken, J. P. (1996) *Yeast* **12**, 1607–1633
24. Restrepo, A., de Bedout, C., Cano, L. E., Arango, M. D. & Bedoya, V. (1981) *Sabouraudia* **19**, 295–300
25. Izacc, S. M., Gomez, F. J., Jesuino, R. S., Fonseca, C. A., Felipe, M. S., Deepe, G. S. & Soares, C. M. (2001) *Med. Mycol.* **39**, 445–455
26. da Silva, S. P., Borges-Walmsley, M. I., Pereira, I. S., Soares, C. M., Walmsley, A. R. & Felipe, M. S. (1999) *Mol. Microbiol.* **31**, 1039–1050
27. Moradas-Ferreira, P. & Costa, V. (2000) *Redox. Rep.* **5**, 277–285
28. Lengeler, K. B., Davidson, R. C., D'Souza, C., Harashima, T., Shen, W. C., Wang, P., Pan, X., Waugh, M. & Heitman, J. (2000) *Microbiol. Mol. Biol. Rev.* **64**, 746–785
29. Hohmann, S. (2002) *Int. Rev. Cytol.* **215**, 149–187
30. Sonneborn, A., Bockmuhl, D. P., Gerads, M., Kurpanek, K., Sanglard, D. & Ernst, J. F. (2000) *Mol. Microbiol.* **35**, 386–396
31. Paris, S. & Duran, S. (1985) *Mycopathologia* **92**, 115–120
32. Kraus, P. R. & Heitman, J. (2003) *Biochem. Biophys. Res. Commun.* **311**, 1151–1157
33. Mosch, H. U., Kubler, E., Krappmann, S., Fink, G. R. & Braus, G. H. (1999) *Mol. Biol. Cell* **10**, 1325–1335
34. Falkow, S. (2004) *Nat. Rev. Microbiol.* **2**, 67–72
35. Xu, J. R. (2000) *Fungal Genet. Biol.* **31**, 137–152
36. Bruckmann, A., Kunkel, W., Hartl, A., Wetzker, R. & Eck, R. (2000) *Microbiology* **146**, 2755–2764
37. Wills, E. A., Redinbo, M. R., Perfect, J. R. & Del Poeta, M. (2000) *Emerg. Therap. Targets* **4**, 1–32
38. Lorenz, M. C. & Fink, G. R. (2001) *Nature* **412**, 83–86
39. Muhlschlegel, F. A. & Fonzi, W. A. (1997) *Mol. Cell. Biol.* **17**, 5960–5967
40. Nathan, C. & Shiloh, M. U. (2000) *Proc. Natl. Acad. Sci. U. S. A.* **97**, 8841–8848
41. Moreira, S. F., Bailao, A. M., Barbosa, M. S., Jesuino, R. S., Felipe, M. S., Pereira, M. & de Almeida Soares, C. M. (2004) *Yeast* **21**, 173–182
42. Cox, G. M., Mukherjee, J., Cole, G. T., Casadevall, A. & Perfect, J. R. (2000) *Infect. Immun.* **68**, 443–448
43. Onyewu, C., Blankenship, J. R., Del Poeta, M. & Heitman, J. (2003) *Agents Chemother.* **47**, 956–964
44. Dickson, R. C. & Lester, R. L. (1999) *Biochim. Biophys. Acta* **1426**, 347–357
45. Kovalchuk, O. & Chakrabarty, K. (1994) *Eur. J. Biochem.* **226**, 133–140
46. Perea, S. & Patterson, T. F. (2002) *Clin. Infect. Dis.* **35**, 1073–1080
47. Gray, C. H., Borges-Walmsley, M. I., Evans, G. J. & Walmsley, A. R. (2003) *Yeast* **20**, 865–880
48. Krishnamurthy, S., Gupta, V., Prasad, R., Panwar, S. L. & Prasad, R. (1998) *FEMS Microbiol. Lett.* **160**, 191–197
49. Goldman, G. H., dos Reis Marques, E., Duarte Ribeiro, D. C., de Souza Bernardes, L. A., Quiapin, A. C., Vitorelli, P. M., Savoldi, M., Semighini, C. P., de Oliveira, R. C., Nunes, L. R., Travassos, L. R., Puccia, R., Batista, W. L., Ferreira, L. E., Moreira, J. C., Bogossian, A. P., Tekaiia, F., Nobrega, M. P., Nobrega, F. G. & Goldman, M. H. (2003) *Eukaryot. Cell* **2**, 34–48

Transcriptional Profiles of the Human Pathogenic Fungus *Paracoccidioides brasiliensis* in Mycelium and Yeast Cells

Maria Sueli S. Felipe, Rosângela V. Andrade, Fabrício B. M. Arraes, André M. Nicola, Andréa Q. Maranhão, Fernando A. G. Torres, Ildinete Silva-Pereira, Márcio J. Poças-Fonseca, Élide G. Campos, Lídia M. P. Moraes, Patrícia A. Andrade, Aldo H. F. P. Tavares, Simoneide S. Silva, Cynthia M. Kyaw, Diorge P. Souza, PbGenome Network, Maristela Pereira, Rosália S. A. Jesuino, Edmar V. Andrade, Juliana A. Parente, Gisele S. Oliveira, Mônica S. Barbosa, Natália F. Martins, Ana L. Fachin, Renato S. Cardoso, Geraldo A. S. Passos, Nalvo F. Almeida, Maria Emília M. T. Walter, Célia M. A. Soares, Maria José A. Carvalho and Marcelo M. Brígido

J. Biol. Chem. 2005, 280:24706-24714.

doi: 10.1074/jbc.M500625200 originally published online April 22, 2005

Access the most updated version of this article at doi: [10.1074/jbc.M500625200](https://doi.org/10.1074/jbc.M500625200)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

Supplemental material:

<http://www.jbc.org/content/suppl/2005/05/05/M500625200.DC1>

This article cites 46 references, 18 of which can be accessed free at <http://www.jbc.org/content/280/26/24706.full.html#ref-list-1>