



UNIVERSIDADE FEDERAL DE GOIÁS
PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO
INSTITUTO DE CIÊNCIAS BIOLÓGICAS
PROGRAMA DE PÓS-GRADUAÇÃO EM ECOLOGIA E EVOLUÇÃO



Leandro Maracahipes dos Santos

ESTRATÉGIAS ECOLÓGICAS DE PLANTAS EM
FLORESTAS ESTACIONAIS E SAVANAS DO CERRADO

Orientador: Dr. Marcus Vinicius Cianciaruso

Coorientador: Dr. Marcos Bergmann Carlucci

GOIÂNIA - GO

ABRIL – 2017

TERMO DE CIÊNCIA E DE AUTORIZAÇÃO PARA DISPONIBILIZAR AS TESES E DISSERTAÇÕES ELETRÔNICAS NA BIBLIOTECA DIGITAL DA UFG

Na qualidade de titular dos direitos de autor, autorizo a Universidade Federal de Goiás (UFG) a disponibilizar, gratuitamente, por meio da Biblioteca Digital de Teses e Dissertações (BDTD/UFG), regulamentada pela Resolução CEPEC nº 832/2007, sem ressarcimento dos direitos autorais, de acordo com a Lei nº 9610/98, o documento conforme permissões assinaladas abaixo, para fins de leitura, impressão e/ou *download*, a título de divulgação da produção científica brasileira, a partir desta data.

1. Identificação do material bibliográfico: ☐ Dissertação ☒ Tese

2. Identificação da Tese ou Dissertação


Nome completo do autor: Leandro Maracahipes dos Santos

Título do trabalho: Estratégias ecológicas de plantas em florestas estacionais e savanas do Cerrado

3. Informações de acesso ao documento:

Concorda com a liberação total do documento ☒ SIM ☐ NÃO¹

Havendo concordância com a disponibilização eletrônica, torna-se imprescindível o envio do(s) arquivo(s) em formato digital PDF da tese ou dissertação.


Assinatura do (a) autor (a)

Data: 19 / 05 / 2017

¹ Neste caso o documento será embargado por até um ano a partir da data de defesa. A extensão deste prazo suscita justificativa junto à coordenação do curso. Os dados do documento não serão disponibilizados durante o período de embargo.

UNIVERSIDADE FEDERAL DE GOIÁS
PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO
INSTITUTO DE CIÊNCIAS BIOLÓGICAS
PROGRAMA DE PÓS-GRADUAÇÃO EM ECOLOGIA E EVOLUÇÃO

Leandro Maracahipes dos Santos

**ESTRATÉGIAS ECOLÓGICAS DE PLANTAS EM
FLORESTAS ESTACIONAIS E SAVANAS DO CERRADO**

Orientador: Dr. Marcus Vinicius Cianciaruso

Coorientador: Dr. Marcos Bergmann Carlucci

Tese apresentada à Universidade Federal de Goiás, como parte das exigências do Programa de Pós-graduação em Ecologia e Evolução para obtenção do título de doutor em Ecologia e Evolução.

GOIÂNIA - GO

ABRIL – 2017

Ficha de identificação da obra elaborada pelo autor, através do
Programa de Geração Automática do Sistema de Bibliotecas da UFG.

Santos, Leandro Maracahipes dos
Estratégias ecológicas de plantas em florestas estacionais e
savanas do Cerrado [manuscrito] / Leandro Maracahipes dos Santos.
- 2017.
vii, 123 f.

Orientador: Prof. Dr. Marcus Vinicius Cianciaruso; co-orientador
Dr. Marcos Bergmann Carlucci.

Tese (Doutorado) - Universidade Federal de Goiás, Instituto de
Ciências Biológicas (ICB), Programa de Pós-Graduação em Ecologia e
Evolução, Goiânia, 2017.

Bibliografia. Anexos.

Inclui mapas, abreviaturas, gráfico, tabelas.

1. preferência de habitats. 2. plasticidade fenotípica. 3. gradiente
edáfico. 4. dano foliar em contexto de vizinhança. 5. distâncias
ecológicas e evolutivas. I. Cianciaruso, Marcus Vinicius, orient. II. Título.

CDU 574




**SERVIÇO PÚBLICO FEDERAL
UNIVERSIDADE FEDERAL DE GOIÁS - UFG
INSTITUTO DE CIÊNCIAS BIOLÓGICAS - ICB
PROGRAMA DE PÓS-GRADUAÇÃO EM ECOLOGIA E EVOLUÇÃO - PPGEE**


ATA DA SESSÃO PÚBLICA DE DEFESA DE TESE Nº 55

Aos treze dias do mês de abril de 2017 (13/04/2017), às oito horas (08h), no Auditório do ICB V, UFG, reuniram-se os componentes da banca examinadora: **Prof. Dr. Marcus Vinicius Cianciaruso, ICB-UFG; Prof. Dr. Paulo De Marco Júnior, ICB-UFG; Dra. Marcia Nunes Macedo, Woods Hole Research Center – Falmouth, EUA; Prof. Dr. Fabrício Barreto Teresa, UEG-Anápolis; Prof. Dr. José Roberto Rodrigues Pinto, UnB;** para, em sessão pública presidida pelo (a) primeiro(a) examinador(a) citado(a), procederem à avaliação da defesa de tese intitulada: **“Estratégias ecológicas de plantas em florestas estacionais e savanas do Cerrado”**, em nível de doutorado, área de concentração em Ecologia e Evolução, de autoria de **Leandro Maracahipes dos Santos**, discente do Programa de Pós-Graduação Ecologia e Evolução da Universidade Federal de Goiás. A sessão foi aberta pelo(a) presidente(a), que fez a apresentação formal dos membros da banca. A palavra, a seguir, foi concedida a(o) autor(a) da tese que, em cerca de 35 minutos, procedeu à apresentação de seu trabalho. Terminada a apresentação, cada membro da banca arguiu a(o) examinada(o), tendo-se adotado o sistema de diálogo sequencial. Terminada a fase de arguição, procedeu-se à avaliação da tese. Tendo-se em vista o que consta na Resolução nº 1127 de dezembro de 2012 do Conselho de Ensino, Pesquisa, Extensão e Cultura (CEPEC), que regulamenta o Programa de Pós-Graduação em Ecologia e Evolução, a tese foi APROVADA, considerando-se integralmente cumprido este requisito para fins de obtenção do título de Doutor(a) em Ecologia e Evolução pela Universidade Federal de Goiás. A conclusão do curso dar-se-á quando da entrega da versão definitiva da tese na secretaria do programa, com as devidas correções sugeridas pela banca examinadora, no prazo de trinta dias a contar da

data da defesa. Cumpridas as formalidades de pauta, às 12 h e 02 min., encerrou-se a sessão de defesa e, para constar, eu, Suely Ana Ribeiro, secretária executiva da Universidade Federal de Goiás - UFG, lavrei a presente ata que, após lida e aprovada, será assinada pelos membros da banca examinadora em três vias de igual teor.



Prof. Dr. Marcus Vinicius Cianciaruso
Presidente da banca
ICB-UFG



Prof. Dr. Paulo De Marco Júnior
ICB-UFG



Dra. Marcia Nunes Macedo
Woods Hole Research Center – Falmouth, EUA



Prof. Dr. Fabrício Barreto Teresa
UEG-Anápolis



Prof. Dr. José Roberto Rodrigues Pinto
UnB

*Aos meus pais, Luiz e Helena
pelo amparo e incentivo*

Agradecimentos

Agradeço à Universidade Federal de Goiás através do Programa de Pós-Graduação em Ecologia e Evolução pelo suporte estrutural e de aprendizado disponibilizado para a formação dos discentes.

Agradeço à Capes pela concessão de bolsa de estudo durante estes quatro anos de doutoramento. Bem como ao CNPq e Fapeg que financiaram o desenvolvimento deste projeto de pesquisa.

Ao prestativo orientador Marcus Cianciaruso pelo grande apoio e suporte para desenvolvimento deste trabalho, e principalmente pelas correções, sugestões, paciência e dedicação de tempo para discutir a direção de cada trabalho. Também, gostaria de agradecer a excelente co-orientação que obtive por parte do Marcos Carlucci, pelas sugestões e correções que certamente enriqueceram o trabalho.

Ao corpo docente do programa que direta ou indiretamente contribuíram para com a minha formação através de disciplinas cursadas, palestras, cursos e conversas específicas. Estendo meus agradecimentos também ao corpo discente que contribuíram bastante com minha formação durante o processo de doutoramento, por meio de palestras e defesas assistidas, e principalmente discussões individuais diversos temas, bem como temas relacionadas ao meu projeto de pesquisa. Especialmente aos colegas Fernando Sobral, Walter Araújo, Leonardo Bergamini, Advaldo Neto e Claudinei (San).

À todos os amigos e colegas que auxiliaram durante a realização das coletas de dados, com excelente trabalho e elevada contribuição para a construção da base de dados que resultou no desenvolvimento desta tese: Edmar Oliveira, Leonardo Maracahipes, Letícia Gomes, Livia Laureto, Danira Padilha, Mônica Forsthofer, Mariângela Abreu, Josias Santos e Fernando Ribeiro. Agradeço também aos taxonomistas Geraldo Franco, Osnir Aguiar, João Baitelo, Natália Ivanauskas e Renato Mello-Silva pela identificação de algumas espécies.

Agradeço à todas as pessoas do antigo (especialmente Priscila, Geisiane e Nathália) e do atual laboratório de comunidades, especialmente ao Fernando, Ana, Cibele, Daniel, Fábio, Fabiane, Francielle, Gabriela, Jaqueline, J. Hidasi, Joyce, Livia, Ingrid, Nubia, Paula, Paulo, Rhayane e Verônica, com os quais compartilhei muitos momentos de alegria, aprendizado, trabalho e especialmente amizade.

Aos queridos “Pragas”, vocês são demais e foram ótimos os momentos que passamos juntos durante boa parte deste doutorado. À “praga mor” Vanessa, Karina, Renata e Letícia, e aos brothers Paulinho, Thiago e L. Brasil.

Ao Leandro Juen pela indicação em 2011 para uma bolsa de pesquisa dentro de um projeto, que se tornou meu projeto de tese. Muito obrigado pela indicação, confiança e amizade adquirida.

Ao pessoal do “futebol de quinta”, que apesar do sugestivo nome jogam um futebol de primeira qualidade. Sentirei muita falta destes dias, e claro vocês terão que conseguir um outro “camisa 10” para o time.

À minha família, meus pais, Luiz Maracahipes e Helena Maria, e irmãos e cunhada, Leonardo, Eliabe e Ângela. E aos sobrinhos, Daniel e Davi, que com os quais convivi pouco em razão da distância, mas que sempre se fizeram presentes em minha vida. Obrigado família, e Parabéns, meus pais por sempre incentivarem e investirem na educação de todos os seus filhos.

Sumário

Resumo	1
Introdução geral	3
 Capítulo I	
Plant ecological strategies in seasonal forests and savannas species	16
 Capítulo II	
Edaphic properties drive functional trait patterns in savannas and seasonal forests plant communities of the Cerrado	53
 Capítulo III	
Insect herbivore damage is not related with host plant ecological and evolutionary distances	95
 Conclusão geral	 122

Resumo – A adoção de diferentes estratégias ecológicas é um fator importante para determinar o estabelecimento e a persistência de espécies em comunidades locais. De maneira geral, o cerrado é caracterizado por uma alta frequência de fogo e solos pobres em nutrientes. Geralmente em condições de baixa fertilidade e alta frequência de fogo as espécies filtradas tendem a possuir características que representam adaptações a estes estresses ambientais. Considerando que as espécies de Cerrado se desenvolvem sob a atuação destes filtros ambientais, nosso objetivo foi avaliar como a adoção de diferentes estratégias ecológicas podem determinar a performance dos atributos funcionais, a estrutura das comunidades e a relação entre uma planta focal e sua vizinhança. Neste trabalho de tese, que está dividido em três capítulos, nós utilizamos três diferentes escalas para avaliar como estratégias ecológicas das espécies podem determinar seu desempenho e estabelecimento em comunidades locais. No primeiro capítulo que está baseado em uma escala de habitat, nós avaliamos como as estratégias ecológicas de espécies generalistas e especialistas de floresta estacional e cerrado sentido restrito são fundamentais para o estabelecimento e a persistência das espécies nestes habitats com diferenças marcantes em relação à frequência de fogo e disponibilidade de nutrientes. Neste capítulo, nós discutimos que as diferentes estratégias adotadas pelas espécies estão de acordo com os fatores limitantes da ocorrência de espécies em cada um destes ambientes. No segundo capítulo, que está baseado em escala de comunidades, nós buscamos compreender como os gradientes ambientais podem determinar diferentes estratégias ecológicas relacionadas aos atributos funcionais e a densidade de espécies. Nós demonstramos que as mudanças nos valores de atributos e densidade de espécies foram mais claras no gradiente de fertilidade do que toxicidade, e que comunidades de floresta estacional foram mais sensíveis a mudanças do que comunidades de cerrado sentido restrito em ambos os gradientes. Nós observamos também que espécies com

atributos conservativos foram associados à solos pobres e espécies com atributos aquisitivos associado à solos mais férteis. Já no terceiro capítulo, que foi desenvolvido na escala de indivíduo, nós discutimos se as características e relação filogenética das plantas vizinhas influenciam o dano foliar em árvores e arbustos do cerrado. Neste capítulo, demonstramos que a distância ecológica e evolutiva entre plantas individuais e as plantas vizinhas não determina o nível de consumo foliar por herbívoros. Nós discutimos que a dominância de herbívoros generalistas, a co-evolução entre plantas e herbívoros especialistas, e o consumo preferencial de folhas jovens podem ser mais importante para determinar o nível de dano foliar do que o contexto de vizinhança em que uma dada planta está inserida.

Palavras-chave: preferência de habitats, plasticidade fenotípica, gradiente edáfico, estratégia aquisitiva e conservativa, dano foliar, distâncias ecológicas e evolutivas, contexto de vizinhança

Introdução Geral

A diversidade do reino vegetal vem fascinando o homem há muito tempo, desde os celebres naturalistas aos renomados ecólogos contemporâneos, proponentes de teorias que buscam compreender quais são os mecanismos estruturadores da riqueza e composição de espécies. A teoria do nicho e todos os seus princípios tem sido aquela mais frequentemente testada ao longo do desenvolvimento da ecologia, essencialmente no que diz respeito às interações competitivas entre espécies e à atuação de filtros ambientais (tais como fogo em savanas do Cerrado). Assim, em função da atuação de filtros ambientes, a adoção de diferentes estratégias ecológicas é um fator importante para determinar o estabelecimento e a persistência de espécies, bem como os valores dos atributos funcionais das espécies capazes de se estabelecer em uma comunidade local.

Breve histórico

Comunidade ecológica é definida como um grupo de espécies que co-ocorre em um dado tempo e lugar (McGill *et al.* 2006). Entender os padrões e os mecanismos reguladores da composição e diversidade das comunidades naturais é um dos principais objetivos da ciência ecológica. Já nos séculos XVIII e XIX, os trabalhos dos naturalistas Alexander von Humboldt e Eugenius Warming buscaram compreender a relação entre plantas, animais e clima, bem como inferir os padrões de diversidade de cada localidade (Hawkins 2001). Os trabalhos de Warming e Humboldt influenciaram notáveis naturalistas do passado, tais como Karl Philipp von Martius, Alfred Wallace e Charles Darwin. Neste sentido, em “A origem das espécies” Darwin postula que indivíduos com alto grau de parentesco tendem a apresentar fortes relações competitivas, introduzindo o

conceito de “competição ecológica” como um dos principais mecanismos reguladores da diversidade e composição de comunidades naturais (Mayfield & Levine 2010).

Na década de 20 e 30, Vito Volterra, Alfred Lotka e Georgii Gause avançaram no entendimento dos mecanismos reguladores da riqueza e composição de espécies ao propor o conceito de nicho ecológico. Estes autores postularam o “princípio da exclusão competitiva” ao afirmarem que duas espécies competindo pelo mesmo recurso não podem coexistir ao longo do tempo em uma mesma comunidade (Volterra 1926; Lotka 1932; Gause 1934). Em meados do século XX, George Hutchinson redefiniu nicho ecológico como um espaço multidimensional composto pelos requerimentos bióticos e abióticos necessários à sobrevivência e reprodução das espécies (Hutchinson 1957). Na década de 60, Robert MacArthur e Edward O. Wilson formularam a “teoria do equilíbrio da biogeografia de ilhas”, a qual exerceu um papel primordial no desenvolvimento do “princípio da partição de nicho”. Na mesma década, MacArthur e Levins formularam a hipótese da similaridade limitante (MacArthur & Levins 1967), que prevê um limite máximo de semelhança morfológica para duas espécies coexistirem numa mesma comunidade. Em hipótese, a coexistência de duas ou mais espécies com diferentes características morfológicas seria possibilitada pela exploração de recursos distintos, resultando numa baixa sobreposição de nicho entre essas espécies.

Atualmente, os cientistas buscam compreender como os diferentes mecanismos reguladores da diversidade e composição de espécies atuam nas diferentes escalas espaciais. Neste sentido, se destacam os trabalhos de David Tilman e Peter Chesson (Tilman 1982, 1990, Chesson 2000), os quais buscam entender como atuam os principais mecanismos reguladores da composição e diversidade de espécies. Mais recentemente, uma nova abordagem baseada na “teoria de biogeografia de ilhas”, denominada “Teoria Neutra da Biodiversidade”, tem sido utilizada para explicar os

padrões de diversidade e composição de espécies nos mais variados grupos de organismos (Hubbell 2001). A teoria neutra assume que as espécies dentro de uma comunidade são ecologicamente equivalentes e que as extinções são balanceadas pelas especiações, e por processos estocásticos de nascimento e imigração.

Teoria de Nicho

A teoria de nicho teve suas bases lançadas por Eugenius Warming ao observar que as plantas possuíam diferentes habilidades fisiológicas e que algumas espécies eram capazes de se estabelecer em determinados locais e incapazes de obter o mesmo sucesso em outros locais (Cavender-Bares *et al.* 2009). Partindo destas observações, Grinnell e Elton foram os primeiros ecólogos a desenvolverem o conceito de nicho ecológico de espécie (Grinnell 1917; Elton 1927). Posteriormente, Hutchinson redefiniu o conceito de nicho (Hutchinson 1957), assumindo que espécies com elevada similaridade ecológica não poderiam coexistir, assim criou-se um paradigma onde processos evolutivos não possuíam qualquer influência na estruturação das comunidades (Cavender-Bares *et al.* 2009). No entanto, atualmente diversos estudos vêm demonstrando a relevância da história evolutiva e das interações interespecíficas como processos fundamentais para compreender a coexistência das espécies (Webb *et al.* 2002; Cahill *et al.* 2008; Cavender-Bares *et al.* 2009; Vamosi *et al.* 2009; Mayfield & Levine 2010).

A teoria do nicho Hutchinsoniano basicamente é constituída de um conjunto de condições (abióticas - também chamado de nicho Grinnelliano) e de recursos (bióticos - nicho Eltoniano) que determinam a capacidade de uma espécie em se estabelecer e manter populações viáveis ao longo do tempo (Hutchinson 1957; Soberón 2007). A premissa básica desta teoria encontra-se fundamentada no Princípio da Exclusão

Competitiva proposto por Gause (Gause 1934). Dentro da teoria do nicho, a atuação de filtros ambientais também pode ser um importante processo regulador do nicho das espécies (Gómez *et al.* 2010). O filtro ambiental determina quais espécies são aptas a se estabelecer em cada local, e é provável que atue sobre escalas espaciais maiores (escala regional), o que resultaria na seleção de espécies mais semelhantes em atributos (Gómez *et al.* 2010; Sobral & Cianciaruso 2012). Enquanto que, após a seleção de determinado conjunto de espécies pelo filtro ambiental, em escala local a similaridade limitante entre as espécies é quem irá atuar e determinar quais espécies deverão coexistir (Gómez *et al.* 2010; Sobral & Cianciaruso 2012).

Entretanto, dentro da teoria de nicho outros processos importantes são a sobreposição e a diferenciação dos nichos (Gilbert 2012). Quando espécies possuem elevado grau de sobreposição de nicho (ou seja, alto compartilhamento de consumo de recursos) isso resulta em fortes interações competitivas que muitas vezes podem levar a exclusão competitiva, ou forçar as espécies a explorarem diferentes recursos (diferenciação de nicho) (Webb *et al.* 2002). A diferenciação de nicho pode ainda levar ao deslocamento de caracteres, caracterizado por uma alta similaridade morfológica quando duas espécies são distribuídas alopatricamente, mas há uma diferenciação em uma ou mais dessas características quando ocorrem simpatricamente (Connell 1980; Schmidt *et al.* 2000). Outra possibilidade é a divisão do recurso disponível entre os competidores (partição de nicho), que consiste em cada competidor ocupar uma porção do recurso disponível, evitando a competição e consequente exclusão de um dos competidores (Webb *et al.* 2002).

Estratégias ecológicas de plantas

Em ecologia funcional, diversos atributos funcionais têm sido utilizados para compreender a estruturação e a funcionalidade das comunidades naturais (atributos foliares, da madeira e regenerativos). Dentre estes, o sistema LHS (*leaf-height-seed*; folha-altura-semente) descreve o nicho funcional das plantas, usando estes três eixos fundamentais e independentes, resumindo todas as principais dimensões de variação na estrutura e funcionamento (Westoby 1998). A representação deste sistema, que consiste em um modelo de classificação tridimensional, é importante por representar demandas conflitantes fundamentais controlando as estratégias vegetais e pelo fato de que cada uma dessas características se correlaciona com outros atributos relevantes (Westoby 1998; Laughlin *et al.* 2010). A área foliar específica é o principal atributo envolvendo o espectro econômico foliar (Wright *et al.* 2004) e representa a demanda conflitante entre aquisição e a estratégia no uso de recurso pelas plantas. A altura máxima da planta indica a habilidade competitiva da planta por luz e, portanto, sua estratégia de assimilação de carbono (Westoby *et al.* 2002). Enquanto que a massa da semente indica a estratégia de regeneração das espécies por meio da demanda conflitante entre massa da semente, habilidade de dispersão e estabelecimento de plântulas (Westoby *et al.* 2002). O sistema LHS vem sendo testado em diversos ambientes e demonstrando ser capaz de prever a funcionalidade das comunidades (Lavergne, Garnier & Debussche 2003; Carly, Marcelo & Jaime 2009).

Em ambientes florestais, em que o acesso à luz é um fator limitante, o investimento de árvores em crescimento vertical é vantajoso, pois árvores que apresentam alturas maiores do que seus vizinhos possuem maior vantagem competitiva devido ao melhor acesso à luz. Ainda, a altura em que as flores e sementes são produzidas também pode influenciar o sucesso reprodutivo dos indivíduos, bem como a

dispersão de sementes (Garnier & Navas 2012), uma vez que a altura da árvore possui uma relação positiva com a distância em que as sementes são dispersas. Por outro lado, plantas que produzem poucas sementes grandes são caracterizadas como espécies com alta habilidade competitiva, enquanto que espécies que produzem elevado número de pequenas sementes são vistas como espécies pioneiras devido à sua alta capacidade de dispersão (Garnier & Navas 2012). Outro importante mecanismo utilizado por plantas é o sistema de rebrota. Diversas plantas do Cerrado apresentam esta característica de resiliência, que permite com que o indivíduo possa rebrotar após um distúrbio (como fogo) a partir de estruturas basais ou subterrâneas (Medeiros & Miranda 2005).

Fogo, consumo foliar e propriedades do solo no Cerrado

O Cerrado é composto por um complexo mosaico vegetacional, com a ocorrência de vários tipos de vegetação (Eiten 1972), variando de savanas abertas a florestas de galeria, florestas secas e semidecíduais. Muitas das espécies são capazes de se estabelecer em diferentes fisionomias (Ratnam *et al.* 2011; Hoffmann *et al.* 2012). No entanto, estratégias relacionadas principalmente a proteção contra o fogo (grande espessura da casca) podem ser cruciais para determinar o estabelecimento e permanência de espécies nos diferentes ambientes savânicos do Cerrado. Além disso, estratégias relacionadas a proteção contra herbívoros (folhas grossas, espinhos e tricomas) e perda de água (folhas grossas, e fechamento de estômatos e esporos) (Silva & Batalha 2011; Dantas, Batalha & Pausas 2013a).

Uma vez que os processos de dispersão e eventos estocásticos determinam o banco regional de espécies, este é composto pelo conjunto de espécies que possuem agentes dispersores que as tornem capazes de chegar a uma área (Lortie *et al.* 2004), possibilitando que um conjunto de espécies melhor adaptado as condições locais seja

selecionado a partir do banco regional de espécies. Inicialmente, em maior escala poderá ser filtrado um conjunto de espécies com atributos funcionais mais similares (por exemplo, pelo clima), e posteriormente por fatores ambientais em menores escalas (por exemplo, características edáficas) e, por último as interações bióticas entre as espécies (tais como relação entre planta-planta e planta-herbívoros) irão determinar a abundância relativa e quais espécies poderão estabelecer e persistir em cada comunidade local (Cain, Milligran & Strand 2000; Dufour *et al.* 2006; Lessard *et al.* 2012; Bello *et al.* 2013).

O fogo é um dos principais fatores estruturadores de comunidades vegetais no cerrado (Hoffmann *et al.* 2012; Dantas *et al.* 2013b, 2016). Os focos de incêndio no cerrado ocorrem com elevada frequência e intensidade, principalmente em localidades em que ocorre grande acúmulo de biomassa durante alguns anos sem fogo. No Cerrado, áreas de savanas possuem grande abundância de gramíneas, com arbustos e árvores coexistindo sobre a camada de herbácea. No momento da passagem do fogo as gramíneas atuam como uma grande fonte de combustível, promovendo um maior avanço do fogo em áreas com maior quantidade (Dantas *et al.* 2013a). Enquanto que, em áreas florestais a maior umidade relativa e a baixa quantidade de gramíneas dificultam a propagação do fogo nestes ambientes (Hoffmann *et al.* 2012).

O fogo atua como um filtro selecionando espécies com características que confirmam resistência contra os efeitos das queimadas ou que possuem estruturas de reserva que permita rebrotar após a passagem do fogo (Dantas *et al.* 2013a; Pausas, Keeley & Schwilk 2016). As espécies evoluíram com a presença constante de fogo, sendo assim diversas adaptações de resistência e resiliência ao fogo foram desenvolvidas ao longo do tempo evolutivo (Simon *et al.* 2009; Pausas *et al.* 2016). A espessura da casca é um dos principais atributos de resistência ao fogo em plantas

(Rosell *et al.* 2014). Hoffmann *et al.* (2012) propuseram um limiar de que espécies com espessura de casca superior à 5,9 mm tem alta probabilidade resistir a queimadas de baixa intensidade. Muitas das espécies exclusivas de savanas possuem espessura de casca igual ou superior a este limiar, no entanto espécies com habito florestal que ocorrem em áreas de savanas não possuem tais características, e muitas vezes são excluídas das comunidades locais após a passagem do fogo.

Considerando que habitats de florestas estacionais e savanas são contrastantes, e que ocorrem lado a lado em paisagens de Cerrado. Isto torna possível encontrar espécies com ocorrência em ambos habitats ou com ocorrência restrita à somente um destes habitats (floresta estacional ou savana). Sendo assim, avaliar como diferentes estratégias ecológicas de espécies generalistas e especialistas de florestas estacional e cerrado sentido restrito é fundamental para compreender o estabelecimento e a persistência de espécies nestes habitats com diferenças marcantes em relação à frequência de fogo e disponibilidade de nutrientes. Nesse primeiro capítulo, nós discutimos como as diferentes estratégias adotadas pelas espécies estão de acordo com os fatores limitantes da ocorrência de espécies em cada um destes ambientes.

De maneira geral, os solos do cerrado são pobres em nutrientes e bem drenados (Furley & Ratter 1988). Desta forma, para que a coexistência de espécies torne-se possível, frequentemente as espécies adotam diferentes estratégias relacionadas ao crescimento e reprodução (Oliveras & Malhi 2016), que podem resultar em estratégias de rápida aquisição ou conservação de recursos. As propriedades edáficas influenciam fortemente a estrutura de comunidade de plantas. Assim, visto que mudanças ao longo de gradientes ambientais são capazes de alterar a composição funcional das comunidades, nós buscamos compreender neste segundo capítulo como o gradiente de fertilidade e toxicidade do solo dentro de cada um dos habitats (floresta estacional e

savana) podem determinar a performance dos atributos funcionais e densidade de espécies. Logo, nós fomos hábeis em discutir se espécies com atributos conservativos foram associados à solos pobres e espécies com atributos aquisitivos associado à solos mais férteis.

Estas condições de baixa fertilidade dos em solos savanas e a alta frequência de fogo, fazem com que as espécies filtradas tenham características adaptativas à estresse ambiental, que também estão relacionadas a proteção contra o ataque por herbívoros. As plantas de savanas desenvolveram diversas estratégias que inibe ou acaba por reduzir o efeito dos herbívoros sobre as plantas (Silva & Batalha 2011). Entretanto, o nível de consumo foliar por herbívoros varia muitas entre as plantas coexistentes em um dado local. Diversos fatores podem explicar esta variação, aqui neste Capítulo 3 nós buscamos compreender se as características das plantas vizinhas poderiam determinar a taxa de consumo foliar de um indivíduo focal. Desta forma, nós buscamos entender como a distância ecológica e evolutiva entre uma planta individual e suas plantas vizinhas podem mediar a nível de consumo foliar.

Em suma, nesta tese nós usamos três diferentes escalas para avaliar como estratégias ecológicas de plantas podem determinar a performance dos atributos funcionais, a estrutura das comunidades e a relação entre uma planta focal e sua vizinhança. O Capítulo 1 está baseado em uma escala de habitat e busca compreender como a preferência de habitat pode determinar a performance dos atributos funcionais em diferentes habitats. Já o Capítulo 2 foi analisado em uma escala de comunidades e busca entender como gradientes ambientais podem determinar diferentes estratégias ecológicas relacionadas aos atributos funcionais e a densidade de espécies. E por fim, o Capítulo 3 foi desenvolvido em uma escala de indivíduo e nós buscamos compreender

como as características e relação filogenética das plantas vizinhas poderiam influenciar na taxa de dano foliar de uma planta focal.

Referências

- Bello, F. de, Lavorel, S., Lavergne, S., Albert, C.H., Boulangeat, I., Mazel, F. & Thuiller, W. (2013) Hierarchical effects of environmental filters on the functional structure of plant communities: a case study in the French Alps. *Ecography*, **36**, 393–402.
- Cahill, J.F., Kembel, S.W., Lamb, E.G. & Keddy, P.A. (2008) Does phylogenetic relatedness influence the strength of competition among vascular plants? *Perspectives in Plant Ecology, Evolution and Systematics*, **10**, 41–50.
- Cain, M.L., Milligran, B.G. & Strand, A.E. (2000) Long-distance seed dispersal in plant populations. *American Journal of Botany*, **87**, 1217–1227.
- Carly, G., Marcelo, S. & Jaime, K. (2009) A community-level test of the leaf-height-seed ecology strategy scheme in relation to grazing conditions. *Journal of Vegetation Science*, **20**, 392–402.
- Cavender-Bares, J., Kozak, K.H., Fine, P.V.A. & Kembel, S.W. (2009) The merging of community ecology and phylogenetic biology. *Ecology Letters*, **12**, 693–715.
- Chesson, P. (2000) Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics*, **31**, 343–366.
- Connell, J.H. (1980) Diversity and the Coevolution of Competitors, or the Ghost of Competition Past. *Oikos*, **35**, 131–138.
- Dantas, V.L., Batalha, M.A. & Pausas, J.G. (2013a) Fire drives functional thresholds on the savanna–forest transition. *Ecology*, **94**, 2454–2463.
- Dantas, V. de L., Hirota, M., Oliveira, R.S. & Pausas, J.G. (2016) Disturbance maintains alternative biome states (ed M Rejmanek). *Ecology Letters*, **19**, 12–19.
- Dantas, V. de L., Pausas, J.G., Batalha, M.A., Loiola, P. de P. & Cianciaruso, M.V. (2013b) The role of fire in structuring trait variability in Neotropical savannas. *Oecologia*, **171**, 487–494.
- Dufour, A., Gadallah, F., Wagner, H.H., Guisan, A. & Buttler, A. (2006) Plant species richness and environmental heterogeneity in a mountain landscape: Effects of variability and spatial configuration. *Ecography*, **29**, 573–584.

- Eiten, G. (1972) The cerrado vegetation of Brazil. *Botanical Reviews*, **38**, 201–341.
- Elton, C. (1927) *Animal Ecology* (ed JS Huxley). Macmillan Co., New York.
- Furley, P.A. & Ratter, J.A. (1988) Soil resources and plant communities of the central Brazilian Cerrado and their development. *Journal of Biogeography*, **15**, 97–108.
- Garnier, E. & Navas, M.L. (2012) A trait-based approach to comparative functional plant ecology: Concepts, methods and applications for agroecology. A review. *Agronomy for Sustainable Development*, **32**, 365–399.
- Gause, G.F. (1934) *The Struggle for Existence*. Willians and Wilkins, Baltimore.
- Gilbert, B. (2012) Joint consequences of dispersal and niche overlap on local diversity and resource use. *Journal of Ecology*, **100**, 287–296.
- Gómez, J.P., Bravo, G.A., Brumfield, R.T., Tello, J.G. & Cadena, C.D. (2010) A phylogenetic approach to disentangling the role of competition and habitat filtering in community assembly of Neotropical forest birds. *Journal of Animal Ecology*, **79**, 1181–1192.
- Grinnell, J. (1917) The niche-relationships of the California thrasher. *Auk*, **34**, 427–433.
- Hawkins, B.A. (2001) Ecology's oldest pattern? *Trends in Ecology & Evolution*, **16**, 470.
- Hoffmann, W. a, Geiger, E.L., Gotsch, S.G., Rossatto, D.R., Silva, L.C.R., Lau, O.L., Haridasan, M. & Franco, A.C. (2012) Ecological thresholds at the savanna-forest boundary: how plant traits, resources and fire govern the distribution of tropical biomes (ed F Lloret). *Ecology Letters*, **15**, 759–768.
- Hubbell, S.P. (2001) *A Unified Neutral Theory of Biodiversity and Biogeography*. Princeton University Press, Princeton.
- Hutchinson, G.E. (1957) Concluding Remarks. *Cold Spring Harbor Symposia on Quantitative Biology*, **22**, 415–427.
- Laughlin, D.C., Leppert, J.J., Moore, M.M. & Sieg, C.H. (2010) A multi-trait test of the leaf-height-seed plant strategy scheme with 133 species from a pine forest flora. *Functional Ecology*, **24**, 493–501.
- Lavergne, S., Garnier, E. & Debussche, M. (2003) Do rock endemic and widespread plant species differ under the Leaf-Height-Seed plant ecology strategy scheme? *Ecology Letters*, **6**, 398–404.
- Lessard, J.P., Belmaker, J., Myers, J.A., Chase, J.M. & Rahbek, C. (2012) Inferring local ecological processes amid species pool influences. *Trends in Ecology and*

- Evolution*, **27**, 600–607.
- Lortie, C.J., Brooker, R.W., Choler, P., Kikvidze, Z., Michalet, R., Pugnaire, F.I. & Callaway, R.M. (2004) Rethinking plant community theory. *Oikos*, **107**, 433–438.
- Lotka, A.J. (1932) The growth of mixed populations: two species competing for a common food supply. *Journal of the Washington Academy of Sciences*, **22**, 461–469.
- MacArthur, R. & Levins, R. (1967) The limiting similarity, convergence, and divergence of coexisting species. *The American Naturalist*, **101**, 377–385.
- Mayfield, M.M. & Levine, J.M. (2010) Opposing effects of competitive exclusion on the phylogenetic structure of communities. *Ecology Letters*, **13**, 1085–1093.
- McGill, B.J., Enquist, B.J., Weiher, E. & Westoby, M. (2006) Rebuilding community ecology from functional traits. *Trends in ecology & evolution*, **21**, 178–85.
- Medeiros, M.B. De & Miranda, H.S. (2005) Mortalidade pós-fogo em espécies lenhosas de campo sujo submetido a três queimadas prescritas anuais. *Acta Botanica Brasilica*, **19**, 493–500.
- Oliveras, I. & Malhi, Y. (2016) Many shades of green: the dynamic tropical forest–savannah transition zones. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **371**, 20150308.
- Pausas, J.G., Keeley, J.E. & Schwillk, D.W. (2016) Flammability as an ecological and evolutionary driver (ed M Rees). *Journal of Ecology*.
- Ratnam, J., Bond, W.J., Fensham, R.J., Hoffmann, W. a., Archibald, S., Lehmann, C.E.R., Anderson, M.T., Higgins, S.I. & Sankaran, M. (2011) When is a “forest” a savanna, and why does it matter? *Global Ecology and Biogeography*, **20**, 653–660.
- Rosell, J.A., Gleason, S., Méndez-Alonzo, R., Chang, Y. & Westoby, M. (2014) Bark functional ecology: evidence for tradeoffs, functional coordination, and environment producing bark diversity. *New Phytologist*, **201**, 486–497.
- Schmidt, K.A., Earnhardt, J.M., Brown, J.S. & Holt, R.D. (2000) Habitat selection under temporal heterogeneity: Exorcizing the Ghost of Competition Past. *Ecology*, **81**, 2622–2630.
- Silva, D.M. da & Batalha, M.A. (2011) Defense syndromes against herbivory in a cerrado plant community. *Plant Ecology*, **212**, 181–193.
- Simon, M.F., Grether, R., de Queiroz, L.P., Skema, C., Pennington, R.T. & Hughes, C.E. (2009) Recent assembly of the Cerrado, a neotropical plant diversity hotspot,

- by in situ evolution of adaptations to fire. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 20359–64.
- Soberón, J. (2007) Grinnellian and Eltonian niches and geographic distributions of species. *Ecology Letters*, **10**, 1115–1123.
- Sobral, F.L. & Cianciaruso, M.V. (2012) Estrutura filogenética e funcional de assembléias: (re)montando a ecologia de comunidades em diferentes escalas espaciais. *Bioscience Journal*, **28**, 617–631.
- Tilman, D. (1982) *Resource Competition and Community Structure*. Princeton University Press, Princeton.
- Tilman, D. (1990) Constraints and tradeoffs: toward a predictive theory of competition and succession. *Oikos*, **58**, 3–15.
- Tilman, D. (1993) The maintenance of species richness in plant communities. *Species diversity in ecological communities: historical and geographical perspectives* (eds R.E. Ricklefs & D. Schluter), pp. 13–25. University Chicago Press, Chicago.
- Tilman, D. & Pacala, S. (1993) The maintenance of species richness in plant communities. *Species diversity in ecological communities* (eds R.E. Ricklefs & D. Schluter), pp. 13–25. University Chicago Press, Chicago.
- Vamosi, S.M., Heard, S.B., Vamosi, J.C. & Webb, C.O. (2009) Emerging patterns in the comparative analysis of phylogenetic community structure. *Molecular Ecology*, **18**, 572–592.
- Volterra, V. (1926) Fluctuations in the abundance of species considered mathematically. *Nature*, **118**, 558–560.
- Webb, C.O., Ackerly, D.D., McPeck, M.A. & Donoghue, M.J. (2002) Phylogenies and community ecology. *Annual Review of Ecology and Systematics*, **33**, 475–505.
- Westoby, M. (1998) A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*, **199**, 213–227.
- Westoby, M., Falster, D.S., Moles, A.T., Vesk, P.A. & Wright, I.J. (2002) Plant ecological strategies: some leading dimensions of variation between species. *Annual Review of Ecology and Systematics*, **33**, 125–159.
- Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-bares, J., Chapin, T., Cornelissen, J.H.C., Diemer, M., Flexas, J., Garnier, E., Groom, P.K. & Gulias, J. (2004) The worldwide leaf economics spectrum. *Nature*, **428**, 821–827.

CAPÍTULO I

Formatado nas normas da revista:

New Phytologist

Plant ecological strategies in seasonal forests and savannas species

Leandro Maracahipes¹, Marcos B. Carlucci^{1,2} and Marcus V. Cianciaruso¹

¹Departamento de Ecologia, Instituto de Ciências Biológicas, Universidade Federal de Goiás, Goiânia, Goiás, CP 131, 74001-970, Brazil.

²Departamento de Botânica, Setor de Ciências Biológicas, Universidade Federal do Paraná, CP 19031, Curitiba, PR 81531-980, Brazil.

Summary

- Seasonal forests and savannas are contrasting habitats that occur side by side in Neotropical savanna landscapes. Thus, it is possible to either find plant species common to both forests and savannas (generalists) or exclusive to one of these habitats (specialists). In this study, we aimed to unravel the ecological strategies that determine the occurrence of species in forests, savannas or in both habitats.
- We used data on habitat preference of 284 woody species of seasonal forest and savanna communities in six sites throughout the Brazilian Cerrado. We evaluated the ecological strategies of generalist and specialist species of forests and savannas using data on ecological traits at inter- and intraspecific level.
- We found that acquisitive traits were associated to forest habitat (forest specialist and forest generalist species), while conservative traits were related to savanna habitat (savanna specialist and savanna generalist species).
- Our results demonstrate how functional traits mediate species habitat preference in contrasting habitats. The different strategies adopted by species are in accordance with the limiting factors of species occurrence in each habitat. While acquisitive traits represent competitive ability and fast resource acquisition for forest-specialist species in an environment where light is a limiting resource, conservative traits promote resistance against fire and other environmental stresses for savanna specialists. Furthermore, phenotypic plasticity appears to underlie the ability of species to persist in the contrasting habitats of seasonal forests and savannas.

Key words: habitat preference, specialist species, generalist species; functional diversity; ecological strategies, phenotypic plasticity

Introduction

Understanding the role of species traits on how species are sorted across habitats is an important question in community ecology (Hoffmann *et al.*, 2005, 2012; Ratnam *et al.*, 2011; Pellegrini *et al.*, 2015; Pellegrini, 2016). Habitats with low soil fertility or hard access to nutrients by plants low soil fertility [as African savannas (Wigley *et al.*, 2016)], high elevation [as French Alps, (de Bello *et al.* (2013))], and high fire frequency [as Australian savannas (Murphy *et al.*, 2010)] represent habitats with strong environmental filters to plants (see de Bello *et al.* (2013) for more details). Thus, plants with a conservation resource strategy are more likely to occur in such environments (Wigley *et al.*, 2016; Pellegrini, 2016). On the other hand, in environments with high resource availability plants with acquisition resource strategy are more likely to prevail (Pellegrini, 2016).

In savannas, recurrent fires and edaphic properties, such as low soil fertility and moisture deficits (Murphy & Bowman, 2012; Dantas *et al.*, 2013a,b), should favour species with conservative strategies (Pausas *et al.*, 2016). Indeed, the predominant strategies of savanna species are related to protection against fire (e.g., thick bark, Hoffmann *et al.*, 2009, 2012; Dantas *et al.*, 2013a) and drought (e.g., water storage in stem and root tissues, thick leaves with trichomes and high leaf carbon-nitrogen ratio, Scholz *et al.*, 2007; Schymanski *et al.*, 2013; Dantas *et al.*, 2013a). On the other hand, the predominant strategies of seasonal forest species are related to competition for light (taller plants, large leaves, and high specific leaf area), nutrient usage (high leaf nutrient content) and structural vigor (high wood density) (Hoffmann & Franco, 2003; Dantas *et al.*, 2013a; Pellegrini, 2016). Light is a limiting resource in forest interior, especially during the phase when tree juveniles grows toward the canopy (Hoffmann & Franco, 2003). These two strategies, rapid acquisition or conservation of resources, can be

assessed through trade-offs in functional traits, thereby helping explain why a given species is able to occur in forests or in savannas (Table 1, Westoby, 1998; Lusk *et al.*, 2008; Ratnam *et al.*, 2011; Dantas *et al.*, 2013a).

Forest and savanna habitats occur intermingle, distributed as patches across Neotropical landscapes with savanna predominance. Given this close co-occurrence of habitats, the regional pool of these savanna-dominated landscapes should have species with traits that allow them to establish and persist in one of these environments (Hoffmann *et al.*, 2012; Pausas & Dantas, 2016) or in both of them (Ratnam *et al.*, 2011; Hoffmann *et al.*, 2012; Pausas & Dantas, 2016). Exclusive forest species are unable to establish in savanna habitats because they are not adapted to survive to frequent and intense fires, low soil water availability and low soil fertility (Goodland & Pollard, 1973; Hoffmann *et al.*, 2003; Dantas *et al.*, 2013a). On the other hand, exclusive savannas species are unable to establish in forest, especially due to shade intolerance and little invest in height growth (Hoffmann & Franco, 2003; Poorter, 2009; Rossatto *et al.*, 2013; Silva *et al.*, 2013). Thus, the establishment of forest species in savanna habitats require long intervals without fire, while disturbance and edge effects, like those caused by fire, may enable the persistence of savanna species in forest habitats (Hoffmann & Franco, 2003).

The boundary between forest and savanna is normally abrupt (Hoffmann *et al.*, 2003; Rossatto *et al.*, 2009) and few species are able to occur in both habitats (Hoffmann *et al.*, 2009, 2012). The capacity of some species occur in both habitats can be related to plastic responses of species to different biotic and abiotic factors (Turcotte & Levine, 2016), or because they have strategies that permit colonizing different habitat types (Hoffmann *et al.*, 2012; Silva *et al.*, 2013; Bowman *et al.*, 2015). The ability of species to exhibit different strategies according to the conditions imposed by the

environment is defined as phenotypic plasticity (Miner *et al.*, 2005). Phenotypic plasticity plays a fundamental role on species interactions and coexistence (Violle *et al.*, 2012; Cianciaruso *et al.*, 2012). Many plant species are highly plastic and are able to establish and persist in various environments (Hoffmann *et al.*, 2009, 2012). Intraspecific variability can determine species persistence in environments with strong differences to plant establishment, as forest and savannas. We highlight that considering species habitat preference (habitat generalist versus specialist) and phenotypic plasticity is a new perspective in the quest to understand the distinct ecological strategies of species in seasonal forest and savannas. Several studies have sought to identify which mechanisms determine the species occurrence in forest and savanna habitats (Hoffmann *et al.*, 2003, 2005; Ratnam *et al.*, 2011; Rossatto *et al.*, 2013; Charles-Dominique *et al.*, 2015; Pellegrini *et al.*, 2015). However, information about how different ecological strategies relate to edaphic proprieties and fire disturbance determine the occurrence of species in forests or savannas is still missing, because the few existing studies evaluated a low number of species and just a few traits (Hoffmann *et al.*, 2003, 2004, 2005, 2012; Hao *et al.*, 2008; Rossatto *et al.*, 2009; Ratnam *et al.*, 2011; Silva *et al.*, 2013; Pellegrini, 2016)

Here we evaluated whether forest-specialist, savanna-specialist and generalist woody species (i.e., occurring in both seasonal forest and savannas) have distinct ecological strategies (Table 1). We aimed to answer the following questions: (i) Do trait values differ between habitat-specialist and habitat-generalist species (i.e. species restricted to either seasonal forest or savanna, and species common to seasonal forest and savanna, respectively)? (ii) Do generalist species exhibit intermediate functional traits values between forest- and savanna-specialist species? (iii) Do the same species occurring in forests and savannas show different functional traits values in each

environment? We hypothesized that savanna-specialist woody species would have trait values more related to a conservative use of resource whereas forest-specialist woody species will have functional traits associated to competitive ability and resource acquisition (Table 1). Generalist species would present intermediate values of functional traits between forest- and savanna-specialist species. Additionally, using site-specific data we expect that when a species occurs at both habitats, its traits values will be plastic and respond to the general strategy of each environment, with acquisitive strategy in seasonal forest and conservative strategy in savanna.

Table 1 Functional significance of traits and basis of trait-based patterns predicted in relation to forest and savanna habitats.

Trait	Abbreviations	Unit	Functional significance	Prediction		Rationale
				Forest	Savanna	
Bark thickness	BT	mm	Protection of vital tissue against damage, like that caused by fire ¹	Low	High	Savannas are fire-prone environments and present many species with fire resistance characteristics
Stem-specific density	SSD	mg mm ⁻³	Resistance to physical damage, and consequent structural vigor ^{1,2,3}	Medium	High	Higher wood density in the savanna because stress tolerant plants have greater construction costs in poor soils, and need protection from fires or herbivores. Forest species have medium wood density due to the mixture of species with high wood density (climax species) and low (pioneer species)
Leaf thickness	LT	mm	Related to species strategies of resource acquisition and use, and resistance to physical damage. Every time is correlated with leaf toughness ^{4,5}	Low	High	Savanna species live under severe environmental stress and nutrient scarcity, so that they invest in leaf thickness to assure greater protection against herbivores and water loss
Leaf carbon/nitrogen ratio	C/N	(%) / g/kg (%)	Low values of the ratio indicate higher nutritional quality and palatability of leaves. Nitrogen is the major limiting macronutrient in plants ^{1,8}	Low	High	C/N ratio would be higher among savanna species because the assimilation of N would be lower in the savanna, leading to a higher C/N ratio. Higher C/N ratio can promote greater leaf toughness and protection against abiotic and biotic stress ¹¹
Maximum plant height	Hmax	m	Competitive plant fitness, plant fecundity, tolerance or resistance to disturbances ¹	High	Low	Forest species are taller due to the greater investment in vertical growth resulting from the strong competition for light in the forest canopy
Leaf area	LA	cm ²	Ecological strategy related to resource acquisition and use, responsible to environmental stress and disturbances. High stress tends to select small leaves ¹	High	Low	Forest species have higher leaf area because light is a limiting resource in forest interiors, and higher leaf laminae enable higher absorption of light per leaf
Specific leaf area	SLA	cm ² g ⁻¹	Related to structural defense, resource uptake, resource use efficiency, and growth strategies. It is negatively correlated with leaf lifespan ¹	High	Low	Species from forest environments rich in nutrients can invest in a larger leaf surface per unit of leaf mass, which would improve their competitive ability in forest interiors
Leaf phosphorus content	LPC	g/kg (%)	Related to growth and productivity of plants ⁶	High	Low	Forest environments have higher concentrations of these nutrients available in the soil than savanna environments and more efficient nutrient cycling processes arising from the litter
Leaf potassium content	LKC	g/kg (%)	Plays a critical role in plant growth and metabolism, and contributes to survival of plants that are under various biotic and abiotic stresses ⁷	High	Low	Same rationale of leaf phosphorous content
Leaf calcium content	LCaC	g/kg (%)	Unique macronutrient with diverse but fundamental physiological roles in plant structure and signaling. Plays an important role in structure and increasing resistance of plants, and helps the leaves to grow strong and health ^{8,9}	High	Low	Same rationale of leaf phosphorous content
Leaf magnesium content	LMgC	g/kg (%)	Involved in photosynthetic processes, related with growth and health of plants ¹⁰	High	Low	Same rationale of leaf phosphorous content

¹ Pérez-Harguindeguy *et al.* (2013); ² Chave *et al.* (2009); ³ Larjavaara & Muller-Landau (2010); ⁴ Cianciaruso *et al.* (2012); ⁵ Vile *et al.* (2005); ⁶ Schachtman *et al.* (1998); ⁷ Tripler *et al.* (2006); ⁸ Chapin *et al.* (2011); ⁹ Gilliham *et al.* (2011); ¹⁰ Shaul (2002); ¹¹ Silva & Batalha (2011).

Methods

Study region

The Cerrado domain (Savannas) is a region of approximately $2 \times 10^6 \text{ km}^2$. The study region ranged from 12°S to 22°S. The soils are predominantly characterized as red-yellow latosol, with oxisols and dystrophic soils (Marimon Junior & Haridasan, 2005; Ruggiero *et al.*, 2006; Silva & Batalha, 2008; Pinheiro & Durigan, 2009). These soils show high water permeability and their water available capacity varies in accordance to texture and clay concentrations (Reatto *et al.*, 2008). For a better characterization of the climate and edaphic proprieties of the studied sites sampled see Table S1. The Cerrado is composed of several vegetation mosaics that are the result of the occurrence of various physiognomies (Eiten, 1972), ranging from open savannas to evergreen gallery forests along streams and upland forests of deciduous and evergreen species (Hoffmann *et al.*, 2012).

Sampling design

We selected six study sites throughout the Brazilian Cerrado phytogeographic domain (Fig. 1). The mean distance between pairs of sites was 12 km. The shortest distance between sites was 150 km (Emas National Park and City of Jataí). We adopted a paired sampling design, which consisted in one block of plots placed in savanna (*cerrado stricto sensu*) and another block of plots placed in seasonally dry forests. The shortest distance within site between a forest and a savanna block was 1 km (in Emas National Park). In each locality we sampled woody communities in 10 100-m² plots in each block. We recorded all individuals with diameter at breast ground height (DGH) $\geq 5 \text{ cm}$ in all savanna plots, and all individuals with a diameter breast height (DBH) $\geq 10 \text{ cm}$ in

all forest plots. All individuals were identified to the species level. Nomenclature follows the Brazilian Flora (Forzza *et al.*, 2012).




Figure 1. Location of six study sites of seasonal forests and savannas sampled in the Brazilian Cerrado. 1. Assis Ecological Station – SP, 2. Vassununga State Park – SP, 3. Emas National Park – GO, 4. Jataí – GO, 5. Bacaba Park – MT, 6. Ribeirão Cascalheira – MT.

Trait sampling

We sampled 11 functional traits of 284 species belonging to 57 families distributed in forest and savannas in six sites throughout the Cerrado (Table S2). The species richness ranged from 21 to 52 species in the forest blocks, and from 43 to 89 in the savanna blocks.

Following Pérez-Harguindeguy et al. (2013), we measured the following traits: bark thickness (BT), stem-specific density (SSD), leaf thickness (LT), leaf carbon/nitrogen ratio (C/N), maximum plant height (Hmax), leaf area (LA), specific leaf area (SLA), leaf phosphorous content (LPC), leaf potassium content (LKC) leaf calcium content (LCaC), and leaf magnesium content (LMgC) (see Table 1 for their ecological significance). Whenever possible we measured 10 individuals for each species in each site. When a given species had less than 10 individuals in our plots, we sampled individuals that were nearby the plot (see Table S3). Leaf nutrient concentration (phosphorous, potassium, calcium, magnesium) and leaf carbon/nitrogen ratio were measured for three individuals of each species in each site. Leaf nutrient concentrations were determined in the Laboratory of Soils of the University of Viçosa. All other traits were quantified in each site for all species by sampling individuals whenever possible. We did not measure stem-specific density for palm species because is impossible to collect it without injuring the whole plant.

To obtain leaf trait values, we scanned three to five leaves of each individual. We measured scanned leaf area through of a script that we build using the “EBImage” package (Pau et al., 2014) with R 3.2.1 (<http://www.R-project.org>). Then, we dried leaves at 70 °C for 48 h, and measured leaf dry mass. Leaf thickness of fresh leaves and bark thickness were measured using a digital micrometer. All traits were collected during the rainy season (December to March), which represents the peak of growth and is also the period of maximum leaf expansion and maturity for the studied species.

We classified species according to their habitat of occurrence according to Mendonça *et al.*, (2008) and the Brazilian Flora database (<http://floradobrasil.jbrj.gov.br/>). We considered as savanna specialists or forest specialists all species with restrict occurrence to savanna or forest, respectively. Species

that occur in both habitats according to the literature, but that were sampled only in forest plots were considered “forest-generalist species”. Following the same rationale species that occurred only in savannas plots, but can also occur in forests according to the literature, were considered “savanna-generalist species”. In order to avoid misinterpretations due to potential sample bias we classified the species as “forest-generalist species” and “savanna-generalist species”, and not only like generalists, because we only found them in only one habitat at the field (Table S2). Only 19 species were sampled in both forest and savanna habitats. These species were analysed separately in order to evaluate the potential intraspecific variation in their ecological strategies (see below).

Statistical analyses

We performed ANOVA to test for differences in functional traits between habitat preferences of species (forest specialists, savanna specialists, forest generalist and savanna generalist) with Tukey’s post hoc comparisons, when appropriate. In order to control for type I error, *P*-values were based on Bonferroni’s correction. The ANOVAs were conducted using “aov” function in R (R Core Team, 2015).

Considering that different sources of variation can act on habitat preferences of species, we evaluated the role of variability among habitat preferences, within habitats and among species. This approach enabled us to assess whether the possible differences related to habitat preference in the ANOVA are a result of variation among different sites or changes in species composition. We fitted a general linear mixed model (GLMM) using site, habitat preference and species as nested random factor in the “lme” function in the “nlme” R package (Pinheiro *et al.*, 2016). We adopted this order, for the nested random factors for considering sites as the coarser and species as the finer factor.

To partition the variation among these three components we used the “varcomp” function in the “ape” R package (Paradis *et al.*, 2004).

For the 19 species sampled in both forest and savanna habitats we performed paired *t* test to evaluate how functional trait of these species respond to forest and savanna habitats.

Results

In general, forest specialists and forest generalists presented traits values representative of an acquisitive strategy, whereas savannas specialists and generalists had trait values more related to a conservative strategy. Forest species had higher maximum height, specific leaf area and leaf nutrient content than savanna species (Table 2; Fig. 2). Conversely, savanna species had thicker barks and leaves and higher leaf carbon/nitrogen ratio (Table 2; Fig. 2). Moreover, the traits of forest specialist and forest generalist species did not differed (Table 2; Fig. 2). However, savanna specialist species presented higher values of bark thickness and leaf thickness than savanna generalist species (Table 2; Fig. 2). We did not find any differences for stem-specific density and leaf area in relation to habitat type (Table 2; Fig. 2).

For all traits, the variation among species and habitats explained trait variability (Table 2). In general, the variability among sites was low, with only carbon/nitrogen, phosphorous and calcium presenting intermediate values of variability (Table 2; Fig. S1).

Table 2 Mean values of functional traits ($\pm$ SD) and results of variance component analysis comparing the variability among sites, habitats and species for traits of seasonal forests and savannas in the Brazilian Cerrado. Tukey's post hoc comparisons with Bonferroni corrections ($\alpha = 0.05$ to 0.0045) reveals significant differences between functional traits of forest and savanna. Different letters indicate significant differences between habitat preferences (Tukey test, $p < 0.05$).

Trait	Forest	Generalist		Savanna	F	P	Variability			
		Sampled only in forest	Sampled only in savanna				Among sites	Among habitats	Among species	Residuals
Bark thickness (mm)	1.73 $\pm$ 1.77 ^a	2.37 $\pm$ 2.23 ^a	6.26 $\pm$ 5.33 ^b	8.19 $\pm$ 4.62 ^c	54.35	<0.001	0.00	0.35	0.65	0.00
Stem-specific density (mg mm ⁻³)	0.54 $\pm$ 0.12	0.53 $\pm$ 0.14	0.54 $\pm$ 0.09	0.51 $\pm$ 0.08	1.95	0.121	0.12	0.12	0.66	0.10
Leaf thickness (mm)	0.17 $\pm$ 0.04 ^a	0.18 $\pm$ 0.05 ^a	0.22 $\pm$ 0.08 ^b	0.27 $\pm$ 0.08 ^c	37.78	<0.001	0.07	0.37	0.48	0.08
Leaf C/N ratio (g/kg(%))	2.55 $\pm$ 0.80 ^a	2.65 $\pm$ 0.79 ^a	3.18 $\pm$ 0.86 ^b	3.47 $\pm$ 0.84 ^b	22.34	<0.001	0.10	0.15	0.65	0.10
Maximum plant height (m)	15.8 $\pm$ 5.5 ^a	15.8 $\pm$ 5.5 ^a	6.7 $\pm$ 2.2 ^b	5.6 $\pm$ 1.8 ^b	134.6	<0.001	0.00	0.73	0.27	0.00
Leaf area (cm ²)	175 $\pm$ 281	181 $\pm$ 360	111 $\pm$ 173	90 $\pm$ 120	2.50	0.059	0.01	0.04	0.84	0.11
Specific leaf area (cm ² g ⁻¹)	117 $\pm$ 39.0 ^a	119 $\pm$ 43.3 ^a	88 $\pm$ 26.55 ^b	81.3 $\pm$ 22.4 ^b	24.91	<0.001	0.00	0.46	0.53	0.01
Leaf phosphorous content (g/kg(%))	1.25 $\pm$ 0.53 ^a	1.30 $\pm$ 0.48 ^a	1.11 $\pm$ 0.34 ^{a,b}	1.01 $\pm$ 0.26 ^b	7.28	<0.001	0.29	0.22	0.42	0.07
Leaf potassium content (g/kg(%))	9.25 $\pm$ 6.23 ^a	9.10 $\pm$ 5.14 ^a	6.19 $\pm$ 3.57 ^b	5.48 $\pm$ 2.72 ^b	12.98	<0.001	0.00	0.49	0.47	0.04
Leaf calcium content (g/kg(%))	12.5 $\pm$ 11.7 ^a	9.57 $\pm$ 9.31 ^a	3.99 $\pm$ 2.86 ^b	3.44 $\pm$ 1.74 ^b	24.72	<0.001	0.26	0.48	0.21	0.05
Leaf magnesium content (g/kg(%))	3.05 $\pm$ 1.90 ^{a,b}	3.29 $\pm$ 1.93 ^a	2.43 $\pm$ 1.29 ^{b,c}	2.17 $\pm$ 1.02 ^c	7.69	<0.001	0.06	0.32	0.53	0.09




Figure 2 Mean value and confidence interval to species habitat preference - specialist of forest (F) and savanna (S), and generalist sampled only in forest (FG) and sampled only in savanna (SG) - for traits of seasonal forests and savannas in the Brazilian Cerrado. Different letters indicate significant differences between habitat preferences (Tukey test, $p < 0.05$).

For the 19 species that occurred in both forest and savanna habitats, we found intraspecific differences for only three traits (Fig. 3). Individuals occurring in forests presented higher values of maximum plant height, specific leaf area, while those occurring in savannas had higher carbon/nitrogen ratio (Fig. 3).




Figure 3 Ecological strategies of 19 species common to seasonal forests and savannas environments in the Brazilian Cerrado. Paired t test with Bonferroni corrections ($\alpha = 0.05$ to 0.0045) reveals significant differences between functional traits of forest and savanna.

Discussion

In general, our results were in accordance to our predictions (see Table 1) and demonstrate that forest and savanna species have different ecological strategies. Forest species had higher values for traits related to competitive ability and fast resource acquisition, and savannas species had higher values for traits associated with physical and biological stress damage. We found evidence for fundamental trade-offs involving major plant traits in contrasting tropical habitats, in which a set of species invests in traits related to rapid acquisition of resources, while another group of species allocates more energy in traits that represent conservation of resources and protection against physical and biological damage (Diaz *et al.*, 2004; Wigley *et al.*, 2016). For example, plants with a resource acquisition strategy have leaves with high nutritional quality and fast growth, with higher plant height and specific leaf area, which represents better capture and use of light (Westoby, 1998; Diaz *et al.*, 2004; Hoffmann *et al.*, 2012; Pellegrini, 2016). Because light is a limited resource in the understory species that invest more in acquisitive strategies tend to be more successful in seasonal forests (Hoffmann *et al.*, 2012). On the other hand, species with traits related to conservative strategy tend to persist in savannas, under poor soils and higher effect of strong environmental filters (Westoby, 1998; Dantas *et al.*, 2013a; Pausas *et al.*, 2016).

The distinction between functional traits of forest and savanna species occurs because the species with different growth strategies respond differently to environments so contrasting as forests and savannas and adopt different strategies for persisting in an environment that is either light limiting or fire-prone (Hoffmann & Franco, 2003; Dantas *et al.*, 2016; Pausas *et al.*, 2016). The adoption of these different ecological strategies by species living in habitats drive by different limiting factors often determine their persistence or exclusion (Hoffmann & Franco, 2003; Laureto & Cianciaruso,

2015). Species that do not have fire-adaptive traits may be excluded from fire-prone habitats like savannas (Simon *et al.*, 2009; Pausas *et al.*, 2016). Likewise, species that not have fast growth or shade tolerance may not persist in light limiting habitats (Hoffmann & Franco, 2003; Ratnam *et al.*, 2011). For instance, savanna species presented higher values for traits related to defense against herbivores and water loss (leaf thickness and carbon/nitrogen ratio) and fire protection (bark thickness) (Silva & Batalha, 2011; Dantas *et al.*, 2013b,a; Pausas *et al.*, 2016), representing strategies of stress tolerance in savanna environments (Hoffmann *et al.*, 2012; Pellegrini *et al.*, 2015). On the other hand, forest species have strategies associated with competitive vigor (taller plants), efficiency in capture and use of the light resource (high specific leaf area), and fertility (high leaf concentrations of P, K, Mg, Ca). Thus, forest specialist species cannot establish in savannas because of the effects of environmental filters, and savanna specialist species are unable to establish in forests due to low competitive ability for light (Silva *et al.*, 2013).

We expected that stem-specific density and leaf area would be different between forest and savannas species (see Table 1). However, these traits did not differ between forest and savanna species. A similar wood density to forest and savanna species can be explained because forest species need a high wood density to support taller trees in an environment where height means competitive ability (Chave *et al.*, 2009; Pérez-Harguindeguy *et al.*, 2013). On their turn, savanna species slow growth in a stressful environment have high construction costs (Larjavaara & Muller-Landau, 2010), which leads to high wood density, that also confers higher advantage in persisting in under strong environmental stress, poor soils and high fire frequency [Dantas *et al.* (2013a); but see Hoffmann *et al.* (2009) and Brando *et al.* (2012)]. Even if some studies showed that leaf area is two times greater for forest than savanna species (Hoffmann *et al.*,

2012; Silva *et al.*, 2013), these studies were conducted with a restricted number of species. Savannas species have large root systems (Oliveira *et al.*, 2005) and lead habitat ranging from evergreens to deciduous species (Cianciaruso *et al.*, 2013). The deep roots make the savanna species able to have access to soil water and, therefore, to survive and grow even during the dry season (Oliveira *et al.*, 2005). Yet, many savannas species concentrate the loss of leaves during dry season whereas forest species generally produce and lose leaves throughout the year (Franco *et al.*, 2005; Lenza & Klink, 2006). However, for Neotropical savanna species, Cianciaruso *et al.* (2013) found a lack of differences in leaf traits (SLA and N) among species with distinct leaf habits. In fact, they found that even evergreens and deciduous have similar leaf life spans. This reinforces the idea that the more relevant trade-offs are in the root systems. Also, the similar leaf area between forest and savannas species (but distinct SLA) indicate that leaf dry mass content is main factor determining acquisitive strategy for forest (high SLA and low carbon/nitrogen rate) and conservative strategy for savannas (low SLA and high carbon/nitrogen rate).

When we evaluate the individuals of the same species inhabiting in seasonal forests and savannas, three out of the 11 functional traits differed between species of forest and savanna. Carbon/nitrogen that represent conservative strategy presented higher values in savanna habitat, while maximum plant height and specific leaf area that indicate acquisitive strategy were higher in forest habitat. These opposite responses of the same species in different environments show that the habitat has a strong influence on intraspecific trait variability of generalist species. Intraspecific variability is evident when a given species occurs in different environments expressing distinct trait values (Miner *et al.*, 2005), because non-plastic species can be removed from communities through environmental filters (as fire) and competition by light (Hoffmann & Franco,

2003; Jung *et al.*, 2010). Leaf traits and plant height are traits that have greater phenotypic plasticity and that can shift according to light and resource availability or environmental stress (Hoffmann & Franco, 2003; Hoffmann *et al.*, 2005, 2012; Nicotra *et al.*, 2010). Assuming that some “dominant plants tend to monopolize light and mineral nutrient capture by the development of extensive leaf canopies and root systems” (Campbell *et al.*, 1991), thus species ability to perform shifts in morphological and physiological leaf traits is key to determine the permanence of generalist species (Schlichting, 1986; Callaway *et al.*, 2003). Usually, savanna species have thicker barks than forest species, because bark promote protection of stem in fire-prone habitat (Hoffmann *et al.*, 2003). However, our results showed that the same species in savanna and forest had similar bark thickness.

Savanna species have thick barks as a fire resistance strategy (Dantas *et al.*, 2013a; Rosell *et al.*, 2014; Pausas *et al.*, 2016). Hoffmann *et al.* (2012) calculated a threshold bark thickness of 5.9 mm needed to survive in a low-intensity fire of forest and savanna species. In our study system, the species with individuals inhabiting in seasonal forests and savannas showed mean values below of threshold bark thickness in both habitats (see Table S4). These values below of threshold bark thickness can be explained by the fact that the most of the 19 species occurring in forest and savanna have the forest habitat as preferential habitat (see list of species in Table S4). Moreover, these 19 species present low percentage of occurrence in 376 areas analysed by Ratter *et al.* (2003) that also can be justified by forest preferential habitat these species, because no included forest areas in analyses. We highlight that the same species with individuals occurring in forest and savannas are not proper generalists, which would promote an adaptive advantage in a phytogeographic region with forests and savannas, and with distinct levels of fire. We verified that some traits respond to habitat preference at

interspecific level, but such differences are lost when intraspecific variation is included. However, changes in SLA, plant height and leaf carbon/nitrogen ratio were similar at both inter and intraspecific levels. Therefore, these are likely to be key traits for adaptation of individuals in forest and savannas.

Concluding remarks

We showed that habitat preference determines the value of functional traits. Traits related to an acquisitive strategy were associated to and higher in the forest habitat, while traits with conservative strategy were related to and higher in the savanna habitat (Diaz *et al.*, 2004). We highlight that these different strategies are related to competitive ability and resource acquisition of forest-specialist species in an environment in which light is a limiting resource as forests, and related to defense against fire and physical and biological stress of savanna-specialist species in an environment with high fire frequency and poor-soils as savannas (Pellegrini, 2016; Pausas *et al.*, 2016). Generalist species with occurrence restrict to forest or savanna in our study generally presented values of functional traits intermediary between forest- and savanna-specialist species. However, generalist species of forests and savannas were more similar to specialist species of each habitat than each other. This indicates that the habitat play an important role on functional traits, determining values according to limiting factors of each environment. This pattern was also observed when we evaluated the same species occurring in forest and savanna, demonstrating that the habitat has also a strong influence on intraspecific trait variability of generalist species. Thus, the phenotypic plasticity of the species may confer ability to persistence in a habitat with stronger competition for light (forest) or in a habitat with strong environmental filters and stress (savanna).

Acknowledgements

This research was by CNPq-Brazil (#563621/2010-9, #478747/2009-8, and PELD - SITE 13), FAPEG-GO (#201110267000130/31-10), and CAPES-Brazil via scholarships to L.M. and a postdoctoral fellowship to M.B.C. (PNPD #1454013). MVC has a productivity grant awarded by CNPq (307796/2015-9). We thank to all field team responsible for sampling data: Edmar Oliveira, Leonardo Maracahipes, Letícia Gomes, Livia Laureto, Danira Padilha, Mônica Forsthofer, Mariângela Abreu, Josias Santos, Fernando Ribeiro. Thank for Vinicius Dantas by help in design of mixed model. We also thank to taxonomists Geraldo Franco, Osnir Aguiar, João Baitelo, Natália Ivanauskas and Renato Mello-Silva for the identification of some species.

References

- Bello F de, Lavorel S, Lavergne S, Albert CH, Boulangeat I, Mazel F, Thuiller W. 2013.** Hierarchical effects of environmental filters on the functional structure of plant communities: a case study in the French Alps. *Ecography* **36**: 393–402.
- Bowman DMJS, Perry GLW, Marston JB. 2015.** Feedbacks and landscape-level vegetation dynamics. *Trends in Ecology & Evolution* **30**: 255–260.
- Brando PM, Nepstad DC, Balch JK, Bolker B, Christman MC, Coe M, Putz FE. 2012.** Fire-induced tree mortality in a neotropical forest: the roles of bark traits, tree size, wood density and fire behavior. *Global Change Biology* **18**: 630–641.
- Callaway RM, Pennings SC, Richards CL. 2003.** Phenotypic plasticity and interactions among plants. *Ecology* **84**: 1115–1128.
- Campbell BD, Grime JP, Mackey JML. 1991.** A trade-off between scale and precision in resource foraging. *Oecologia* **87**: 532–538.
- Chapin FS, Matson PA, Vitousek PM. 2011.** Plant Nutrient Use. Principles of terrestrial ecosystem ecology. New York, NY: Springer New York, 229–258.
- Charles-Dominique T, Beckett H, Midgley GF, Bond WJ. 2015.** Bud protection: a key trait for species sorting in a forest-savanna mosaic. *New Phytologist* **207**: 1052–1060.
- Chave J, Coomes D, Jansen S, Lewis SL, Swenson NG, Zanne AE. 2009.** Towards a worldwide wood economics spectrum. *Ecology Letters* **12**: 351–366.
- Cianciaruso M V., Silva IA, Batalha MA, Gaston KJ, Petchey OL. 2012.** The influence of fire on phylogenetic and functional structure of woody savannas: moving

from species to individuals. *Perspectives in Plant Ecology, Evolution and Systematics* **14**: 205–216.

Cienciaruso M V., Silva IA, Manica LT, Souza JP. 2013. Leaf habit does not predict leaf functional traits in cerrado woody species. *Basic and Applied Ecology* **14**: 404–412.

Dantas VL, Batalha MA, Pausas JG. 2013a. Fire drives functional thresholds on the savanna–forest transition. *Ecology* **94**: 2454–2463.

Dantas V de L, Hirota M, Oliveira RS, Pausas JG. 2016. Disturbance maintains alternative biome states (M Rejmanek, Ed.). *Ecology Letters* **19**: 12–19.

Dantas V de L, Pausas JG, Batalha MA, Loiola P de P, Cienciaruso MV. 2013b. The role of fire in structuring trait variability in Neotropical savannas. *Oecologia* **171**: 487–494.

Diaz S, Hodgson JG, Thompson K, Cabido M, Cornelissen JHC, Jalili A, Montserrat-Martí G, Grime JP, Zarrinkamar F, Asri Y, et al. 2004. The plant traits that drive ecosystems: evidence from three continents. *Journal of Vegetation Science* **15**: 295–304.

Eiten G. 1972. The cerrado vegetation of Brazil. *Botanical Reviews* **38**: 201–341.

Forzza RC, Baumgratz JFA, Bicudo CEM, Canhos DAL, Carvalho AA, Coelho MAN, Costa AF, Costa DP, Hopkins MG, Leitman PM, et al. 2012. New Brazilian floristic list highlights conservation challenges. *BioScience* **62**: 39–45.

Franco AC, Bustamante M, Caldas LS, Goldstein G, Meinzer FC, Kozovits AR, Rundel P, Coradin VTR. 2005. Leaf functional traits of Neotropical savanna trees in relation to seasonal water deficit. *Trees* **19**: 326–335.

Gilliam M, Dayod M, Hocking BJ, Xu B, Conn SJ, Kaiser BN, Leigh RA, Tyerman SD. 2011. Calcium delivery and storage in plant leaves: exploring the link with water flow. *Journal of Experimental Botany* **62**: 2233–2250.

Goodland R, Pollard R. 1973. The Brazilian cerrado vegetation: a fertility gradient. *Journal of Ecology* **61**: 219–224.

Hao G-Y, Hoffmann WA, Scholz FG, Bucci SJ, Meinzer FC, Franco AC, Cao K-F, Goldstein G. 2008. Stem and leaf hydraulics of congeneric tree species from adjacent tropical savanna and forest ecosystems. *Oecologia* **155**: 405–415.

Hoffmann WA, Adasme R, Haridasan M, Carvalho MT, Geiger EL, Pereira M a B, Gotsch SG, Franco AC. 2009. Tree topkill, not mortality, governs the dynamics of savanna-forest boundaries under frequent fire in central Brazil. *Ecology* **90**: 1326–37.

Hoffmann WA, Franco AC. 2003. Comparative growth analysis of tropical forest and savanna woody plants using phylogenetically independent contrasts. *Journal of Ecology* **91**: 475–484.

Hoffmann WA, Franco AC, Moreira MZ, Haridasan M. 2005. Specific leaf area explains differences in leaf traits between congeneric savanna and forest trees. *Functional Ecology* **19**: 932–940.

Hoffmann W a, Geiger EL, Gotsch SG, Rossatto DR, Silva LCR, Lau OL, Haridasan M, Franco AC. 2012. Ecological thresholds at the savanna-forest boundary: how plant traits, resources and fire govern the distribution of tropical biomes (F Lloret,

Ed.). *Ecology Letters* **15**: 759–768.

Hoffmann WA, Orthen B, Franco AC. 2004. Constraints to seedling success of savanna and forest trees across the savanna-forest boundary. *Oecologia* **140**: 252–260.

Hoffmann WA, Orthen B, Nascimento PKV Do. 2003. Comparative fire ecology of tropical savanna and forest trees. *Functional Ecology* **17**: 720–726.

Jung V, Violle C, Mondy C, Hoffmann L, Muller S. 2010. Intraspecific variability and trait-based community assembly. *Journal of Ecology* **98**: 1134–1140.

Larjavaara M, Muller-Landau HC. 2010. Rethinking the value of high wood density. *Functional Ecology* **24**: 701–705.

Laureto LMO, Cianciaruso M V. 2015. Trait distribution patterns in savanna and forest plant assemblages and their relationship with soil features. *Plant Ecology* **216**: 629–639.

Lenza E, Klink CA. 2006. Comportamento fenológico de espécies lenhosas em um cerrado sentido restrito de Brasília, DF. *Revista Brasileira de Botânica* **29**: 627–638.

Lusk C, Reich P, Montgomery R, Ackerly D, Cavender-Bares J. 2008. Why are evergreen leaves so contrary about shade? *Trends in Ecology & Evolution* **23**: 299–303.

Marimon Junior BH, Haridasan M. 2005. Comparação da vegetação arbórea e características edáficas de um cerrado e um cerrado sensu stricto em áreas adjacentes sobre solo distrófico no leste de Mato Grosso, Brasil. *Acta Botanica Brasilica* **19**: 913–926.

Mendonça RC, Felfili JM, Walter BMT, Silva Junior MC, Rezende A V., Filgueiras TS, Nogueira PE, Fagg CW. 2008. *Cerrado: ecologia e flora* (SM Sano, SP Almeida, and JF Ribeiro, Eds.). Brasília-DF: Embrapa Cerrados.

Miner BG, Sultan SE, Morgan SG, Padilla DK, Relyea RA. 2005. Ecological consequences of phenotypic plasticity. *Trends in Ecology & Evolution* **20**: 685–692.

Murphy BP, Bowman DMJS. 2012. What controls the distribution of tropical forest and savanna? *Ecology Letters* **15**: 748–758.

Murphy BP, Russell-Smith J, Prior LD. 2010. Frequent fires reduce tree growth in northern Australian savannas: implications for tree demography and carbon sequestration. *Global Change Biology* **16**: 331–343.

Nicotra AB, Atkin OK, Bonser SP, Davidson AM, Finnegan EJ, Mathesius U, Poot P, Purugganan MD, Richards CL, Valladares F, et al. 2010. Plant phenotypic plasticity in a changing climate. *Trends in Plant Science* **15**: 684–692.

Oliveira RS, Bezerra L, Davidson EA, Pinto F, Klink CA, Nepstad DC, Moreira A. 2005. Deep root function in soil water dynamics in cerrado savannas of central Brazil. *Functional Ecology* **19**: 574–581.

Paradis E, Claude J, Strimmer K. 2004. APE: analyses of phylogenetics and evolution in R language. : 289–290.

Pausas JG, Dantas V de L. 2016. Scale matters: fire-vegetation feedbacks are needed to explain tropical tree cover at the local scale. *Global Ecology and Biogeography*.

Pausas JG, Keeley JE, Schwilk DW. 2016. Flammability as an ecological and

evolutionary driver (M Rees, Ed.). *Journal of Ecology*.

Pellegrini AFA. 2016. Nutrient limitation in tropical savannas across multiple scales and mechanisms. *Ecology* **97**: 313–324.

Pellegrini AFA, Franco AC, Hoffmann WA. 2015. Shifts in functional traits elevate risk of fire-driven tree dieback in tropical savanna and forest biomes. *Global Change Biology*: n/a-n/a.

Pérez-Harguindeguy N, Díaz S, Garnier E, Lavorel S, Poorter H, Jaureguiberry P, Bret-Harte MS, Cornwell WK, Craine JM, Gurvich DE, et al. 2013. New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany* **61**: 167–234.

Pinheiro J, Bates D, DebRoy D, Team C. 2016. nlme: Linear and Nonlinear Mixed Effects Models.

Pinheiro EDS, Durigan G. 2009. Dinâmica espaço-temporal (1962-2006) das fitofisionomias em unidade de conservação do Cerrado no sudeste do Brasil. *Revista Brasileira de Botânica* **32**: 441–454.

Poorter L. 2009. Leaf traits show different relationships with shade tolerance in moist versus dry tropical forests. *New Phytologist* **181**: 890–900.

Ratnam J, Bond WJ, Fensham RJ, Hoffmann W a., Archibald S, Lehmann CER, Anderson MT, Higgins SI, Sankaran M. 2011. When is a ‘forest’ a savanna, and why does it matter? *Global Ecology and Biogeography* **20**: 653–660.

Ratter JA, Bridgewater S, Ribeiro JF. 2003. Analysis of the floristic composition of the Brazilian cerrado vegetation III: comparison of the woody vegetation of 376 areas. *Edinburgh Journal of Botany* **60**: 57–109.

Reatto A, Correia JR, Spera ST, Martins É de S. 2008. Solos do bioma Cerrado: aspectos pedológicos. In: Sano SM,, In: Almeida SP,, In: Ribeiro JF, eds. Cerrado: ecologia e flora. Brasília-DF: Embrapa Cerrados, 107–149.

Rosell JA, Gleason S, Méndez-Alonzo R, Chang Y, Westoby M. 2014. Bark functional ecology: evidence for tradeoffs, functional coordination, and environment producing bark diversity. *New Phytologist* **201**: 486–497.

Rossatto DR, Hoffmann WA, de Carvalho Ramos Silva L, Haridasan M, Sternberg LSL, Franco AC. 2013. Seasonal variation in leaf traits between congeneric savanna and forest trees in Central Brazil: implications for forest expansion into savanna. *Trees* **27**: 1139–1150.

Rossatto DR, Hoffmann WA, Franco AC. 2009. Differences in growth patterns between co-occurring forest and savanna trees affect the forest-savanna boundary. *Functional Ecology* **23**: 689–698.

Ruggiero PGC, Pivello VR, Sparovek G, Teramoto E, Pires Neto AG. 2006. Relação entre solo, vegetação e topografia em área de cerrado (Parque Estadual de Vassununga, SP): como se expressa em mapeamentos? *Acta Botanica Brasilica* **20**: 383–394.

Schachtman D, Reid RJ, Ayling SM. 1998. Phosphorus uptake by plants: from soil to cell. *Plant Physiology* **116**: 447–453.

- Schlichting CD. 1986.** The evolution of phenotypic plasticity in plants. *Annual Review of Ecology and Systematics* **17**: 667–693.
- Scholz FG, Bucci SJ, Goldstein G, Meinzer FC, Franco AC, Miralles-Wilhelm F. 2007.** Biophysical properties and functional significance of stem water storage tissues in Neotropical savanna trees. *Plant, Cell & Environment* **30**: 236–248.
- Schymanski SJ, Or D, Zwieniecki M. 2013.** Stomatal Control and Leaf Thermal and Hydraulic Capacitances under Rapid Environmental Fluctuations (W Bauerle, Ed.). *PLoS ONE* **8**: e54231.
- Shaul O. 2002.** Magnesium transport and function in plants: the tip of the iceberg. *BioMetals* **15**: 307–321.
- Silva DM da, Batalha MA. 2008.** Soil–vegetation relationships in cerrados under different fire frequencies. *Plant and Soil* **311**: 87–96.
- Silva DM da, Batalha MA. 2011.** Defense syndromes against herbivory in a cerrado plant community. *Plant Ecology* **212**: 181–193.
- Silva LCR, Hoffmann WA, Rossatto DR, Haridasan M, Franco AC, Horwath WR. 2013.** Can savannas become forests? A coupled analysis of nutrient stocks and fire thresholds in central Brazil. *Plant and Soil* **373**: 829–842.
- Simon MF, Grether R, de Queiroz LP, Skema C, Pennington RT, Hughes CE. 2009.** Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by in situ evolution of adaptations to fire. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 20359–64.
- Tripler CE, Kaushal SS, Likens GE, Todd Walter M. 2006.** Patterns in potassium dynamics in forest ecosystems. *Ecology Letters* **9**: 451–466.
- Turcotte MM, Levine JM. 2016.** Phenotypic plasticity and species coexistence. *Trends in Ecology & Evolution* **31**: 803–813.
- Vile D. 2005.** Specific leaf area and dry matter content estimate thickness in laminar leaves. *Annals of Botany* **96**: 1129–1136.
- Violle C, Enquist BJ, McGill BJ, Jiang L, Albert CH, Hulshof C, Jung V, Messier J. 2012.** The return of the variance: intraspecific variability in community ecology. *Trends in Ecology & Evolution* **27**: 244–252.
- Westoby M. 1998.** A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil* **199**: 213–227.
- Wigley BJ, Slingsby JA, Díaz S, Bond WJ, Fritz H, Coetsee C. 2016.** Leaf traits of African woody savanna species across climate and soil fertility gradients: evidence for conservative vs . acquisitive resource use strategies. *Journal of Ecology*.

Supporting Information

Table S1. Climate characteristics and edaphic proprieties of the six sites sampled in Brazilian Cerrado. The values represented a range, when available, of 20 years of data collected in the nearest weather station of each site. The fire frequency applies only to savannas sites. A low fire frequency indicates an interval about of 5 years among fire events, and high fire frequency of 10 years.

Environmental characteristics	Assis Ecological State - SP	Vassununga State Park - SP	Emas National Park - GO	Jataí - GO	Bacaba Park - MT	Ribeirão Cascalheira - MT
Coordinates	50W 22', 22S 35'	47W 37', 21S 36'	52W 59', 17S 54'	51W 32', 17S 45'	52W 21', 14S 42'	51W 46', 12S 49'
Fire frequency	low	low	High	high	high	low
Climate Type	Cfa	Cwa	Aw	Aw	Aw	Aw
Altitude (m)	560	740	840	860	325	374
Mean Annual Temp. (°C)	23.6	22.3	24.6	22.6	24.8	25.4
Mean Minimum Temp. (°C)	19.0	16.6	18.9	17.0	18.9	20.1
Precipitation (mm yr ⁻¹)	1352	1437	1745	1605	1433	1822
Evapotranspiration (mm)	129	92	82	121	134	136
Relative Moisture (%)	64	70	69	69	77	71
Soil characteristics	Forest savanna	Forest savanna	Forest savanna	Forest savanna	Forest savanna	Forest savanna
N	0.32(0.14); 0.25(0.10)	0.22(0.06); 0.12(0.04)	0.26(0.09); 0.09(0.04)	0.18(0.03); 0.17(0.04)	0.11(0.02); 0.08(0.01)	0.10(0.01); 0.05(0.01)
P	8.39(4.26); 3.91(0.31)	4.51(4.58); 3.89(3.13)	1.79(0.65); 0.56(0.20)	1.57(0.66); 1.03(0.76)	1.15(0.57); 2.03(0.14)	1.75(0.25); 2.04(0.63)
K	46.3(16.9); 11.8(3.82)	11.2(39.8); 13.1(3.33)	42.4(7.43); 31.4(8.06)	28.4(12.0); 22.3(5.51)	46.7(10.1); 46.2(6.65)	17.7(3.20); 18.3(7.42)
SOM	3.10(1.12); 1.70(0.60)	5.13(0.96); 3.10(0.50)	5.87(1.64); 3.08(1.30)	6.22(1.13); 4.79(0.83)	1.07(0.56); 1.73(0.31)	2.49(0.28); 1.83(0.57)
Clay	11.2(2.64); 8.0(1.79)	44.7(10.8); 10.6(0.80)	26.1(7.13); 29.4(18.7)	54.6(7.32); 51.0(3.69)	20.1(3.42); 12.6(2.33)	16.7(2.10); 2.9(1.14)
pH	7.09(0.67); 4.85(0.13)	6.16(0.65); 4.51(0.09)	4.20(0.42); 4.30(0.25)	5.26(0.15); 5.26(0.11)	4.65(0.09); 5.04(0.13)	4.10(0.08); 4.95(0.22)
Al	0; 0.88(0.15)	0.12(0.32); 1.70(0.20)	1.65(0.37); 1.19(0.27)	1.03(0.37); 0.65(0.26)	0.85(0.18); 1.16(0.13)	1.12(0.11); 0.77(0.11)

Table S2 List of species and families classified according to habitat preference. The species are ordered according to family. In bold the 19 species that occurred in both forest and savanna habitats in our study and for which we have intraspecific trait variability.

Species	Family	Habitat preference
<i>Anacardium occidentale</i> L.	Anacardiaceae	Generalist savanna
<i>Astronium fraxinifolium</i> Schott	Anacardiaceae	Generalist
<i>Astronium graveolens</i> Jacq.	Anacardiaceae	Generalist forest
<i>Myracrodruon urundeuva</i> Allemão	Anacardiaceae	Generalist forest
<i>Tapirira guianensis</i> Aubl.	Anacardiaceae	Generalist
<i>Tapirira obtusa</i> (Benth.) J.D.Mitch.	Anacardiaceae	Generalist forest
<i>Tapirira</i> sp.	Anacardiaceae	Forest
<i>Annona cacans</i> Warm.	Annonaceae	Generalist forest
<i>Annona coriacea</i> Mart.	Annonaceae	Savanna
<i>Annona crassiflora</i> Mart.	Annonaceae	Generalist savanna
<i>Annona sylvatica</i> A. St.-Hil.	Annonaceae	Generalist forest
<i>Bocageopsis mattogrossensis</i> (R.E.Fr.) R.E.Fr.	Annonaceae	Generalist forest
<i>Duguetia marcgraviana</i> Mart.	Annonaceae	Generalist forest
<i>Ephedranthus parviflorus</i> S.Moore	Annonaceae	Generalist forest
<i>Guatteria blepharophylla</i> Mart.	Annonaceae	Forest
<i>Xylopia amazonica</i> R.E.Fr.	Annonaceae	Forest
<i>Xylopia aromatica</i> (Lam.) Mart.	Annonaceae	Generalist
<i>Xylopia frutescens</i> Aubl.	Annonaceae	Forest
<i>Xylopia sericea</i> A.St.-Hil.	Annonaceae	Generalist savanna
<i>Aspidosperma desmanthum</i> Benth. ex Müll.Arg.	Apocynaceae	Forest
<i>Aspidosperma discolor</i> A.DC.	Apocynaceae	Forest
<i>Aspidosperma macrocarpon</i> Mart.	Apocynaceae	Generalist savanna
<i>Aspidosperma multiflorum</i> A.DC.	Apocynaceae	Savanna
<i>Aspidosperma nobile</i> Müll.Arg.	Apocynaceae	Generalist savanna
<i>Aspidosperma polyneuron</i> Müll.Arg.	Apocynaceae	Generalist forest
<i>Aspidosperma ramiflorum</i> Müll.Arg.	Apocynaceae	Forest
<i>Aspidosperma tomentosum</i> Mart.	Apocynaceae	Generalist savanna
<i>Hancornia speciosa</i> Gomes	Apocynaceae	Savanna
<i>Himatanthus obovatus</i> (Müll.Arg.) Woodson	Apocynaceae	Savanna
<i>Himatanthus sucuuba</i> (Spruce ex Müll.Arg.) Woodson	Apocynaceae	Forest
<i>Schefflera macrocarpa</i> (Cham. & Schltdl.) Frodin	Araliaceae	Generalist savanna
<i>Schefflera morototoni</i> (Aubl.) Maguire, Steyerl. & Frodin	Araliaceae	Generalist forest
<i>Schefflera vinosa</i> (Cham. & Schltdl.) Frodin & Fiaschi	Araliaceae	Savanna
<i>Euterpe edulis</i> Mart.	Arecaceae	Forest
<i>Oenocarpus distichus</i> Mart.	Arecaceae	Generalist forest
<i>Syagrus comosa</i> (Mart.) Mart.	Arecaceae	Savanna
<i>Syagrus flexuosa</i> (Mart.) Becc.	Arecaceae	Savanna

Species	Family	Habitat preference
<i>Moquiniastrum polymorphum</i> (Less.) G. Sancho	Asteraceae	Generalist savanna
<i>Piptocarpha rotundifolia</i> (Less.) Baker	Asteraceae	Savanna
<i>Vernonia</i> sp.	Asteraceae	Savanna
<i>Handroanthus heptaphyllus</i> (Vell.) Mattos	Bignoniaceae	Forest
<i>Handroanthus ochraceus</i> (Cham.) Mattos	Bignoniaceae	Savanna
<i>Handroanthus serratifolius</i> (Vahl) S.O.Grose	Bignoniaceae	Generalist forest
<i>Jacaranda copaia</i> (Aubl.) D.Don	Bignoniaceae	Forest
<i>Jacaranda micrantha</i> Cham.	Bignoniaceae	Forest
<i>Tabebuia aurea</i> (Silva Manso) Benth. & Hook.f. ex S.Moore	Bignoniaceae	Savanna
<i>Zeyheria tuberculosa</i> (Vell.) Bureau ex Verl.	Bignoniaceae	Forest
<i>Cordia sellowiana</i> Cham.	Boraginaceae	Generalist forest
<i>Cordia trichotoma</i> (Vell.) Arráb. ex Steud.	Boraginaceae	Generalist forest
<i>Protium pilosissimum</i> Engl.	Burseraceae	Forest
<i>Protium spruceanum</i> (Benth.) Engl.	Burseraceae	Forest
<i>Protium unifoliolatum</i> Engl.	Burseraceae	Forest
<i>Tetragastris altissima</i> (Aubl.) Swart	Burseraceae	Forest
<i>Trattinnickia glaziovii</i> Swart	Burseraceae	Forest
<i>Trattinnickia</i> sp.	Burseraceae	Forest
<i>Kielmeyera coriacea</i> Mart.	Calophyllaceae	Savanna
<i>Kielmeyera rubriflora</i> Cambess.	Calophyllaceae	Savanna
<i>Jacaratia spinosa</i> (Aubl.) A.DC.	Caricaceae	Forest
<i>Caryocar brasiliense</i> A.St.-Hil.	Caryocaraceae	Savanna
<i>Cheiloclinium cognatum</i> (Miers) A.C.Sm.	Celastraceae	Forest
<i>Plenckia populnea</i> Reissek	Celastraceae	Savanna
<i>Salacia crassifolia</i> (Mart. ex Schult.) G. Don	Celastraceae	Savanna
<i>Couepia grandiflora</i> (Mart. & Zucc.) Benth. ex Hook.f.	Chrysobalanaceae	Generalist savanna
<i>Hirtella glandulosa</i> Spreng.	Chrysobalanaceae	Generalist
<i>Hirtella racemosa</i> Lam.	Chrysobalanaceae	Generalist forest
<i>Licania apetala</i> (E.Mey.) Fritsch	Chrysobalanaceae	Forest
<i>Licania blackii</i> Prance	Chrysobalanaceae	Forest
<i>Licania gardneri</i> (Hook.f.) Fritsch.	Chrysobalanaceae	Forest
<i>Licania humilis</i> Cham. & Schltdl.	Chrysobalanaceae	Savanna
<i>Licania kunthiana</i> Hook.f.	Chrysobalanaceae	Generalist forest
<i>Licania minutiflora</i> (Sagot) Fritsch	Chrysobalanaceae	Forest
<i>Buchenavia tetraphylla</i> (Aubl.) R.A.Howard	Combretaceae	Generalist forest
<i>Buchenavia tomentosa</i> Eichler	Combretaceae	Generalist
<i>Terminalia argentea</i> Mart.	Combretaceae	Generalist savanna
<i>Connarus perrottetii</i> (DC.) Planch.	Connaraceae	Forest
<i>Connarus suberosus</i> Planch.	Connaraceae	Savanna
<i>Rourea induta</i> Planch.	Connaraceae	Savanna
<i>Curatella americana</i> L.	Dilleniaceae	Savanna
<i>Davilla elliptica</i> A.St.-Hil.	Dilleniaceae	Savanna
<i>Diospyros hispida</i> A.DC.	Ebenaceae	Generalist savanna

Species	Family	Habitat preference
<i>Sloanea monosperma</i> Vell.	Elaeocarpaceae	Forest
<i>Sloanea pubescens</i> Benth.	Elaeocarpaceae	Forest
<i>Sloanea sinemariensis</i> Aubl.	Elaeocarpaceae	Forest
<i>Erythroxylum engleri</i> O.E.Schulz	Erythroxylaceae	Savanna
<i>Erythroxylum suberosum</i> A.St.-Hil.	Erythroxylaceae	Savanna
<i>Erythroxylum testaceum</i> Peyr.	Erythroxylaceae	Savanna
<i>Erythroxylum tortuosum</i> Mart.	Erythroxylaceae	Savanna
<i>Alchornea glandulosa</i> Poepp.	Euphorbiaceae	Generalist forest
<i>Chaetocarpus echinocarpus</i> (Baill.) Ducke	Euphorbiaceae	Generalist
<i>Croton floribundus</i> Spreng.	Euphorbiaceae	Generalist forest
<i>Croton piptocalyx</i> Müll.Arg.	Euphorbiaceae	Forest
<i>Mabea fistulifera</i> Mart.	Euphorbiaceae	Generalist forest
<i>Mabea paniculata</i> Spruce ex Benth.	Euphorbiaceae	Forest
<i>Maprounea guianensis</i> Aubl.	Euphorbiaceae	Generalist savanna
<i>Micrandra elata</i> (Didr.) Müll.Arg.	Euphorbiaceae	Forest
<i>Pera coccinea</i> (Benth.) Müll.Arg.	Euphorbiaceae	Forest
<i>Pera glabrata</i> (Schott) Poepp. ex Baill.	Euphorbiaceae	Generalist savanna
<i>Pera bicolor</i> (Klotzsch) Müll.Arg.	Euphorbiaceae	Forest
<i>Albizia niopoides</i> (Spruce ex Benth.) Burkart	Fabaceae	Forest
<i>Anadenanthera peregrina</i> (L.) Speg.	Fabaceae	Generalist savanna
<i>Andira cujabensis</i> Benth.	Fabaceae	Generalist savanna
<i>Andira vermifuga</i> Benth.	Fabaceae	Generalist savanna
<i>Apuleia leiocarpa</i> (Vogel) J.F.Macbr.	Fabaceae	Forest
<i>Bowdichia virgilioides</i> Kunth	Fabaceae	Generalist savanna
<i>Copaifera langsdorffii</i> Desf.	Fabaceae	Generalist
<i>Dalbergia miscolobium</i> Benth.	Fabaceae	Savanna
<i>Dimorphandra mollis</i> Benth.	Fabaceae	Savanna
<i>Diptotropis purpurea</i> (Rich.) Amshoff	Fabaceae	Forest
<i>Dipteryx alata</i> Vogel	Fabaceae	Generalist savanna
<i>Diptychandra aurantiaca</i> Tul.	Fabaceae	Generalist savanna
<i>Enterolobium gummiiferum</i> (Mart.) J.F.Macbr.	Fabaceae	Savanna
<i>Holocalyx balansae</i> Micheli	Fabaceae	Forest
<i>Hymenaea courbaril</i> L.	Fabaceae	Forest
<i>Hymenaea stigonocarpa</i> Hayne	Fabaceae	Savanna
<i>Inga heterophylla</i> Willd.	Fabaceae	Forest
<i>Inga striata</i> Benth.	Fabaceae	Forest
<i>Inga thibaudiana</i> DC.	Fabaceae	Forest
<i>Leptolobium dasycarpum</i> Vogel	Fabaceae	Savanna
<i>Leptolobium elegans</i> Vogel	Fabaceae	Savanna
<i>Luetzelburgia praecox</i> (Harms) Harms	Fabaceae	Generalist savanna
<i>Machaerium acutifolium</i> Vogel	Fabaceae	Generalist savanna
<i>Machaerium brasiliense</i> Vogel	Fabaceae	Generalist savanna
<i>Mimosa laticifera</i> Rizzini & A.Mattos	Fabaceae	Savanna
<i>Myroxylon peruiferum</i> L.f.	Fabaceae	Forest

Species	Family	Habitat preference
<i>Ormosia arborea</i> (Vell.) Harms	Fabaceae	Forest
<i>Ormosia fastigiata</i> Tul.	Fabaceae	Forest
<i>Ormosia paraensis</i> Ducke	Fabaceae	Forest
<i>Peltogyne confertiflora</i> (Hayne) Benth.	Fabaceae	Generalist savanna
<i>Piptadenia gonoacantha</i> (Mart.) J.F.Macbr.	Fabaceae	Forest
<i>Plathymenia reticulata</i> Benth.	Fabaceae	Savanna
<i>Platypodium elegans</i> Vogel	Fabaceae	Generalist savanna
<i>Pterodon emarginatus</i> Vogel	Fabaceae	Generalist savanna
<i>Pterodon pubescens</i> (Benth.) Benth.	Fabaceae	Generalist savanna
<i>Senegalia polyphylla</i> (DC.) Britton	Fabaceae	Generalist forest
<i>Stryphnodendron adstringens</i> (Mart.) Coville	Fabaceae	Savanna
<i>Stryphnodendron rotundifolium</i> Mart.	Fabaceae	Savanna
<i>Tachigali aurea</i> Tul.	Fabaceae	Savanna
<i>Tachigali vulgaris</i> L.F. Gomes da Silva & H.C. Lima	Fabaceae	Generalist
<i>Vatairea macrocarpa</i> (Benth.) Ducke	Fabaceae	Savanna
<i>Sacoglottis guianensis</i> Benth.	Humiriaceae	Generalist forest
<i>Emmotum nitens</i> (Benth.) Miers	Icacinaeae	Generalist savanna
<i>Aegiphila sellowiana</i> Cham.	Lamiaceae	Forest
<i>Cryptocarya moschata</i> Nees & Mart.	Lauraceae	Forest
<i>Endlicheria paniculata</i> (Spreng.) J.F.Macbr.	Lauraceae	Forest
<i>Mezilaurus crassiramea</i> (Meisn.) Taub. ex Mez	Lauraceae	Generalist savanna
<i>Nectandra cuspidata</i> Nees & Mart.	Lauraceae	Generalist forest
<i>Nectandra megapotamica</i> (Spreng.) Mez	Lauraceae	Forest
<i>Ocotea corymbosa</i> (Meisn.) Mez	Lauraceae	Generalist savanna
<i>Ocotea dispersa</i> (Nees & Mart.) Mez	Lauraceae	Forest
<i>Ocotea guianensis</i> Aubl.	Lauraceae	Forest
<i>Ocotea indecora</i> (Schott) Mez	Lauraceae	Forest
<i>Ocotea leucoxydon</i> (Sw.) Laness.	Lauraceae	Forest
<i>Cariniana legalis</i> (Mart.) Kuntze	Lecythidaceae	Forest
<i>Antonia ovata</i> Pohl	Loganiaceae	Savanna
<i>Strychnos pseudoquina</i> A. St.-Hil.	Loganiaceae	Savanna
<i>Lafoensia pacari</i> A. St.-Hil.	Lythraceae	Savanna
<i>Byrsonima basiloba</i> A.Juss.	Malpighiaceae	Savanna
<i>Byrsonima coccolobifolia</i> Kunth	Malpighiaceae	Savanna
<i>Byrsonima intermedia</i> A.Juss.	Malpighiaceae	Generalist forest
<i>Byrsonima pachyphylla</i> A.Juss.	Malpighiaceae	Savanna
<i>Byrsonima verbascifolia</i> (L.) DC.	Malpighiaceae	Generalist savanna
<i>Heteropterys byrsonimifolia</i> A.Juss.	Malpighiaceae	Generalist savanna
<i>Ceiba speciosa</i> (A.St.-Hil.) Ravenna	Malvaceae	Forest
<i>Eriotheca candolleana</i> (K.Schum.) A.Robyns	Malvaceae	Forest
<i>Eriotheca globosa</i> (Aubl.) A.Robyns	Malvaceae	Generalist forest
<i>Eriotheca gracilipes</i> (K.Schum.) A.Robyns	Malvaceae	Savanna
<i>Eriotheca pubescens</i> (Mart. & Zucc.) Schott & Endl.	Malvaceae	Generalist savanna

Species	Family	Habitat preference
<i>Guazuma ulmifolia</i> Lam.	Malvaceae	Forest
<i>Luehea grandiflora</i> Mart.	Malvaceae	Generalist savanna
<i>Mollia lepidota</i> Spruce ex Benth.	Malvaceae	Forest
<i>Pseudobombax longiflorum</i> (Mart. & Zucc.) A.Robyns	Malvaceae	Generalist
<i>Bellucia grossularioides</i> (L.) Triana	Melastomataceae	Forest
<i>Miconia albicans</i> (Sw.) Steud.	Melastomataceae	Savanna
<i>Miconia latecrenata</i> (DC.) Naudin	Melastomataceae	Forest
<i>Miconia ligustroides</i> (DC.) Naudin	Melastomataceae	Generalist savanna
<i>Miconia pyrifolia</i> Naudin	Melastomataceae	Forest
<i>Miconia rubiginosa</i> (Bonpl.) DC.	Melastomataceae	Savanna
<i>Miconia</i> sp.	Melastomataceae	Forest
<i>Mouriri apiranga</i> Spruce ex Triana	Melastomataceae	Forest
<i>Mouriri elliptica</i> Mart.	Melastomataceae	Savanna
<i>Mouriri pusa</i> Gardner ex Gardner	Melastomataceae	Savanna
<i>Cabralea canjerana</i> (Vell.) Mart.	Meliaceae	Forest
<i>Cedrela fissilis</i> Vell.	Meliaceae	Forest
<i>Guarea guidonia</i> (L.) Sleumer	Meliaceae	Forest
<i>Guarea kunthiana</i> A.Juss.	Meliaceae	Forest
<i>Trichilia casaretti</i> C.DC.	Meliaceae	Forest
<i>Trichilia catigua</i> A.Juss.	Meliaceae	Forest
<i>Trichilia clausenii</i> C.DC.	Meliaceae	Forest
<i>Trichilia elegans</i> A.Juss.	Meliaceae	Forest
<i>Trichilia micrantha</i> Benth.	Meliaceae	Forest
<i>Trichilia pallida</i> Sw.	Meliaceae	Forest
<i>Mollinedia widgrenii</i> A.DC.	Monimiaceae	Forest
<i>Brosimum gaudichaudii</i> Trécul	Moraceae	Savanna
<i>Brosimum rubescens</i> Taub.	Moraceae	Forest
<i>Ficus eximia</i> Schott	Moraceae	Forest
<i>Pseudolmedia laevigata</i> Trécul	Moraceae	Forest
<i>Virola sebifera</i> Aubl.	Myristicaceae	Generalist
<i>Campomanesia adamantium</i> (Cambess.) O.Berg	Myrtaceae	Savanna
<i>Campomanesia xanthocarpa</i> (Mart.) O.Berg	Myrtaceae	Generalist forest
<i>Eugenia aurata</i> O.Berg	Myrtaceae	Savanna
<i>Eugenia blastantha</i> (O.Berg) D.Legrand	Myrtaceae	Forest
<i>Eugenia dysenterica</i> DC.	Myrtaceae	Generalist savanna
<i>Eugenia florida</i> DC.	Myrtaceae	Generalist forest
<i>Eugenia gemmiflora</i> O.Berg	Myrtaceae	Savanna
<i>Eugenia livida</i> O.Berg	Myrtaceae	Savanna
<i>Eugenia puniceifolia</i> (Kunth) DC.	Myrtaceae	Generalist savanna
<i>Eugenia ternatifolia</i> Cambess.	Myrtaceae	Savanna
<i>Eugenia uniflora</i> L.	Myrtaceae	Generalist forest
<i>Myrcia bella</i> Cambess.	Myrtaceae	Savanna
<i>Myrcia camapuanensis</i> N.Silveira	Myrtaceae	Savanna

Species	Family	Habitat preference
<i>Myrcia guianensis</i> (Aubl.) DC.	Myrtaceae	Savanna
<i>Myrcia lanuginosa</i> O.Berg	Myrtaceae	Savanna
<i>Myrcia multiflora</i> (Lam.) DC.	Myrtaceae	Generalist savanna
<i>Myrcia rimosa</i> Cambess.	Myrtaceae	Savanna
<i>Myrcia</i> sp.	Myrtaceae	Savanna
<i>Myrcia splendens</i> (Sw.) DC.	Myrtaceae	Generalist savanna
<i>Myrcia tomentosa</i> (Aubl.) DC.	Myrtaceae	Generalist forest
<i>Myrcia uberavensis</i> O.Berg	Myrtaceae	Savanna
<i>Myrcia variabilis</i> Mart. ex DC.	Myrtaceae	Savanna
<i>Myrcia venulosa</i> DC.	Myrtaceae	Generalist savanna
<i>Myrciaria delicatula</i> (DC.) O.Berg	Myrtaceae	Savanna
<i>Myrciaria floribunda</i> (H.West ex Willd.) O.Berg	Myrtaceae	Generalist forest
<i>Neomitranthes glomerata</i> (D.Legrand) Govaerts	Myrtaceae	Forest
<i>Plinia cauliflora</i> (Mart.) Kausel	Myrtaceae	Forest
<i>Psidium firmum</i> O.Berg	Myrtaceae	Savanna
<i>Psidium laruotteanum</i> Cambess.	Myrtaceae	Savanna
<i>Psidium</i> sp.	Myrtaceae	Forest
<i>Guapira graciliflora</i> (Mart. ex J.A.Schmidt) Lundell	Nyctaginaceae	Generalist savanna
<i>Guapira noxia</i> (Netto) Lundell	Nyctaginaceae	Savanna
<i>Neea theifera</i> Oerst.	Nyctaginaceae	Savanna
<i>Ouratea discophora</i> Ducke	Ochnaceae	Forest
<i>Ouratea hexasperma</i> (A. St.-Hil.) Baill.	Ochnaceae	Savanna
<i>Ouratea spectabilis</i> (Mart. ex Engl.) Engl.	Ochnaceae	Savanna
<i>Quiina florida</i> Tul.	Ochnaceae	Forest
<i>Heisteria ovata</i> Benth.	Olacaceae	Generalist savanna
<i>Minquartia guianensis</i> Aubl.	Olacaceae	Forest
<i>Agonandra brasiliensis</i> Miers ex Benth.	Opiliaceae	Generalist savanna
<i>Savia dictyocarpa</i> Müll.Arg.	Phyllanthaceae	Generalist forest
<i>Gallesia integrifolia</i> (Spreng.) Harms	Phytolaccaceae	Forest
<i>Cybianthus</i> sp.	Primulaceae	Forest
<i>Myrsine guianensis</i> (Aubl.) Kuntze	Primulaceae	Generalist forest
<i>Myrsine umbellata</i> Mart.	Primulaceae	Generalist
<i>Euplassa inaequalis</i> (Pohl) Engl.	Proteaceae	Generalist savanna
<i>Roupala montana</i> Aubl.	Proteaceae	Generalist
<i>Colubrina glandulosa</i> G.Perkins	Rhamnaceae	Forest
<i>Prunus myrtifolia</i> (L.) Urb.	Rosaceae	Generalist
<i>Amaioua guianensis</i> Aubl.	Rubiaceae	Generalist forest
<i>Cordia sessilis</i> (Vell.) Kuntze	Rubiaceae	Generalist
<i>Ferdinandusa rudgeoides</i> (Benth.) Wedd.	Rubiaceae	Forest
<i>Genipa americana</i> L.	Rubiaceae	Forest
<i>Tocoyena formosa</i> (Cham. & Schltdl.) K.Schum.	Rubiaceae	Savanna
<i>Metrodorea nigra</i> A. St.-Hil.	Rutaceae	Forest
<i>Zanthoxylum monogynum</i> A. St.-Hil.	Rutaceae	Forest
<i>Casearia gossypiosperma</i> Briq.	Salicaceae	Generalist forest

Species	Family	Habitat preference
<i>Casearia grandiflora</i> Cambess.	Salicaceae	Generalist forest
<i>Casearia lasiophylla</i> Eichler	Salicaceae	Generalist savanna
<i>Casearia sylvestris</i> Sw.	Salicaceae	Generalist
<i>Prockia crucis</i> P.Browne ex L.	Salicaceae	Forest
<i>Xylosma pseudosalzmanii</i> Sleumer	Salicaceae	Forest
<i>Cupania vernalis</i> Cambess.	Sapindaceae	Generalist forest
<i>Magonia pubescens</i> A. St.-Hil.	Sapindaceae	Generalist savanna
<i>Matayba guianensis</i> Aubl.	Sapindaceae	Generalist
<i>Matayba</i> sp.	Sapindaceae	Forest
<i>Chrysophyllum gonocarpum</i> (Mart. & Eichler ex Miq.) Engl.	Sapotaceae	Forest
<i>Micropholis venulosa</i> (Mart. & Eichler ex Miq.) Pierre	Sapotaceae	Generalist forest
<i>Pouteria ramiflora</i> (Mart.) Radlk.	Sapotaceae	Generalist
<i>Pouteria</i> sp.	Sapotaceae	Forest
<i>Pouteria torta</i> (Mart.) Radlk.	Sapotaceae	Generalist
<i>Schoepfia brasiliensis</i> A.DC.	Schoepfiaceae	Generalist
<i>Simarouba versicolor</i> A. St.-Hil.	Simaroubaceae	Generalist savanna
<i>Styrax ferrugineus</i> Nees & Mart.	Styracaceae	Savanna
<i>Styrax</i> sp.	Styracaceae	Forest
<i>Cecropia pachystachya</i> Trécul	Urticaceae	Generalist forest
<i>Urera baccifera</i> (L.) Gaudich. ex Wedd.	Urticaceae	Generalist forest
<i>Callisthene fasciculata</i> Mart.	Vochysiaceae	Generalist savanna
<i>Qualea cordata</i> Spreng.	Vochysiaceae	Generalist savanna
<i>Qualea grandiflora</i> Mart.	Vochysiaceae	Savanna
<i>Qualea multiflora</i> Mart.	Vochysiaceae	Savanna
<i>Qualea parviflora</i> Mart.	Vochysiaceae	Savanna
<i>Salvertia convallariodora</i> A. St.-Hil.	Vochysiaceae	Savanna
<i>Vochysia cinnamomea</i> Pohl	Vochysiaceae	Savanna
<i>Vochysia rufa</i> Mart.	Vochysiaceae	Savanna
<i>Vochysia tucanorum</i> Mart.	Vochysiaceae	Savanna
<i>Vochysia vismiifolia</i> Spruce ex Warm.	Vochysiaceae	Forest

Table S3 Number of species sampled to each site of seasonal forests and savannas environments in the Brazilian Cerrado. Density ≤ 3 represents the number and percentage of species with individual density equal or less than three individuals in each site, and density ≥ 4 denotes the number and percentage (in parentheses) of species with individual density equal or greater than four individuals in each site.

Site	Number of species	density ≤ 3	density ≥ 4
<i>Forest</i>			
Assis Ecological State - SP	39	31 (79)	8 (21)
Vassununga State Park - SP	31	18 (58)	13 (42)
Emas National Park - GO	21	15 (71)	6 (29)
Jataí – GO	29	12 (41)	17 (59)
Bacaba Park - MT	28	16 (57)	12 (43)
Ribeirão Cascalheira - MT	52	20 (38)	32 (62)
<i>Savanna</i>			
Assis Ecological State - SP	44	16 (36)	28 (64)
Vassununga State Park - SP	47	16 (34)	31 (66)
Emas National Park - GO	51	34 (67)	17 (33)
Jataí – GO	43	7 (16)	36 (84)
Bacaba Park - MT	90	15 (17)	74 (83)
Ribeirão Cascalheira - MT	66	24 (36)	42 (64)

Table S4 Percentage of occurrence in 376 areas evaluated by Ratter et al. (2003) and bark thickness of the 19 species that occurred in both seasonal forest and savanna habitats in the Brazilian Cerrado. Bold indicate the species that were more important (38 species) in Ratter et al. (2003), and threshold bark thickness of 5.9 mm in Hoffmann *et al.* (2012).

Species	Family	Percentage of occurrence in Ratter et al. (2003)	Bark thickness (mm)	
			Forest	Savanna
<i>Astronium fraxinifolium</i>	Anacardiaceae	47.3	0.7	0.60
<i>Tapirira guianensis</i>	Anacardiaceae	28.5	1.4	0.78
<i>Xylopia aromatica</i>	Annonaceae	49.2	5.9	3.8
<i>Hirtella glandulosa</i>	Chrysobalanaceae	16.5	7.6	2.0
<i>Buchenavia tomentosa</i>	Combretaceae	21.5	3.3	2.8
<i>Chaetocarpus echinocarpus</i>	Euphorbiaceae	1.0	2.7	1.6
<i>Copaifera langsdorffii</i>	Fabaceae	39.1	2.6	10.4
<i>Tachigali vulgaris</i>	Fabaceae	42.3	3.1	0.52
<i>Pseudobombax longiflorum</i>	Malvaceae	39.6	2.7	5.6
<i>Virola sebifera</i>	Myristicaceae	15.2	2.3	1.7
<i>Myrsine umbellata</i>	Primulaceae	5.3	1.0	0.9
<i>Roupala montana</i>	Proteaceae	57.4	8.2	10.4
<i>Prunus myrtifolia</i>	Rosaceae	0.8	1.3	1.3
<i>Cordia sessilis</i>	Rubiaceae	10.6	1.5	0.66
<i>Casearia sylvestris</i>	Salicaceae	58.0	1.1	14.9
<i>Matayba guianensis</i>	Sapindaceae	26.9	3.3	1.6
<i>Pouteria ramiflora</i>	Sapotaceae	55.9	5.4	9.9
<i>Pouteria torta</i>	Sapotaceae	24.2	3.1	13.8
<i>Schoepfia brasiliensis</i>	Schoepfiaceae	0	1.4	1.1




Figure S1 Mean values of functional traits to each group of habitat preference - specialist of forest (F) and savanna (S), and generalist sampled only in forest (FG) and sampled only in savanna (SG) - for the sites of seasonal forests and savannas in the Brazilian Cerrado.

CAPÍTULO II

Formatado nas normas da revista:

Journal of Vegetation Science

Edaphic properties drive functional trait patterns in savannas and seasonal forests plant communities of the Cerrado

Leandro Maracahipes, Marcos B. Carlucci & Marcus V. Cianciaruso

Key-words

acquisitive strategies, conservative strategies, soil nutrients, edaphic gradients, leaf nutrient content

Maracahipes, L. (lmaracahipes@gmail.com)¹

Cianciaruso, M.V. (corresponding author, cianciaruso@gmail.com)¹

Carlucci, M.B. (marcos.carlucci@gmail.com)^{1,2}

¹Departamento de Ecologia, Instituto de Ciências Biológicas, Universidade Federal de Goiás, Goiânia, Goiás, CP 131, 74001-970, Brazil.

²Departamento de Botânica, Setor de Ciências Biológicas, Universidade Federal do Paraná, CP 19031, Curitiba, PR 81531-980, Brazil.

Abstract

Question: Do gradients in soil fertility and toxicity drive shifts in density of individuals and community trait means in savannas and seasonal forests? Are communities with conservative traits associated with poor soils and communities with acquisitive traits associated with more fertile soils?

Location: Savannas and seasonal forests plant communities in the Cerrado phytogeographic domain, in Brazil.

Methods: We measured seven functional traits from 286 species occurring in 120 plots distributed in six forest and savanna sites throughout the Cerrado. In each vegetation, we recorded tree and shrub species and collected surface soil samples in 10 200-m² plots. We measured maximum plant height (Hmax), leaf area (LA), specific leaf area (SLA), leaf phosphorous content (LPC), stem-specific density (SSD), leaf thickness (LT), and leaf carbon-nitrogen ratio (C/N) to compute community-weighted means for each trait. We used linear regressions to model community trait means and density of individuals by edaphic gradients.

Results: We found lower density of individuals in more fertile soils in forests and savannas. In forests communities, our results suggest that an acquisitive strategy is related to toxicity (lower LA, SLA and PLC in more toxic soils) and fertility (high Hmax and LA in richer soils) gradients, and that a conservative strategy is related to toxicity gradient (high SSD and C/N in more toxic soils). In savanna communities, the patterns were less evident, with fewer traits indicating a strategy of resource acquisition (high LA in richer soils) or a strategy of resource conservation (high LT in toxic soil).

Conclusions: We brought evidence that forest and savanna species indeed adopt different strategies related to resource use in contrasting communities of the Cerrado

domain. We highlighted that density of individuals and community trait means shift along edaphic gradients in savannas and seasonal forest plant communities in the Cerrado domain. We show a stronger negative relationship of density of individuals with fertility and toxicity gradient, that could be result of dominance of some species with high competitive ability. For community trait mean, we demonstrate that in plots with fertile soils and less toxic have species with traits related to fast acquisitive strategy of resources, while in soil more toxic and with low fertility have species with traits related to conservative strategy.

Introduction

Understanding what are the processes and factors determining ecological strategies plants use to respond to abiotic and biotic factors and their consequences to community composition, structure and functioning is a fundamental goal in vegetation science (McGill et al. 2006; Vellend 2010). In order to coexist, species often adopt different strategies of growth and reproduction (Oliveras & Malhi 2016) leading to a trade-off between rapid acquisition and resource conservation strategies (Diaz et al. 2004; Poorter 2009). Such a trade-off may be assessed by evaluating the distribution of traits along environmental gradients at the community scale (Goodland & Pollard 1973; Furley & Ratter 1988; Ordoñez et al. 2009; Baxendale et al. 2014). Therefore, exploring the role of environmental gradients on different traits (Lavorel & Garnier 2002; Carvalho & Batalha 2013; Wigley et al. 2016) is key to understanding vegetation dynamics in contrasting environments such as forests and savannas.

Several studies have highlighted that edaphic properties strongly influence plant community structure (Wright et al. 2004; de Bello et al. 2006; Dantas & Batalha 2011; Zava & Cianciaruso 2014; Laureto & Cianciaruso 2015; Barros et al. 2015; Moraes et

al. 2015). Functional traits influence directly the responses of plants to the environment as well as plant-plant interactions (Díaz & Cabido 2001; Lavorel & Garnier 2002). Soil properties can determine the performance and distribution of functional traits, directly affecting the growth and reproduction of plants (Wright et al. 2004; Swenson & Enquist 2007; Freschet et al. 2011; Moles et al. 2014). Low stem density, high maximum height, high specific leaf area, and leaves of high nutritional quality characterize plants with an acquisitive strategy, typical of nutrient-rich environments. On the other hand, plants with a conservative strategy, typical nutrient-poor environments, can be characterized by thick and tough leaves, low specific leaf area, and leaves of low nutritional quality (Wright et al. 2004; Diaz et al. 2004; Ordoñez et al. 2009). Therefore, functional trait-based approaches enable us to evaluate the responses of plants to environmental conditions and interactions among species (Lavorel & Garnier 2002; Götzenberger et al. 2012).

One way to evaluate the dominant strategies in plant communities, is with community trait means weighted by abundance (CWM, Garnier et al. 2004). Ecosystem processes are mainly determined by dominant species (Grime 1998; Garnier et al. 2004), thus, CWM is useful to account the effect of dominant species on environmental change and ecosystem processes (Garnier et al. 2004; Pakeman & Quested 2007; Garnier et al. 2007; Garnier & Navas 2012). Because it is expected that CWM vary along environmental gradients (Diaz et al. 2004; Garnier et al. 2004) this approach should allow us to understand how changes in the functional composition along of environmental gradients reflect in acquisitive and conservative strategies in forest and savanna communities. Furthermore, trait mean values can provide insights on different trait compositions across environmental gradients.

In Neotropical savannas different habitats, such as savannas and seasonal forests, occur side by side under the same climate (Oliveras & Malhi 2016). The occurrence of such contrasting environments under the same climatic conditions has been identified as an effect of spatial variation in soil nutrient availability (Ruggiero et al. 2002). Several nutrients play important roles in the structure and composition of these communities (Furley & Ratter 1988; Ruggiero et al. 2002; Dantas & Batalha 2011; Pellegrini 2016). In savannas, an important determinant of the structure and composition of communities is related to the ability of acquisition of soil nutrients and water by plants (Oliveira et al. 2005). Many savanna species have a root system that enable them to extract nutrients and water from great depths (Goedert 1983). While high soil aluminum concentration may inhibit nutrients absorption by plants, being toxic for many forest species, savanna species are usually adapted to high levels of aluminum and some are even aluminum accumulators (Furley & Ratter 1988). Soil moisture, fire, and competition for nutrients determine plant density in savannas (Furley & Ratter 1988; Marimon Junior & Haridasan 2005). High frequency of fire can turn savannas into grasslands, while fire exclusion may lead to the development of seasonal forests (Hoffmann et al. 2012). On the other hand, forest habitats often have higher soil nutrient availability that favours tree growth, because plants with strategy of fast growth can avoid light limitation in the understory (Ruggiero et al. 2002; Hoffmann et al. 2012; Oliveras & Malhi 2016). Moreover, resource availability together with disturbance regimes have been appointed as a determining factor for the occurrence of forest and savanna habitats under same climate (Dantas et al. 2016).

We answered the following questions to each study habitat (forest and savanna):

- (i) Do gradients in soil fertility and toxicity reduce the number of individuals in seasonal forests and savannas?
- (ii) Do gradients in soil fertility and toxicity drive shifts

in community trait means? Our expectation is that communities with higher soil fertility and lower toxicity will have higher mean values of maximum plant height (Hmax), leaf area (LA), specific leaf area (SLA), leaf phosphorous content (LPC), but lower mean values of stem-specific density (SSD), leaf thickness (LT) and leaf carbon-nitrogen ratio (C/N), reflecting a dominance of species with rapid acquisition strategy (Westoby 1998; Pérez-Harguindeguy et al. 2013; Wigley et al. 2016). On the other hand, communities in soils with low fertility and high toxicity, we expect to observe lower mean values of Hmax, LA, SLA and LPC, but high mean values of SSD, LT and C/N, reflecting a conservative strategy (Table 1). Furthermore, forest environments with high fertility and low toxicity may promote the establishment of species with fast growth and better competitive ability, reducing the number of individuals by competitive exclusion (Mayfield & Levine 2010; Dantas & Batalha 2011; Carvalho & Batalha 2013).

Table 1. Functional significance of traits and trait-based patterns predicted in relation to edaphic gradients in forests and savannas of the Cerrado.

Trait	Acronym	Functional significance	Prediction	References
Maximum plant height	Hmax	Competitive plant fitness, plant fecundity, tolerance or resistance to disturbances.	We expected that trees in more fertile soil will be taller. Considering that soils richer in nutrients support higher plant biomass, the variation of Hmax along soil gradients may be related to a trade-off between resource conservation and competitive ability.	(Westoby 1998; Pérez-Harguindeguy et al. 2013; Laureto & Cianciaruso 2015)
Leaf area	LA	Strategies of resource acquisition and use. Ecological strategy adopted with respect to environmental nutrients stress and disturbances.	Leaf area has consequences for the leaf energy and water balance. We expected that plants present high leaf area in more fertile habitats than infertile habitats. The slower growth of plants on infertile soils can favor a smaller investment in leaf area, which minimizes water loss.	(McDonald et al. 2003; Pérez-Harguindeguy et al. 2013)
Specific leaf area	SLA	Related to structural defense, resource uptake and use efficiency and growth strategies; negatively correlated with leaf lifespan.	We expected that communities in resource-rich environments have higher SLA than in poor habitats. This reflects the trade-off between fast growth and nutrient conservation, and is related with leaf nitrogen concentration. This indicate that environments with fertile soils provide rapid resource capture and high relative growth, while the opposite is expected for a conservative strategy.	(Wright et al. 2004; Pérez-Harguindeguy et al. 2013)
Leaf phosphorus content	LPC	Phosphorous is a limiting resource related to growth and productivity in plants.	We expected that plants in more fertile soils show higher nutrients concentrations in their leaves. Fertile soils are generally associated with high nutritional quality and plant growth due to higher soil phosphorous availability.	(Schachtman et al. 1998; Lloyd et al. 2009; Ordoñez et al. 2009; Pérez-Harguindeguy et al. 2013)
Stem-specific density	SSD	Structural vigor, and consequent resistance to physical damage.	We expected to find a negative relationship between SSD and soil fertility. This relation is associated to the growth-survival trade-off, in which low stem density (with large vessels) leads to a fast growth, because of large hydraulic capacity and conductive tissue is less expensive to construct, whereas a high stem density (with small vessels) leads to a high survival by providing a stronger defense against physical and biological damage.	(Baker et al. 2004; Chave et al. 2006; Lloyd et al. 2009; Chave et al. 2009; Pérez-Harguindeguy et al. 2013)
Leaf thickness	LT	Related to species strategies of resource acquisition and use, and with resistance to physical damage, correlated with leaf toughness.	LT is one of the key components of SLA and photosynthetic capacity, and predict that LT should be higher in less fertile habitats, because balancing photosynthetic benefits against C costs of respiration and transpiration. Besides, LT indicates the largest relative carbon investment in structural protection of photosynthetic tissues.	(Vile 2005; Pérez-Harguindeguy et al. 2013)
Leaf carbon-nitrogen ratio	C/N	Nutritional quality and palatability of plant, leaf decomposition rate, N is the major limiting macronutrient in plants.	We expected a negative relationship between C/N ratio and soil fertility. In less-fertile soils high values of this ratio suggest a resource conservation strategy. Moreover, more fertile soils tend to present soil organic matter, and this might enhance the nitrogen absorption by plants, thereby increasing the nitrogen content in the leaves of plants in more fertile soils.	(Ordoñez et al. 2009; Pérez-Harguindeguy et al. 2013; Apaza-Quevedo et al. 2015)

Material and Methods

Study region

The Cerrado is a phytogeographical domain characterized by complex vegetational mosaics (Ribeiro & Walter 2008) composed by savannas, seasonal forests and tropical grasslands (Batalha 2011; Dantas et al. 2016). The Cerrado receives strong floristic influence of the neighboring phytogeographical domains of Amazônia, Mata Atlântica and Caatinga (Méio et al. 2003; Souza-Neto et al. 2016). The great variety of habitats, ranging from grasslands to forests, can result from the diverse climatic conditions, edaphic properties and fire frequency registered throughout the phytogeographical domains (Oliveira-Filho & Ratter 2002; Dantas et al. 2013).

Sampling design

We surveyed six regions located throughout the Cerrado domain in order to capture a great range of phytogeographical variation. We adopt a paired design sampling one savanna (cerrado *stricto sensu*) and one forest (seasonally dry forest) site in each region (Fig. 1). Average distance between sites was 12 km, and the latitudinal range varied from 12°S to 22°S (Fig. 1 and Table S1). The smallest distance between regions was 150 km (between Emas National Park and Jataí), while the smallest distance within pairs of forest and savanna blocks was 1 km (in Emas National Park). In general, the mean values and coefficient of variation of the edaphic characteristics varied little between the plots of either forests or savannas at each site (Table S2).




Figure 1. Location of six study regions with seasonal forest and savanna sites in the Brazilian Cerrado. 1. Assis Ecological Station (lat/long: 22°35'S/50°22'W), 2. Vassununga State Park (21°36'S/47°37'W), 3. Emas National Park (17°54'S/52°59'W), 4. Jataí (17°45'S/51°32'W), 5. Bacaba Park (14°42'S/52°21'W), 6. Ribeirão Cascalheira (12°49'S/51°46'W).

In each savanna and forest site we sampled all tree- and shrubs in 10 100-m² plots. In savanna plots we recorded all individuals with diameter at ground height ≥ 5 cm, whereas in forest plots all individuals with diameter at breast height ≥ 10 cm were sampled. All individuals were identified following the Angiosperm Phylogeny Group

classification (APG-IV 2016). Additionally, we revisited the species names in the Brazilian Flora (Forzza et al. 2012).

Trait sampling

We measured the following traits at species-level according to Pérez-Harguindeguy et al. (2013): maximum plant height (Hmax), leaf area (LA), specific leaf area (SLA), leaf nutrient content of phosphorous (LPC), leaf thickness (LT), stem specific density (SSD) and carbon/nitrogen ratio (C/N) (see Table 1 for their ecological significance). Whenever possible we measured 10 individuals for each species in each site. When a given species had less than 10 individuals in our plots, we sampled individuals that were nearby the plot. When that was not possible we included all individuals that were available (see Table S3). LPC and C/N were measured for three individuals of each species in each site. To obtain the values of leaf traits, first we scanned five leaves of each individual, and posteriorly calculated area leaf through of a script that we build using the “EBImage” package (Pau et al., 2014) with R 3.2.1 (<http://www.R-project.org>). After scanning, the leaves were dried at 80 °C for 48 h to acquire leaf dry mass. Other five leaves were utilized to calculate the leaf thickness values using a digital micrometer. All traits were collected during the rainy season (December to March 2011-2012), which represents the period of maximum leaf expansion and maturity and growth and reproductive peak for most of the studied plants (Lenza & Klink 2006).

Edaphic variables

Composite soil samples (0-10 cm depth) were collected in all forest and savanna plots (three samples per plot). We obtained the following soil variables (Embrapa 1999): clay

concentration, pH, soil organic matter (SOM), phosphorus concentration (P), nitrogen concentration (N), exchangeable contents of K^+ and Al^{3+} .

Spatial autocorrelation

Our aim was to evaluate the role of fertility gradient on community trait means of plant communities. Communities at short distances from one another tend to be more similar than distant communities (Hawkins 2012) due to shared historical processes and spatially-structured biological processes such as dispersal and species interactions (Peres-Neto & Legendre 2010). Therefore, response variables such as community trait means may be structured in the space, which affect statistical assumptions such as independence of residuals (Legendre 1993).

In order to take the spatial autocorrelation into account in our analysis we used the method of principal coordinates of neighbourhood matrices (PCNM) (Borcard & Legendre 2002). The PCNM is based on a PCoA of geographical coordinates truncated by distance in order to generate axes (spatial filters) representing vectors of spatial variation at increasing scales. Here, we used the spatial filters as explanatory variables together with the environmental predictors to minimize the residual autocorrelation (Rangel et al. 2006). These analyses were conducted at the plot level and the truncation distance that represent the maximum distance connecting all communities, which defines the neighbourhood boundaries (Rangel et al. 2006), was 5.5 km for forest plots and 2.5 km for savanna plots. We conducted the PCNM analyses in software SAM 4.0 (Rangel et al. 2006).

Statistical analyses

All analyses were performed separately at the plot level for each of communities of forests (60 plots) and savannas (60 plots). To reduce the dimensionality of edaphic parameters we computed principal component analysis (PCA). The number of components retained for further analyses was determined on the basis of the broken stick criteria, in which the eigenvalue must be greater than the values of broken stick eigenvalue (Frontier 1976). Moreover, we evaluated the effect of each variable on each principal component (PC) through bootstrap randomization in the software Past (Hammer et al. 2001). These analyses were conducted separately for forest (Fig. S1) and savanna plots (Fig. S2).

Initially, we seek to understand if the fertility and toxicity gradients determine the density of individuals of the communities of forest and savannas. We related the density of individuals with edaphic gradients controlling the spatial autocorrelation of predictors variables through spatial filters.

In order to evaluate shifts of trait means at the community level along the gradient, we computed community-weighted trait means (CWM), defined as the mean traits values present in the community weighted by the relative abundance of species (Garnier et al. 2004).

To investigate the shifts of community trait means along the edaphic gradient, we fitted a linear regression analysis (ordinary least squares - OLS) with PC1, PC2 and PCNM (spatial filters) as predictors variables (Rangel et al. 2006; Peres-Neto & Legendre 2010). Spatial filters were used as predictor variables to account for spatial autocorrelation in residuals. When necessary, we used an algorithm that added the spatial filter that accounted for the most of the spatial autocorrelation in the residuals remaining in the model, repeating this process until the autocorrelation in the residuals

reached its minimum (Rangel et al. 2010; Peres-Neto & Legendre 2010). All response variables of community trait means were spatially structured. Thus, we accounted for spatial autocorrelation in the residuals of the models by including spatial filters as predictors (Table 2 and Appendix 1).

Results

Environmental gradients - For the forests, the first two principal components (PC) together accounted for 70.4% of the variance in the edaphic variables (Fig. 2). PC1 accounted for 42.3% of the explained variance and represents mainly a soil toxicity gradient, characterized mostly by Al and low pH (Fig. 2 and S1A). PC2 accounted for 28.1% of the explained variance and represents a soil fertility gradient, characterized mostly by SOM and clay (Fig. 2 and S1B). For savannas, the first two principal components together accounted for 54.5% of the variance in the edaphic variables (Fig. 3). The PC1, accounted for 30.3% explained variance and represented a soil fertility, characterized mostly by clay and SOM (Fig. 3 and S2A). On the other hand, PC2 accounted for 24.2% of the explained variance and represented a toxicity gradient, characterized by pH and Al (Fig. 3 and S2B).




Figure 2. Distribution of the plots and sites of seasonal forests along the two axes of a principal component analysis (PCA) based on ten edaphic properties. Each principal component represents a soil gradient: PC1, toxicity gradient, and PC2, fertility gradient. Different symbols and colors represent different sites.




Figure 3. Distribution of the plots and sites of savannas along the two axes of a principal component analysis (PCA) based on ten edaphic properties. Each principal component represents a soil gradient: PC1, fertility gradient, and PC2, toxicity gradient. Different symbols and colors represent different sites.

Density of individuals – The density of individuals decreased along the toxicity and fertility gradients in the forests (Table 2, Fig. S3). In savannas, the density of individuals decreased only with fertility, however spatial filters showed high explanatory power for this model (Table 2, Fig. S3).

Community trait means – In seasonal forests, considering species with traits with acquisitive strategy, only maximum height increased along the toxicity gradient, while leaf area, specific leaf area and leaf phosphorous content decreased along the same gradient (Table 2, Fig. S4). On fertility gradient, maximum height and leaf area increased along the fertility gradient, while specific leaf area and leaf phosphorous

content decreased along the same gradient (Table 2, Fig. S4). Considering species with traits with conservative strategy, stem-specific density and carbon/nitrogen ratio increased along the toxicity gradient (Table 2, Fig. S4). While, stem-specific density, leaf thickness and carbon/nitrogen ratio also increased along the fertility gradient (Table 2, Fig. S4).

In savannas, the fertility gradient was more important than toxicity to community trait mean. Only leaf thickness increased along the toxicity gradients considering conservative strategy (Table 2, Fig. S5). Considering traits with acquisitive strategy, maximum height increased along the fertility gradient, while leaf area decreased along the same gradient (Table 2, Fig. S5). While considering conservative strategy, only leaf thickness increased along the fertility gradient (Table 2, Fig. S5). The others traits were significant with edaphic gradient (Table 2, Fig. S5).

Table 2. Models of linear regression analysis for toxicity and fertility gradients in seasonal forests and savannas in the Brazilian Cerrado. Spatial filters (F) were computed using principal coordinates of neighbourhood matrices (PCNM; Peres-Neto & Legendre 2010) and were included in each model as predictors to account for spatial autocorrelation in the residuals. The variables toxicity and fertility are principal component from PCA computed separately for seasonal forest and savannas. Numbers stressed in bold indicate significant regression coefficients (β) with $p < 0.05$ and with explanatory power of predictors ≥ 0.19 .

	Model	toxicity - β	fertility - β	R^2	P model	predictors	shared	filters
Density of individuals								
<i>Forest</i>	density ~ toxicity + fertility + F1 + F5	-0.31	-0.73	0.52	0.001	0.45	-0.09	0.18
<i>Savanna</i>	density ~ fertility + toxicity + F2 + F3 + F4 + F5 + F6	0.11	-0.69	0.80	0.001	0.27	0.16	0.40
Community-weighted means								
<i>Forest - acquisitive strategy</i>	Hmax ~ toxicity + fertility + F1 + F2	0.60	0.34	0.36	0.001	0.23	-0.07	0.23
	LA ~ toxicity + fertility + F1	-0.35	0.33	0.38	0.001	0.23	-0.04	0.21
	SLA ~ toxicity + fertility + F3	-0.48	-0.25	0.64	0.001	0.29	0.20	0.16
	LPC ~ toxicity + fertility + F3 + F4 + F5	-0.58	-0.21	0.91	0.001	0.27	0.44	0.21
<i>Forest - conservative strategy</i>	SSD ~ toxicity + fertility + F3	0.44	0.15	0.82	0.001	0.21	0.36	0.26
	LT ~ toxicity + fertility + F1 + F2 + F3	0.27	0.73	0.52	0.001	0.43	-0.17	0.29
	C/N ~ toxicity + fertility + F2 + F3	0.71	0.59	0.75	0.001	0.48	0.16	0.11
<i>Savanna - acquisitive strategy</i>	Hmax ~ fertility + toxicity + F2 + F3 + F5	-0.16	-0.71	0.84	0.001	0.45	0.19	0.21
	LA ~ fertility + toxicity + F2 + F4	-0.15	0.43	0.51	0.001	0.19	0.06	0.29
	SLA ~ fertility + toxicity + F2 + F4 + F5	-0.05	-0.38	0.81	0.001	0.13	0.18	0.51
	LPC ~ fertility + toxicity + F2 + F3 + F5	-0.04	0.16	0.90	0.001	0.02	0.16	0.74
<i>Savanna - conservative strategy</i>	SSD ~ fertility + toxicity + F1	-0.29	0.14	0.74	0.001	0.10	0.40	0.25
	LT ~ fertility + toxicity + F2 + F4 + F5	0.22	0.53	0.86	0.001	0.28	0.30	0.29
	C/N ~ fertility + toxicity + F2 + F3	-0.01	0.02	0.86	0.001	0.01	0.25	0.63

Discussion

In general, our results support that forest and savanna plants have different strategies related to rapid acquisition and conservative strategy of resources. We showed that communities with more toxic soil determined lower density of individuals in forest. In contrast, more fertile soils determined lower density of individuals in forest and savanna environments. For community trait mean, traits values were more sensible to shift in fertility gradient than toxicity gradient, mainly in forest environments. These different strategies (rapid acquisitive and conservative) observed are responses of traits in relation to resources acquisition for establish and develop in environments with different light and soil resources availability as seasonal forest and savanna (Reich et al. 1999; Diaz et al. 2004; Svenning et al. 2004; Apaza-Quevedo et al. 2015; Wigley et al. 2016).

Density of individuals

Density of individuals decreased with the increasing in soil fertility and toxicity in forest communities. This could be related to high competitive ability of some species, that on fertile soils become dominant producing a larger canopy cover, avoiding the establishment or excluding others species (Carvalho & Batalha 2013; Martorell et al. 2015). In theory, a greater resource available in the soil tend to promote species coexistence through resources partitioning (Barot 2004; Dantas & Batalha 2011), as even forest species are adapted to experience habitats with certain nutrient limitation conditions (Furley & Ratter 1988), which would lead sites with more richer soils for have high competitive effect resulting in low density of individuals. Neri *et al.* (2012) also showed a negative relationship between density of individuals and aluminium in the soil from savanna-woodland. Al is the main element indicator of toxicity in our

study system (Goodland & Pollard 1973) because it compete with other essential elements for absorption sites (Wagatsuma 1983). Thus, even with many Al-tolerant species, in general the aluminium toxicity may act on establishment of species, determining a lower density of plants.

Community-level trait shifts

Generally, higher values of leaf nutrient content, SLA, large leaves (Ruggiero et al. 2002; Carvalho & Batalha 2013; Wigley et al. 2016), and tall plants (Westoby 1998; Schamp & Aarssen 2009) grow fast and have a great competitive ability, thus being related to rapid acquisition of resources. On the other hand, large values of wood density, leaf thickness and C/N ratio are related to conservative strategy, protection against fire and water loss (Table 1) (Baker et al. 2004; Hoffmann et al. 2005; Chave et al. 2006; Wigley et al. 2016). Based on such strategies, our expectations were that the soil fertility and toxicity would influence the trait mean values along these gradients (Ruggiero et al. 2002; Silva et al. 2010; Carvalho & Batalha 2013; Baxendale et al. 2014; Wigley et al. 2016). However, our results showed that trait related to acquisitive strategies tended to have larger values in more benign environments (high fertility and low toxic soil), while trait related for conservative strategies tended to have larger values in more stressful environments (low fertility and high toxic soil). Additionally, our results showed that these strategies were more evident in the seasonal forests than on the savannas of Cerrado.

In general, we demonstrate that seasonal forest in richer soils have higher leaf phosphorous content and tall plants, while that low leaf area, specific leaf area and leaf phosphorous content, high woody density, carbon/nitrogen ratio and leaf thickness in soils with higher toxicity. Likewise, the savanna plots with richer soils have higher leaf

area, while that high leaf thickness in toxic soils. These findings are according for the fast acquisitive strategy of resources adopted by plants in conditions of large resource available and lesser environmental stress (Diaz et al. 2004; Wigley et al. 2016; Pellegrini 2016). This is possible because more fertile soils support higher plant biomass due to higher resource available, making possible that species with strategy of fast growing, that consequently have larger nutrient demand, may have an investment in traits related to fast growth (Chapin III et al. 1993; Westoby 1998; Pellegrini 2016), and produce litter with high-quality, that is quickly recycled (Baxendale et al. 2014).

In fact, the nutrient available is an important environmental driver involving the rapid acquisition of nutrients and conservation of resources in plant, determining functional traits and community composition (Wright et al. 2004; Lloyd et al. 2009; Ordoñez et al. 2009; Baxendale et al. 2014; Wigley et al. 2016; Oliveras & Malhi 2016). Plants that explore water- and nutrient stress tend to have convergent traits that indicate better use and conservation of resources, promoting defences and long leaf lifespans (Chapin III et al. 1993; Diaz et al. 2004), and are represented by small, thick and tough leaves, low SLA and leaf nutrient content, and lower plant (Diaz et al. 2004; Pérez-Harguindeguy et al. 2013), and high woody density, leaf thickness and carbon/nitrogen ratio (Table 1). Strikingly, for some leaf traits and wood density the results were opposite to our predictions. We registered low SLA and LPC and high SSD, LT and C/N (in forest), and lower plants and high LT (in savannas) on more fertility soils. These results appoint to conservative strategy on fertile soils, in which normally is expected strategy related to rapid acquisition of resources (Diaz et al. 2004; Liu et al. 2012; Jager et al. 2015; Wigley et al. 2016). We expected that communities in resource-rich environments would have higher SLA than in poor habitats (Knops & Reinhart 2000; Wright et al. 2004). However, our results were the opposite and can be related to

access to light in forests because there is evidence that plants decrease SLA as they increase in height (in order to reach the canopy where light is fully available) (Fahey et al. 1998; Klooster et al. 2007). Thus, more richer soils promoting higher growth of plants, as seen in maximum height in forest, leading to greater light available in taller plants. Like the solar radiation is intense during most of the year (Miranda et al. 1997), when plants reaches the top of the canopy they adopt strategies that prevents water loss altering yours morphological characteristics (low SLA, high LT and C/N). Some studies have showed that SLA generally increase with increasing foil fertility in response to soil nutrient availability and related to rapid acquisition strategy (Ordoñez et al. 2009; Wang et al. 2012; Li et al. 2015).

Plant height is related with the nutrient transport capacity, and maximum plant height is limited by hydraulic and mechanical constraints (Koch et al. 2004; Niklas 2007), moreover the costs with nutrients transports in taller trees can resulting reduction in leaf photosynthesis (Ryan & Yoder 1997). Thus, we expected that plants in less productive and more toxic environment tend to be smaller (Laureto & Cianciaruso 2015; Sfair et al. 2016). Our results indicate that soil more fertile and less toxic promotes in taller plants in forest and savanna, which can reflect in high competitive vigor and tolerance or avoidance of environmental stress (as fire) (Westoby 1998). However, in this study the poor soil supported tall trees on savannas, this may be related to disturbance regime (fire) in savannas (Jager et al. 2015). In savanna communities, the grass are often tall, but in richer soils can accumulate large herbaceous layer increasing the frequency and intensity of fires and decreasing the plant height (Dantas, Pausas, et al. 2013; Dantas, Batalha, et al. 2013), because fertile soils can result in more heterogeneous canopy height (Jager et al. 2015). On the other hand, poor soils can result in more varied plant height, thus fire events often cause damage mainly on herbaceous

layer, favoring the taller plants (Dantas, Pausas, et al. 2013; Jager et al. 2015). Likewise, seasonal forest registered taller plant on more toxicity soil, and its can be negatives effects from fragmentation, because the sites of seasonal forest from São Paulo State showed smaller toxicity and presented small plants, since that fragmentation makes the forest more susceptible to falling trees (Ferreira & Laurance 1997).

Conclusion

We highlighted that density of individuals and community trait means shift along edaphic gradients in savannas and seasonal forest plant communities in the Cerrado domain. We show a stronger negative relationship of density of individuals with fertility and toxicity gradient, that could be result of dominance of some species with high competitive ability. For community trait mean, we showed that the changes were more clear on fertility gradient than toxicity gradient, and that forest environments was more sensible to shift in both gradients than savannas. We demonstrate that in plots with fertile soils and less toxic have traits related to fast acquisitive strategy of resources reflecting in traits related to fast growth competitive ability, and that the plants adopt this strategy in conditions of large resource available and lesser environmental stress (Diaz et al. 2004; Wigley et al. 2016; Pellegrini 2016). On the other hand, soil more toxic and with low fertility have species with traits related to conservative strategy, indicating better use and conservation of resources and promoting defences and long leaf lifespans (Diaz et al. 2004; Pérez-Harguindeguy et al. 2013). However, the relation of some traits with the edaphic gradient were contrary to our expeditions, mainly on fertility gradient.

Acknowledgments

This research was supported by CNPq-Brazil (#563621/2010-9, #478747/2009-8 and PELD - SITE 13) and FAPEG-GO (#201110267000130/31-10). L.M. was supported by scholarships from Capes-Brazil, M.V.C. received PQ fellowships from CNPq (#999999/999-9) and M.B.C received a postdoctoral fellowship from Capes-Brazil (PNPD ##1454013). We thank to all field team responsible for sampling data: Edmar Oliveira, Leonardo Maracahipes, Letícia Gomes, Livia Laureto, Danira Padilha, Mônica Forsthofer, Mariângela Abreu, Josias Santos and Fernando Ribeiro. We also thank to taxonomists Geraldo Franco, Osnir Aguiar, João Baitelo, Natália Ivanauskas and Renato Mello-Silva for the identification of some species.

References

- Apaza-Quevedo, A., Lippok, D., Hensen, I., Schleuning, M., & Both, S. 2015. Elevation, topography, and edge effects drive functional composition of woody plant species in tropical montane forests. *Biotropica* 47: 449–458.
- APG-IV. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181: 1–20.
- Baker, T.R., Phillips, O.L., Malhi, Y., Almeida, S., Arroyo, L., Di Fiore, A., Erwin, T., Killeen, T.J., Laurance, S.G., Laurance, W.F., Lewis, S.L., Lloyd, J., Monteagudo, A., Neill, D.A., Patino, S., Pitman, N.C.A., M. Silva, J.N., & Vasquez Martinez, R. 2004. Variation in wood density determines spatial patterns in Amazonian forest biomass. *Global Change Biology* 10: 545–562.
- Barot, S. 2004. Mechanisms promoting plant coexistence: can all the proposed processes be reconciled? *Oikos* 106: 185–192.
- Barros, M.J.F., Silva-Arias, G.A., Fregonezi, J.N., Turchetto-Zolet, A.C., Iganci, J.R.V., Diniz-Filho, J.A.F., & Freitas, L.B. 2015. Environmental drivers of diversity in Subtropical Highland Grasslands. *Perspectives in Plant Ecology, Evolution and Systematics*. doi: 10.1016/j.ppees.2015.08.001
- Batalha, M.A. 2011. O cerrado não é um bioma. *Biota Neotropica* 11: 21–24.
- Baxendale, C., Orwin, K.H., Poly, F., Pommier, T., & Bardgett, R.D. 2014. Are plant-soil feedback responses explained by plant traits? *New Phytologist* 204: 408–423.
- de Bello, F., Lepš, J., & Sebastià, M.-T. 2006. Variations in species and functional plant diversity along climatic and grazing gradients. *Ecography* 29: 801–810.

- Borcard, D., & Legendre, P. 2002. All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices. *Ecological Modelling* 153: 51–68.
- Carvalho, G.H., & Batalha, M.A. 2013. The drivers of woody species richness and density in a Neotropical savannah. *Biology Letters* 9: 1–4.
- Chapin III, F.S., Autumn, K., & Pugnaire, F. 1993. Evolution of suites of traits in response to environmental stress. *The American Naturalist* 142: S78–S92.
- Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G., & Zanne, A.E. 2009. Towards a worldwide wood economics spectrum. *Ecology Letters* 12: 351–366.
- Chave, J., Muller-Landau, H.C., Baker, T.R., Easdale, T. a, ter Steege, H., & Webb, C.O. 2006. Regional and phylogenetic variation of wood density across 2456 Neotropical tree species. *Ecological applications : a publication of the Ecological Society of America* 16: 2356–67.
- Dantas, V.D.L., & Batalha, M.A. 2011. Vegetation structure: Fine scale relationships with soil in a cerrado site. *Flora - Morphology, Distribution, Functional Ecology of Plants* 206: 341–346.
- Dantas, V.L., Batalha, M.A., & Pausas, J.G. 2013. Fire drives functional thresholds on the savanna–forest transition. *Ecology* 94: 2454–2463.
- Dantas, V. de L., Hirota, M., Oliveira, R.S., & Pausas, J.G. 2016. Disturbance maintains alternative biome states (M. Rejmanek, Ed.). *Ecology Letters* 19: 12–19.
- Dantas, V. de L., Pausas, J.G., Batalha, M.A., Loiola, P. de P., & Cianciaruso, M.V. 2013. The role of fire in structuring trait variability in Neotropical savannas. *Oecologia* 171: 487–494.
- Díaz, S., & Cabido, M. 2001. Vive la différence : plant functional diversity matters to ecosystem processes. 16: 646–655.
- Díaz, S., Hodgson, J.G., Thompson, K., Cabido, M., Cornelissen, J.H.C., Jalili, A., Montserrat-Martí, G., Grime, J.P., Zarrinkamar, F., Asri, Y., Band, S.R., Basconcelo, S., Castro-Díez, P., Funes, G., Hamzehee, B., Khoshnevi, M., Pérez-Harguindeguy, N., Pérez-Rontomé, M.C., Shirvany, F.A., Vendramini, F., Yazdani, S., Abbas-Azimi, R., Bogaard, A., Boustani, S., Charles, M., Dehghan, M., de Torres-Espuny, L., Falczuk, V., Guerrero-Campo, J., Hynd, A., Jones, G., Kowsary, E., Kazemi-Saeed, F., Maestro-Martínez, M., Romo-Díez, A., Shaw, S., Siavash, B., Villar-Salvador, P., & Zak, M.R. 2004. The plant traits that drive ecosystems: evidence from three continents. *Journal of Vegetation Science* 15: 295–304.
- Embrapa. 1999. *Sistema brasileiro de classificação de solos*. Embrapa Produção de Informação, Rio de Janeiro.
- Fahey, T.J., Battles, J.J., & Wilson, G.F. 1998. Responses of early successional northern hardwood forests to changes in nutrient availability. *Ecological Monographs* 68: 183–212.
- Ferreira, L. V., & Laurance, W.F. 1997. Effects of Forest Fragmentation on Mortality and Damage of Selected Trees in Central Amazonia. *Conservation Biology* 11: 797–801.
- Forzza, R.C., Baumgratz, J.F.A., Bicudo, C.E.M., Canhos, D.A.L., Carvalho, A.A., Coelho, M.A.N., Costa, A.F., Costa, D.P., Hopkins, M.G., Leitman, P.M.,

- Lohmann, L.G., Lughadha, E.N., Maia, L.C., Martinelli, G., Menezes, M., Morim, M.P., Peixoto, A.L., Pirani, J.R., Prado, J., Queiroz, L.P., Souza, S., Souza, V.C., Stehmann, J.R., Sylvestre, L.S., Walter, B.M.T., & Zappi, D.C. 2012. New Brazilian floristic list highlights conservation challenges. *BioScience* 62: 39–45.
- Freschet, G.T., Dias, A.T.C., Ackerly, D.D., Aerts, R., van Bodegom, P.M., Cornwell, W.K., Dong, M., Kurokawa, H., Liu, G., Onipchenko, V.G., Ordoñez, J.C., Peltzer, D.A., Richardson, S.J., Shidakov, I.I., Soudzilovskaia, N.A., Tao, J., & Cornelissen, J.H.C. 2011. Global to community scale differences in the prevalence of convergent over divergent leaf trait distributions in plant assemblages. *Global Ecology and Biogeography* 20: 755–765.
- Frontier, S. 1976. Étude de la décroissance des valeurs propres dans une analyse en composantes principales: comparaison avec le modèle de baton brisé. *Journal of Experimental Marine Biology and Ecology* 25: 1–26.
- Furley, P.A., & Ratter, J.A. 1988. Soil resources and plant communities of the central Brazilian Cerrado and their development. *Journal of Biogeography* 15: 97–108.
- Garnier, E., Cortez, J., Billès, G., Navas, M.-L., Roumet, C., Debussche, M., Laurent, G., Blanchard, A., Aubry, D., Bellmann, A., Neill, C., & Toussaint, J.-P. 2004. Plant functional markers capture ecosystem properties during secondary succession. *Ecology* 85: 2630–2637.
- Garnier, E., Lavorel, S., Ansquer, P., Castro, H., Cruz, P., Dolezal, J., Eriksson, O., Fortunel, C., Freitas, H., Golodets, C., Grigulis, K., Jouany, C., Kazakou, E., Kigel, J., Kleyer, M., Lehsten, V., Leps, J., Meier, T., Pakeman, R., Papadimitriou, M., Papanastasis, V.P., Quested, H., Quetier, F., Robson, M., Roumet, C., Rusch, G., Skarpe, C., Sternberg, M., Theau, J.-P., Thebault, A., Vile, D., & Zarovali, M.P. 2007. Assessing the effects of land-use change on plant traits, communities and ecosystem functioning in grasslands: a standardized methodology and lessons from an application to 11 European sites. *Annals of Botany* 99: 967–985.
- Garnier, E., & Navas, M.L. 2012. A trait-based approach to comparative functional plant ecology: Concepts, methods and applications for agroecology. A review. *Agronomy for Sustainable Development* 32: 365–399.
- Goedert, W.J. 1983. Management of the Cerrado soils of Brazil: a review. *Journal of Soil Science* 34: 405–428.
- Goodland, R., & Pollard, R. 1973. The Brazilian cerrado vegetation: a fertility gradient. *Journal of Ecology* 61: 219–224.
- Götzenberger, L., de Bello, F., Bråthen, K.A., Davison, J., Dubuis, A., Guisan, A., Lepš, J., Lindborg, R., Moora, M., Pärtel, M., Pellissier, L., Pottier, J., Vittoz, P., Zobel, K., & Zobel, M. 2012. Ecological assembly rules in plant communities-approaches, patterns and prospects. *Biological Reviews* 87: 111–127.
- Grime, J.P. 1998. Benefits of plant diversity to ecosystems: immediate, filter and founder effects. *Journal of Ecology* 86: 902–910.
- Hammer, O., Harper, D.A.T., & Ryan, P.D. 2001. PAST: paleontological statistics software package for education and data analysis. 9.
- Hawkins, B.A. 2012. Eight (and a half) deadly sins of spatial analysis. *Journal of Biogeography* 39: 1–9.

- Hoffmann, W.A., Franco, A.C., Moreira, M.Z., & Haridasan, M. 2005. Specific leaf area explains differences in leaf traits between congeneric savanna and forest trees. *Functional Ecology* 19: 932–940.
- Hoffmann, W. a, Geiger, E.L., Gotsch, S.G., Rossatto, D.R., Silva, L.C.R., Lau, O.L., Haridasan, M., & Franco, A.C. 2012. Ecological thresholds at the savanna-forest boundary: how plant traits, resources and fire govern the distribution of tropical biomes (F. Lloret, Ed.). *Ecology Letters* 15: 759–768.
- Jager, M.M., Richardson, S.J., Bellingham, P.J., Clearwater, M.J., & Laughlin, D.C. 2015. Soil fertility induces coordinated responses of multiple independent functional traits (G. De Deyn, Ed.). *Journal of Ecology* 103: 374–385.
- Klooster, S.H.J.-T., Thomas, E.J.P., & Sterck, F.J. 2007. Explaining interspecific differences in sapling growth and shade tolerance in temperate forests. *Journal of Ecology* 95: 1250–1260.
- Knops, J.M.H., & Reinhart, K. 2000. Specific leaf area along a nitrogen fertilization gradient. *The American Midland Naturalist* 144: 265–272.
- Koch, G.W., Sillett, S.C., Jennings, G.M., & Davis, S.D. 2004. The limits to tree height. *Nature* 428: 851–854.
- Laureto, L.M.O., & Cianciaruso, M. V. 2015. Trait distribution patterns in savanna and forest plant assemblages and their relationship with soil features. *Plant Ecology* 216: 629–639.
- Lavorel, S., & Garnier, E. 2002. Predicting changes in community composition and ecosystem functioning from plant traits: revisiting the Holy Grail. *Functional Ecology* 16: 545–556.
- Legendre, P. 1993. Spatial autocorrelation: trouble or new paradigm? *Ecology* 74: 1659–1673.
- Lenza, E., & Klink, C.A. 2006. Comportamento fenológico de espécies lenhosas em um cerrado sentido restrito de Brasília, DF. *Revista Brasileira de Botânica* 29: 627–638.
- Li, W., Cheng, J.-M., Yu, K.-L., Epstein, H.E., & Du, G.-Z. 2015. Niche and neutral processes together determine diversity loss in response to fertilization in an Alpine Meadow community (S. Hu, Ed.). *PLOS ONE* 10: e0134560.
- Liu, X., Swenson, N.G., Wright, S.J., Zhang, L., Song, K., Du, Y., Zhang, J., Mi, X., Ren, H., & Ma, K. 2012. Covariation in plant functional traits and soil fertility within two species-rich forests (K. O. Reinhart, Ed.). *PLoS ONE* 7: e34767.
- Lloyd, J., Goulden, M.L., Ometto, J.P., Patiño, S., Fyllas, N.M., & Quesada, C.A. 2009. Ecophysiology of forest and savanna vegetation. In Keller, M., Bustamante, M., Gash, J., & Dias, P.S. (eds.), *Amazonia and global change*, pp. 463–484. American Geophysical Union, Washington, DC.
- Marimon Junior, B.H., & Haridasan, M. 2005. Comparação da vegetação arbórea e características edáficas de um cerrado e um cerrado sensu stricto em áreas adjacentes sobre solo distrófico no leste de Mato Grosso , Brasil. *Acta Botanica Brasilica* 19: 913–926.
- Martorell, C., Almanza-Celis, C.A.I., Pérez-García, E.A., & Sánchez-Ken, J.G. 2015. Co-existence in a species-rich grassland: competition, facilitation and niche

- structure over a soil depth gradient (M. De Cáceres, Ed.). *Journal of Vegetation Science* 26: 674–685.
- Mayfield, M.M., & Levine, J.M. 2010. Opposing effects of competitive exclusion on the phylogenetic structure of communities. *Ecology Letters* 13: 1085–1093.
- McDonald, P.G., Fonseca, C.R., Overton, J.M., & Westoby, M. 2003. Leaf-size divergence along rainfall and soil-nutrient gradients: is the method of size reduction common among clades? *Functional Ecology* 17: 50–57.
- McGill, B.J., Enquist, B.J., Weiher, E., & Westoby, M. 2006. Rebuilding community ecology from functional traits. *Trends in ecology & evolution* 21: 178–85.
- Méio, B.B., Freitas, C. V, Jatobá, L., Silva, M.E.F., & Ribeiro, J.F. 2003. Influência da flora das florestas Amazônica e Atlântica na vegetação do cerrado sensu stricto. *Atlantic*
- Miranda, A.C., Miranda, H.S., Lloyd, J., Grace, J., Francey, R.J., McIntyre, J.A., Meir, P., Riggan, P., Lockwood, R., & Brass, J. 1997. Fluxes of carbon, water and energy over Brazilian cerrado: an analysis using eddy covariance and stable isotopes. *Plant, Cell and Environment* 20: 315–328.
- Moles, A.T., Perkins, S.E., Laffan, S.W., Flores-Moreno, H., Awasthy, M., Tindall, M.L., Sack, L., Pitman, A., Kattge, J., Aarssen, L.W., Anand, M., Bahn, M., Blonder, B., Cavender-Bares, J., Cornelissen, J.H.C., Cornwell, W.K., Díaz, S., Dickie, J.B., Freschet, G.T., Griffiths, J.G., Gutierrez, A.G., Hemmings, F.A., Hickler, T., Hitchcock, T.D., Keighery, M., Kleyer, M., Kurokawa, H., Leishman, M.R., Liu, K., Niinemets, Ü., Onipchenko, V., Onoda, Y., Penuelas, J., Pillar, V.D., Reich, P.B., Shiodera, S., Siefert, A., Sosinski, E.E., Soudzilovskaia, N.A., Swaine, E.K., Swenson, N.G., van Bodegom, P.M., Warman, L., Weiher, E., Wright, I.J., Zhang, H., Zobel, M., & Bonser, S.P. 2014. Which is a better predictor of plant traits: temperature or precipitation? (A. Helm, Ed.). *Journal of Vegetation Science* 25: 1167–1180.
- Moraes, D.A., Cavalin, P.O., Moro, R.S., Oliveira, R.A.C., Carmo, M.R.B., & Marques, M.C.M. 2015. Edaphic filters and the functional structure of plant assemblages in grasslands in southern Brazil (L. Duarte, Ed.). *Journal of Vegetation Science*. doi: 10.1111/jvs.12331
- Neri, A. V., Schaefer, C.E.G.R., Silva, A.F., Souza, A.L., Ferreira-Junior, W.G., & Meira-Neto, J.A.A. 2012. The influence of soils on the floristic composition and community structure of an area of Brazilian Cerrado vegetation. *Edinburgh Journal of Botany* 69: 1–27.
- Niklas, K.J. 2007. Maximum plant height and the biophysical factors that limit it. *Tree Physiology* 27: 433–440.
- Oliveira-Filho, A.T., & Ratter, J.A. 2002. Vegetation physiognomies and woody flora of the Cerrado biome. In Oliveira, P.S. & Marquis, R.J. (eds.), *The Cerrados of Brazil: ecology and natural history of a Neotropical Savanna*, pp. 91–120. Columbia University Press, New York.
- Oliveira, R.S., Bezerra, L., Davidson, E.A., Pinto, F., Klink, C.A., Nepstad, D.C., & Moreira, A. 2005. Deep root function in soil water dynamics in cerrado savannas of central Brazil. *Functional Ecology* 19: 574–581.
- Oliveras, I., & Malhi, Y. 2016. Many shades of green: the dynamic tropical forest–

- savannah transition zones. *Philosophical Transactions of the Royal Society B: Biological Sciences* 371: 20150308.
- Ordoñez, J.C., van Bodegom, P.M., Witte, J.-P.M., Wright, I.J., Reich, P.B., & Aerts, R. 2009. A global study of relationships between leaf traits, climate and soil measures of nutrient fertility. *Global Ecology and Biogeography* 18: 137–149.
- Pakeman, R.J., & Quested, H.M. 2007. Sampling plant functional traits: What proportion of the species need to be measured? *Applied Vegetation Science* 10: 91–96.
- Pellegrini, A.F.A. 2016. Nutrient limitation in tropical savannas across multiple scales and mechanisms. *Ecology* 97: 313–324.
- Peres-Neto, P.R., & Legendre, P. 2010. Estimating and controlling for spatial structure in the study of ecological communities. *Global Ecology and Biogeography* 19: 174–184.
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M.S., Cornwell, W.K., Craine, J.M., Gurvich, D.E., Urcelay, C., Veneklaas, E.J., Reich, P.B., Poorter, L., Wright, I.J., Ray, P., Enrico, L., Pausas, J.G., de Vos, A.C., Buchmann, N., Funes, G., Quétier, F., Hodgson, J.G., Thompson, K., Morgan, H.D., ter Steege, H., Sack, L., Blonder, B., Poschlod, P., Vaieretti, M. V., Conti, G., Staver, A.C., Aquino, S., & Cornelissen, J.H.C. 2013. New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany* 61: 167–234.
- Poorter, L. 2009. Leaf traits show different relationships with shade tolerance in moist versus dry tropical forests. *New Phytologist* 181: 890–900.
- Rangel, T.F., Diniz-filho, J.A.F., & Bini, L.M. 2010. SAM : a comprehensive application for Spatial Analysis in Macroecology. *Ecography*.
- Rangel, T.F.L.V.B., Diniz-Filho, J.A.F., & Bini, L.M. 2006. Towards an integrated computational tool for spatial analysis in macroecology and biogeography. *Global Ecology and Biogeography* 15: 321–327.
- Reich, P.B., Ellsworth, D.S., Walters, M.B., Vose, J.M., Gresham, C., Volin, J.C., & Bowman, W.D. 1999. Generality of leaf trait relationships: a test across six biomes. *Ecology* 80: 1955–1969.
- Ribeiro, J.F., & Walter, B.M.T. 2008. As principais fitofisionomias do bioma Cerrado. In Sano, S.M., Almeida, S.P., & Ribeiro, J.F. (eds.), *Cerrado: ecologia e flora*, pp. 151–212. Embrapa Cerrados, Brasília.
- Ruggiero, P.G.C., Batalha, M.A., Pivello, V.R., & Meirelles, S.T. 2002. Soil-vegetation relationships in cerrado (Brazilian savanna) and semideciduous forest, Southeastern Brazil. *Plant Ecology* 160: 1–16.
- Ryan, M.G., & Yoder, B.J. 1997. Hydraulic Limits to Tree Height and Tree Growth. *BioScience* 47: 235–242.
- Schachtman, D., Reid, R.J., & Ayling, S.M. 1998. Phosphorus uptake by plants: from soil to cell. *Plant Physiology* 116: 447–453.
- Schamp, B.S., & Aarssen, L.W. 2009. The assembly of forest communities according to maximum species height along resource and disturbance gradients. *Oikos* 118: 564–572.

- Sfair, J.C., Rosado, B.H.P., & Tabarelli, M. 2016. The effects of environmental constraints on plant community organization depend on which traits are measured (N. Mason, Ed.). *Journal of Vegetation Science*. doi: 10.1111/jvs.12453
- Silva, L.C.R., Haridasan, M., Sternberg, L.S.L., Franco, A.C., & Hoffmann, W.A. 2010. Not all forests are expanding over central Brazilian savannas. *Plant and Soil* 333: 431–442.
- Souza-Neto, A.C., Cianciaruso, M. V., & Collevatti, R.G. 2016. Habitat shifts shaping the diversity of a biodiversity hotspot through time: insights from the phylogenetic structure of Caesalpinioideae in the Brazilian Cerrado. *Journal of Biogeography* 43: 340–350.
- Svenning, J.-C., Kinner, D. a., Stallard, R.F., Engelbrecht, B.M.J., & Wright, S.J. 2004. Ecological Determinism in Plant Community Structure Across a Tropical Forest Landscape. *Ecology* 85: 2526–2538.
- Swenson, N.G., & Enquist, B.J. 2007. Ecological and evolutionary determinants of a key plant functional trait: wood density and its community-wide variation across latitude and elevation. *American Journal of Botany* 94: 451–459.
- Vellend, M. 2010. Conceptual synthesis in community ecology. *The Quarterly review of biology* 85: 183–206.
- Vile, D. 2005. Specific leaf area and dry matter content estimate thickness in laminar leaves. *Annals of Botany* 96: 1129–1136.
- Wagatsuma, T. 1983. Characterization of absorption sites for aluminum in the roots. *Soil Science and Plant Nutrition* 29: 499–515.
- Wang, D., Maughan, M.W., Sun, J., Feng, X., Miguez, F., Lee, D., & Dietze, M.C. 2012. Impact of nitrogen allocation on growth and photosynthesis of *Miscanthus* (*Miscanthus* × *giganteus*). *GCB Bioenergy* 4: 688–697.
- Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil* 199: 213–227.
- Wigley, B.J., Slingsby, J.A., Díaz, S., Bond, W.J., Fritz, H., & Coetsee, C. 2016. Leaf traits of African woody savanna species across climate and soil fertility gradients: evidence for conservative vs . acquisitive resource use strategies. *Journal of Ecology*. doi: 10.1111/1365-2745.12598
- Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-bares, J., Chapin, T., Cornelissen, J.H.C., Diemer, M., Flexas, J., Garnier, E., Groom, P.K., & Gulias, J. 2004. The worldwide leaf economics spectrum. *Nature* 428: 821–827.
- Zava, P.C., & Cianciaruso, M. V. 2014. Can we use plant traits and soil characteristics to predict leaf damage in savanna woody species? *Plant Ecology* 215: 625–637.

Supplementary Material

Supplementary table 1. Climate characteristics of the six sites sampled in Brazilian Savanna. The values represented a range, when available, of 20 years of data collected in the nearest weather station of each site. The fire frequency is related only to savannas sites. A low fire frequency indicates an interval about of five years among fire events, and high fire frequency of 10 years.


Environmental characteristics	Assis Ecological State - SP	Vassununga State Park - SP	Emas National Park - GO	Jataí - GO	Bacaba Park - MT	Ribeirão Cascalheira - MT
Coordinates	50W 22', 22S 35'	47W 37', 21S 36'	52W 59', 17S 54'	51W 32', 17S 45'	52W 21', 14S 42'	51W 46', 12S 49'
Fire frequency	low	low	high	high	high	low
Climate Type	Cfa	Cwa	Aw	Aw	Aw	Aw
Altitude (m)	560	740	840	860	325	374
Mean Annual Temp. (°C)	23.6	22.3	24.6	22.6	24.8	25.4
Mean Minimum Temp. (°C)	19.0	16.6	18.9	17.0	18.9	20.1
Precipitation (mm yr ⁻¹)	1352	1437	1745	1605	1433	1822
Evapotranspiration (mm)	129	92	82	121	134	136
Mean Monthly Relative Humidity (%)	64	70	69	69	77	71

Supplementary table 2. Mean values and coefficient of variation (in parentheses) of the soil variables to sites of seasonal forests and savannas environments in the Brazilian Cerrado. These values were calculated based in values of parameters in ten plots each site.


Soil characteristics	Assis Ecological State - SP		Vassununga State Park - SP		Emas National Park - GO		Jataí - GO		Bacaba Park - MT		Ribeirão Cascalheira - MT	
	Forest	Savanna	Forest	Savanna	Forest	Savanna	Forest	Savanna	Forest	Savanna	Forest	Savanna
N	0.32 (0.46)	0.25 (0.40)	0.22 (0.29)	0.12 (0.36)	0.26 (0.36)	0.09 (0.45)	0.18 (0.15)	0.17 (0.25)	0.11 (0.23)	0.08 (0.18)	0.10 (0.14)	0.05 (0.31)
P	8.39 (0.54)	3.91 (0.83)	4.51 (1.07)	3.89 (0.85)	1.79 (0.38)	0.56 (0.37)	1.57 (0.44)	1.03 (0.78)	1.15 (0.53)	2.03 (0.07)	1.75 (0.15)	2.04 (0.33)
K	46.3 (0.38)	11.8 (0.34)	11.2 (0.37)	13.1 (0.27)	42.4 (0.18)	31.4 (0.27)	28.4 (0.44)	22.3 (0.26)	46.7 (0.23)	46.2 (0.15)	17.7 (0.19)	18.3 (0.43)
SMO	3.10 (0.38)	1.70 (0.37)	5.13 (0.20)	3.10 (0.17)	5.87 (0.29)	3.08 (0.44)	6.22 (0.19)	4.79 (0.18)	1.07 (0.55)	1.73 (0.19)	2.49 (0.12)	1.83 (0.33)
pH	7.09 (0.10)	4.85 (0.03)	6.16 (0.11)	4.51 (0.02)	4.20 (0.11)	4.30 (0.06)	5.26 (0.03)	5.26 (0.02)	4.65 (0.02)	5.04 (0.03)	4.10 (0.02)	4.95 (0.05)
Al	0	0.88 (0.18)	0.12 (2.87)	1.70 (0.12)	1.65 (0.24)	1.19 (0.24)	1.03 (0.38)	0.65 (0.42)	0.85 (0.22)	1.16 (0.12)	1.12 (0.10)	0.77 (0.15)
Clay	11.2 (0.25)	8.0 (0.24)	44.7 (0.25)	10.6 (0.08)	26.1 (0.29)	29.4 (0.67)	54.6 (0.14)	51.0 (0.08)	20.1 (0.18)	12.6 (0.20)	16.7 (0.13)	2.9 (0.41)

Supplementary table 3. Mean traits and number of species sampled to each sites of the soil variables to sites of seasonal forests and savannas environments in the Brazilian Cerrado. Density ≤ 3 represents the number and percentage of species with individual density equal or less that three individuals in each site, and density ≥ 4 denotes the number and percentage (in parentheses) of species with individual density equal or less that four individuals in each site.


Site	Number of species	density ≤ 3	density ≥ 4	Height	LA	SLA	LPC	SSD	LT	C/N
<i>Forest</i>										
Assis Ecological State - SP	39	31 (79)	8 (21)	13.0	132.2	133.8	1.48	0.51	0.16	2.13
Vassununga State Park - SP	31	18 (58)	13 (42)	9.5	276.1	159.3	1.99	0.42	0.15	2.07
Emas National Park - GO	21	15 (71)	6 (29)	15.7	128.9	105.8	0.84	0.57	0.20	3.36
Jataí - GO	29	12 (41)	17 (59)	13.1	200.6	113.6	1.28	0.59	0.19	3.07
Bacaba Park - MT	28	16 (57)	12 (43)	11.6	78.1	110.8	1.06	0.62	0.15	2.78
Ribeirão Cascalheira - MT	52	20 (38)	32 (62)	14.6	138.7	95.6	0.86	0.59	0.18	2.73
<i>Savanna</i>										
Assis Ecological State - SP	44	16 (36)	28 (64)	4.5	76.8	85.3	0.98	0.51	0.21	2.83
Vassununga State Park - SP	47	16 (34)	31 (66)	4.3	71.2	80.9	1.22	0.48	0.25	3.30
Emas National Park - GO	51	34 (67)	17 (33)	2.7	70.0	68.4	0.83	0.48	0.34	3.92
Jataí - GO	43	7 (16)	36 (84)	3.1	125.2	71.0	0.99	0.46	0.32	3.73
Bacaba Park - MT	90	15 (17)	74 (83)	4.3	139.3	84.7	1.10	0.53	0.23	3.55
Ribeirão Cascalheira - MT	66	24 (36)	42 (64)	5.0	81.8	78.0	0.91	0.56	0.24	3.32




Supplementary figure 1. Contribution of edaphic parameters for the toxicity (PC1) and fertility (PC2) gradients in seasonal forest in the Brazilian Cerrado. Bootstrap confidence intervals when do not overlap zero indicates significant relationship (positive or negative) with the component principal. The numbers on the bars indicates the correlation values of each variable with the PCs.




Supplementary figure 2. Contribution of edaphic parameters for the fertility (PC1) and toxicity (PC2) gradients in savannas in the Brazilian Cerrado. Bootstrap confidence intervals when do not overlap zero indicates significant relationship (positive or negative) with the component principal. The numbers on the bars indicates the correlation values of each variable with the PCs.



Supplementary figure 3. Relationship between individual density and soil proprieties of seasonal and savanna in Brazilian Savanna. Positive values in the PCs indicates greater levels of toxicity or fertility on the gradient.



Supplementary figure 4. Shifts in community-weighted trait means along soil toxicity (PC1) and fertility (PC2) gradients in seasonal forests in the Brazilian Cerrado. Beta regression coefficient (β) indicates the intensity and direction of each regression with the gradients. The β coefficients were derived from linear regression analyses. Hmax = maximum plant height (m); LA = leaf area (cm^2); SLA = specific leaf area (cm^2); LPC = leaf phosphorous content ($\text{g/kg}(\%)$); SSD = Stem-specific density (mg mm^{-3}); LT = leaf thickness (mm); C/N = leaf carbon-nitrogen ratio. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, NS non-significant. Positive values in the PCs indicates greater levels of toxicity or fertility on the gradient.




Supplementary figure 5. Shifts in community-weighted trait means along soil fertility (PC1) and toxicity (PC2) gradients in savannas in the Brazilian Cerrado. Beta regression coefficient (β) indicates the intensity and direction of each regression with the gradients. The β coefficients were derived from linear regression analyses. Hmax = maximum plant height (m); LA = leaf area (cm^2); SLA = specific leaf area (cm^2); LPC = leaf phosphorous content ($\text{g/kg}(\%)$); SSD = Stem-specific density (mg mm^{-3}); LT = leaf thickness (mm); C/N = leaf carbon-nitrogen ratio. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, NS non-significant, † indicate low explanatory power of predictors. Positive values in the PCs indicates greater levels of toxicity or fertility on the gradient.

Appendix 1. Set of graphics representing the spatial patterns of all the predictor variables sampled in savanna and seasonal forests environments, displayed according to acquisitive or conservative strategy of resources. Spatial autocorrelogram of the patterns in the observed (red lines), expected (blue lines), and residuals (green lines).


Community-weighted means

Forest – acquisitive strategy


Hmax – maximum plant height




LA – leaf area



SLA – specific leaf area




LPC – leaf phosphorous content




Forest – conservative strategy


SSD – stem-specific density



LT – leaf thickness




C/N – leaf carbon-nitrogen ratio




Savanna – acquisitive strategy


Hmax – Maximum plant height




LA – leaf area



SLA – specific leaf area




LPC – leaf phosphorous content




Savanna – conservative strategy


SSD – stem-specific density




LT – leaf thickness




C/N – leaf carbon-nitrogen ratio



Density forest



Density savanna



CAPÍTULO III

Formatado nas normas da revista:

Journal of Ecology

Insect herbivore damage is not related with host plant ecological and evolutionary distances

**Leandro Maracahipes^{1*}, Fernando Landa Sobral¹, Leonardo Lima Bergamini¹,
Marcus Vinicius Cianciaruso¹, Mário Almeida-Neto¹, Walter Santos de Araújo^{1,2}**

¹Departamento de Ecologia, Instituto de Ciências Biológicas, Universidade Federal de Goiás, Goiânia, GO, 74001-970, Brazil; ²Departamento de Biologia Geral, Centro de Ciências Biológicas, Universidade Estadual de Montes Claros, Montes Claros, MG, 39401-089, Brazil

***Correspondence author (lmaracahipes@gmail.com)**

Summary

1. The spatial distribution and the identity of co-existing plants individuals should play an important role to herbivory levels. There is a growing interest in understand how the variation on functional traits and phylogenetic relationship in the neighboring affects the herbivory on focal plants.
2. To investigate whether ecological and evolutionary distances between individual plants and their neighbouring plants mediates leaf herbivory levels, we sampled 815 individuals belonging to 27 woody plant species in 49 savanna plots. For each individual, we quantified data on leaf damage (%), specific leaf area ($\text{cm}^2 \text{ g}^{-1}$), leaf toughness (mm), plant height (cm), and abundance. To evaluate the herbivory effects at the individual level, we quantified relative trait values for each individual and the phylogenetic isolation of each individual in relation to the neighboring plants.
3. Contrary to our predictions, we found that ecological and evolutionary distances between focal plants and their neighbouring plants did not mediate leaf herbivory levels in the studied plants.
4. We showed that that the phylogenetic isolation, specific leaf area, leaf toughness, plant height and plant abundance did not influence the herbivore damage on the woody plants of Neotropical savannas, both in the individual context as at neighboring plants. Our results suggest that dominance of herbivore generalists, coevolution between plant and specialist herbivores and preferential consume of young leaves can be more important for determine the leaf damage level of the focal plant in savanna woody species, than the neighboring context.
5. *Synthesis.* We demonstrated that ecological and evolutionary distances between individual plants and their neighbouring plants did not mediates leaf herbivory levels in savanna woody species. These findings suggest that dominance of herbivore generalists,

coevolution between plant and specialist herbivores and preferential consume of young leaves can be more important than the neighboring context for determine the leaf damage by herbivores in savanna woody species. Furthermore, we suggest that this approach be replicated to other plant groups and environments.

Key-words: neighboring plants, herbivory damage, plant functional trait, phylogenetic and ecological isolation, Janzen-Connell hypothesis, Neotropical savanna, cerrado

Introduction

Variation in leaf herbivory levels is a common phenomenon among co-occurring plant species (Cornelissen, Wilson Fernandes & Vasconcellos-Neto 2008; Carmona, Lajeunesse & Johnson 2011; Schuldt *et al.* 2012; Cárdenas *et al.* 2014). Potential explanations to such variation include differences in plant defensive traits, such as type and amount of secondary metabolites and leaf toughness, as well as differences in leaf nutritional content (Agrawal 2007; Carmona *et al.* 2011; Ali & Agrawal 2012; Loranger *et al.* 2012; Cárdenas *et al.* 2014). For example, leaves with high nitrogen concentration and high specific leaf area are more palatable to herbivores (Hanley *et al.* 2007) than leaves with high carbon-nitrogen ratio (Coley & Barone 1996). At the individual scale height influences the accessibility and localization of the plant by herbivores (Loranger *et al.* 2012) whereas at the assemblage level plant density is a signal of resource concentration (Hambäck & Englund 2005). Yet several studies investigated the direct effect of these factors on herbivory (Janzen 1970; Twigg & Socha 1996; Coley & Barone 1996; Agrawal 2007; Hanley *et al.* 2007) there is a growing interest in

understand how the variation of these factors in the neighboring affects the herbivory on focal plants (Barbosa *et al.* 2009; Underwood *et al.* 2014; Kim & Underwood 2015).

Because plants are sessile organisms their spatial distribution and the identity of co-existing individuals should play an important role to herbivory levels. Neighboring plants can attract, retain or repulse herbivores (Root 1973; Hambäck & Englund 2005; Hakes & Cronin 2011) affecting the damage that a focal plant may experience. Damage to neighboring plants can also promote volatile emissions that elicit the induction of defenses in a given focal plant, leading to a reduction in herbivory (Barbosa *et al.* 2009). On the other hand, neighboring properties, such as plant abundance and identity may facilitate the survival of a focal plant by altering its conspicuity (Barbosa *et al.* 2009; Kim & Underwood 2015; Champagne, Tremblay & Côté 2016; Murphy, Xu & Comita 2016). The Janzen-Connell hypothesis predict that herbivore pressure should be positively correlated with conspecific neighbor abundance because specialist insect herbivores can be attract to high-density patches with preferred resource (Janzen 1970; Comita *et al.* 2014; Murphy *et al.* 2016). Palatable plants in an assemblage of unpalatable plants can remain undetected by herbivores and thereby escape from consumption (Callaway, Kikvidze & Kikodze 2000; Baraza, Zamora & Hódar 2006). Thus, studies of this nature are important because the neighboring effects on focal plant may influences their growth and survival, and determine species coexistence and diversity patterns in the local communities (Queenborough *et al.* 2007; Kim & Underwood 2015; Champagne *et al.* 2016).

The phylogenetic relationship between a given plant individual and its neighbors can act as an important mediator of plant–herbivore interactions (Gilbert *et al.* 2012; Lebrija-Trejos *et al.* 2014). This is because key plant functional traits can be phylogenetically conserved (Lebrija-Trejos *et al.* 2014) and closely related plant species

often share more coevolutionary relationships with herbivore consumers and other natural enemies (Cavender-Bares *et al.* 2009; Futuyma & Agrawal 2009; Pearse & Hipp 2009; Gilbert *et al.* 2012). Thus, herbivores may limit the co-existence of closely related plants (Becerra 2007; Futuyma & Agrawal 2009). In this context, focal plants co-occurring with phylogenetically distant neighbors are expected to show lower herbivory levels than those co-occurring with closely related plants (Gilbert & Webb 2007; Chen *et al.* 2016).

In this study, we investigated whether ecological and evolutionary distances between individual plants and their neighbouring plants mediates leaf herbivory levels. We hypothesized that functionally and phylogenetically more similar neighbouring plants will show higher herbivore damage than functionally and phylogenetically less similar neighbouring focal plants. Specifically, we answered the following questions: (1) Does herbivory damage level in a focal plant is mediate by specific neighbour plant functional traits? (2) Do focal plants with leaf traits similar to neighbors plants (specific leaf area and leaf toughness) have higher herbivory damage levels than focal plants with distinct leaf traits to neighbouring? (3) Do focal plants larger than their neighbor plants have higher herbivory damage levels than focal plants smaller than neighborhood plants? (4) Do focal plants closely related to neighbouring plants have higher herbivory damage levels than focal plant distantly related to neighbouring plants?

Material and methods

STUDY SITE

The Cerrado is the largest savanna region of South America (Ratter, Ribeiro & Bridgewater 1997), and one of the richest savanna in the world, with a high habitat diversity and a large number of endemic plants and animals species (Klink & Machado

2005). The predominant vegetation of the Cerrado is a Neotropical savanna (cerrado *stricto sensu*) that is composed by shrubs, tree and grasses coexisting with the herbaceous layer (Ratter *et al.* 1997). The soils of the savannas of the Cerrado are nutrient-poor and well-drained and act as an environmental filter on species establishment (Furley & Ratter 1988). Another important filter is fire, that through time has selected species with traits of resistance or resilience to stresses caused by burning (Simon *et al.* 2009). The low soil fertility and high fire frequency filters species with adaptive traits to environmental stress and with anti-herbivore characteristics (Silva & Batalha 2011). Environments with high environmental stress, as the Cerrado, offer an under explored opportunity to understand how ecological and evolutionary relationships between neighboring species may determine the damage level caused by herbivores.

We conducted this study in the Emas National Park (ENP; 17°54'S, 52°59'W) in central Brazilian plateau, a region included predominantly in the Cerrado domain. The ENP occupies an area of 133,000 ha of mixed formations of savannas, and with elevations ranging between 720 to 890 m. The climate of the region is defined as Aw of Köppen (Peel *et al.* 2007), characterized by a mean annual temperature of 24.6 °C and a mean annual precipitation ranging from 1,200 to 2,000 mm (Ramos-Neto & Pivello 2000). The soils are predominantly characterized as oxisols and dystrophic (Silva & Batalha 2008). The vegetation in the park comprises forest formations (e.g., *semidecidual and riparian forest*) and savannic formations (e.g., dense and open savannas) (Ribeiro & Walter 2008), wherein the last occupy 78.5% of the area of the park (Ramos-Neto & Pivello 2000).

ECOLOGICAL DISTANCES AND PHYLOGENETIC ISOLATION

We used data from 49 plots of 10 x 10 m where all woody plant individuals with a stem circumference larger than 10 cm at soil level were sampled (Zava & Cianciaruso 2014). These plots were randomly distributed in a gradient of open to dense savannas (39 plots in open savannas and 10 plots in dense savannas). We used only those species with at least ten individuals and that occurred in at least five plots. Overall, we analyzed 815 individuals of the 27 most abundant plant species (Table S1). *Anadenanthera peregrina* (Fabaceae) was excluded from the analysis due to its tiny folioles, which makes virtually impossible to measure leaf damage for this species. For each individual we sampled data on leaf damage (%), specific leaf area ($\text{cm}^2 \text{g}^{-1}$), leaf toughness (mm), and plant height (cm). Additionally, we calculated plant abundance per plot. For each individual we collected 10 mature leaves, of which five leaves were used to calculate leaf toughness, and the other five to calculate leaf damage, leaf area and specific leaf area. All trait measurements followed the protocols presented in Pérez-Harguindeguy *et al.* (2013).

To evaluate the herbivory effects at the level of individual, we quantified relative trait values for each individual (Table S2). Thus, for example, positive values for height indicate a higher-than-average individual and probably should be more attacked by herbivores, whereas negative values indicate lower-than-average ones and less attacked. To estimate phylogenetic isolation that is, the phylogenetic distance between each focal individual and the other co-occurring individuals, we used a dated phylogeny recently constructed by Zanne *et al.* (2014). From this phylogeny, we extracted the phylogenetic relationships of all 27 plants species analyzed (Fig. 1). However, 20 species were absent of the original phylogeny, we included these species in the phylogeny as polytomies at the genus (18 species) and family (2 species) level.

We quantified the phylogenetic isolation of each individual in relation to the neighboring plants as the sum of the branch lengths in the phylogeny connecting the focal individual and all the others plant individuals co-occurring in the same plot. Conspecific individuals had their phylogenetic distance set to zero. Therefore, this measure included the abundance of the neighboring plants at each plot, given that every single individual contributed to the sum. Individuals inserted in a local community comprising distantly-related plants will have high values of phylogenetic isolation.

STATISTICAL ANALYSIS

To investigate the factors that influence the leaf herbivore damage in each individual we built a model including SLA, relative SLA, LT, relative LT, conspecific abundance, height, relative height and phylogenetic isolation (Table 1). We build a random intercept linear mixed model relating the percentual herbivory of each individual and its trait values, as well as various measures of neighborhood context (Supporting Information Table S2), with plant species nested by plot identity as a random effect. Random effects structure was tested by likelihood ratio tests. We then computed the standardized coefficients with $\pm 1.96 * SE$ confidence intervals to assess the relative contribution of each variable to the differences in herbivory. Degrees of freedom for the *t-values* p computations were computed using the Satterthwaite's approximation (Satterthwaite 1946). The models were fit with the “lmer” function in the “lme4 package in R (R Core Team 2016).

Results

Leaf damage *per* plant individual ranged from 0.005% to 29.6%, with 95% of plant individuals having up to 10% of their leaf surface consumed by herbivores (Fig. 1). We found that the leaf toughness and relative leaf toughness were related to differences in leaf herbivory (Table 1, Fig. S1). However, the explanatory power of the global mixed effects model was very low. The marginal R^2 that are associated with fixed effects was of 0.07, while the conditional R^2 that considers both fixed and random effects was 0.42. Due to the low explanatory power of the fixed variables, we consider that all the predictor variables were not related to leaf damage (Table 1, Fig. 2).




Fig. 1. The phylogenetic tree assembled for 27 cerrado species, with respective abundance and mean leaf damage for each species, utilized in the analyzes in Emas National Park, Brazil. Abundance is presented in log (for original values see Table 2S). The relationship among species was based on dated phylogeny constructed by Zanne *et al.* (2014). The scale is in million years and the values of abundance are logarithmic.

Table 1. Results of the global mixed effects model relating the predictors and relative herbivory. Standardized coefficients and standard errors are shown for the fixed effects conspecific abundance, phylogenetic isolation, relative specific leaf area, relative leaf toughness, and relative height. The conditional R^2 was of 0.42 and marginal R^2 was of 0.07. P values computed using the Satterthwaite's approximation (815 observations; 27 species). Ma: millions of years before of present; n: number of individuals.

	<i>Groups</i>	Variance	Std.Dev.		
Random effects	Plot	0.74	0.86		
	Species Identity	5.10	2.26		
	Residual	9.78	3.13		
	<i>Variables</i>	<i>Std. Coeff.</i>	<i>Std. Error</i>	<i>t-value</i>	<i>p</i>
Fixed effects	SLA (cm ² g ⁻¹)	0.040	0.039	1.023	0.313
	Relative SLA (cm ²)	-0.066	0.039	-1.715	0.094
	Leaf toughness (mm)	1.321	0.385	3.432	0.002
	Relative leaf toughness (mm)	-1.070	0.441	-2.425	0.018
	Phylogenetic isolation (Ma)	-2.57x10 ⁻⁴	2.53x10 ⁻⁴	-1.018	0.315
	Conspecific abundance (n)	-0.072	0.065	-1.105	0.272
	Height (cm)	-1.44x10 ⁻³	5.11x10 ⁻³	-0.282	0.780
	Relative height (cm)	-1.67x10 ⁻³	5.28x10 ⁻³	0.317	0.753




Fig. 2. Standardized effect size (mean $\pm$ 95% confidence intervals) computed under a global mixed effects model relating the predictors and relative herbivory for 27 cerrado species in Emas National Park, Brazil. Black line represents predictors related to effects at the level of individual, while white line indicates effects at the level of plots. Confidence intervals when do not overlap zero indicates significant relationship (positive or negative), but the marginal R^2 of the model was very low (0.07).

Discussion

Neighboring plants can affect the herbivore damage levels, attracting, retaining or repulsing herbivores (Root 1973; Hambäck & Englund 2005; Hakes & Cronin 2011). We hypothesized that focal plants more similar to neighboring plants will show higher herbivory damage than focal plants less similar to neighbouring plants. However, our results did not corroborate this expectation, because we found that the herbivory level of focal plants were not affected by traits of neighboring plants. This result can be explained by three principal factors: (1) dominance of generalist herbivores in savannas, (2) coevolution of plants and specialist herbivores, and (3) preferential consume of immature leaves by herbivores.

The herbivorous fauna in Neotropical savannas is composed predominantly by generalist insects, such as leaf-cutter ants, grasshoppers and caterpillars (Diniz & Morais 1997; Ribeiro, Carneiro & Fernandes 1998; Mundim, Costa & Vasconcelos 2009). These generalist herbivores can consume indiscriminately different plant types (Meijden 1996; Diniz & Morais 1997; Ali & Agrawal 2012), and range seasonally in the their diet breadth (Scherrer *et al.* 2016). The indiscriminate consume of savanna plants by generalist herbivores can explain the lack of effect of neighbor plants on leaf damage of focal plants.

Several herbivores insects coevolved with their host plants (Coley & Barone 1996) and this represent evolutionary and ecological consequences for plant-herbivore interactions (Becerra 2003, 2007; Futuyma & Agrawal 2009). These species of herbivores usually are specialized in their host plants, consuming their leaves independently of the traits from neighbor individuals. Another potential explanation may be related to the preferential consume of young leaves by herbivore insects. When young, savanna species leaves have high nutritional quality and are vulnerable to

herbivores, contrary to mature leaves that have low nutrient concentrations and high protection against herbivores (Coley & Barone 1996; Awmack & Leather 2002; Lewinsohn, Freitas & Prado 2005; Mundim *et al.* 2009). Mature leaves of savanna plant species have strategy of protection against herbivores, such as thick leaves with trichomes and high leaf carbon-nitrogen ratio (Dantas & Batalha 2012; Dantas, Batalha & Pausas 2013). Moreover, savanna plants synchronize the leaf production at the end of dry season (Marquis, Diniz & Morais 2001; Mundim *et al.* 2009), which can affect the seasonal supply of resources for herbivores (Scherrer *et al.* 2016). Thus, insect herbivores in Neotropical savannas tend to consume indiscriminately immature leaves independent of neighbouring context, mainly in stage of high leaf production.

Our results also showed that phylogenetic isolation, leaf toughness, plant height and conspecific abundance do not influence the herbivore damage on the plants of this Neotropical savanna. The non-effect of phylogenetic isolation on the herbivory can be explained that, even for important traits in the determination of leaf damage by herbivore, the leaf damage level did not have relationship with the kinship between species (Agrawal 2007; Pearse & Hipp 2009; Uriarte *et al.* 2010; Gilbert *et al.* 2012), because same species phylogenetically more related or distant are consumed by herbivores. Pearse and Hipp (2009) argument that leaf volatile chemicals can be more phylogenetically informative than structural defence traits.

A negative effect of conspecific abundance via shared enemies is the main mechanism behind the well-tested Janzen-Connell hypothesis (Janzen 1970; Comita *et al.* 2014; Kim & Underwood 2015). However, we did not find effects of conspecific abundance at the plot on the levels of herbivory. The relationship between plant patch size and insect abundance is highly variable and dependent on the group specificities (Bukovinszky *et al.* 2005; Hambäck & Englund 2005). Therefore, this lack of effect

may be a reflection of the response pattern of visual and olfactory searchers of the most important chewing groups in our study system, that can be able to reduce differences between small and large patches, such as orthopteran and larval lepidopteran (Hambäck & Englund 2005).

This is the first study that investigate the leaf damage caused by insect herbivores in a neighboring context in Neotropical savannas. In conclusion, we highlight that the phylogenetic isolation, specific leaf area, leaf toughness, plant height and plant abundance not influenced the herbivore damage on the woody plants of Neotropical savannas, both in the individual context as at neighboring plants. Overall, our results suggest that dominance of herbivore generalists, coevolution between plant and specialist herbivores and preferential consume of young leaves can be more important than the neighboring context for determine the leaf damage by herbivores in savanna woody species studied. As savanna plant species have leaves with low quality nutritional, herbivores compensate increasing the consumption rates to obtain more nitrogen from leaves. Furthermore, generalist herbivores consume immature leaves because they are vulnerable and present high-quality nutrients (Marquis *et al.* 2001). Future studies may improve these investigations, adding more species and develop this approach with other plant groups, including herbaceous and grasses, and species from other Neotropical vegetation types adjacent to savannas, such as forests and grassland environments.

Acknowledgements

The authors are thankful to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the grant to LM, FLS, LLB and WSA and to the Brazilian Research Council (CNPq) for a productivity grant to MVC and MAN. We thank the

CNPq/Fapeg-Brasil by supported to search by PELD (Sites 13) and GENPAC program (306573/2009-1).

References

- Agrawal, A.A. (2007) Macroevolution of plant defense strategies. *Trends in Ecology & Evolution*, **22**, 103–109.
- Ali, J.G. & Agrawal, A.A. (2012) Specialist versus generalist insect herbivores and plant defense. *Trends in Plant Science*, **17**, 293–302.
- APG-IV. (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society*, **181**, 1–20.
- Awmack, C.S. & Leather, S.R. (2002) Host plant quality and fecundity in herbivorous insects. *Annual Review of Entomology*, **47**, 817–844.
- Baraza, E., Zamora, R. & Hódar, J.A. (2006) Conditional outcomes in plant-herbivore interactions: neighbours matter. *Oikos*, **113**, 148–156.
- Barbosa, P., Hines, J., Kaplan, I., Martinson, H., Szczepaniec, A. & Szendrei, Z. (2009) Associational resistance and associational susceptibility: having right or wrong neighbors. *Annual Review of Ecology, Evolution, and Systematics*, **40**, 1–20.
- Becerra, J.X. (2003) Synchronous coadaptation in an ancient case of herbivory. *Proceedings of the National Academy of Sciences*, **100**, 12804–12807.
- Becerra, J.X. (2007) The impact of herbivore-plant coevolution on plant community structure. *Proceedings of the National Academy of Sciences*, **104**, 7483–7488.
- Bukovinszky, T., Potting, R.P.J., Clough, Y., van Lenteren, J.C. & Vet, L.E.M. (2005) The role of pre- and post- alighting detection mechanisms in the responses to patch size by specialist herbivores. *Oikos*, **109**, 435–446.

- Callaway, R.M., Kikvidze, Z. & Kikodze, D. (2000) Facilitation by unpalatable weeds may conserve plant diversity in overgrazed meadows in the Caucasus Mountains. *Oikos*, **89**, 275–282.
- Cárdenas, R.E., Valencia, R., Kraft, N.J.B., Argoti, A. & Dangles, O. (2014) Plant traits predict inter- and intraspecific variation in susceptibility to herbivory in a hyperdiverse Neotropical rain forest tree community (ed N Swenson). *Journal of Ecology*, **102**, 939–952.
- Carmona, D., Lajeunesse, M.J. & Johnson, M.T.J. (2011) Plant traits that predict resistance to herbivores. *Functional Ecology*, **25**, 358–367.
- Cavender-Bares, J., Kozak, K.H., Fine, P.V.A. & Kembel, S.W. (2009) The merging of community ecology and phylogenetic biology. *Ecology Letters*, **12**, 693–715.
- Champagne, E., Tremblay, J.-P. & Côté, S.D. (2016) Spatial extent of neighboring plants influences the strength of associational effects on mammal herbivory. *Ecosphere*, **7**, e01371.
- Chen, Y., Wright, S.J., Muller-Landau, H.C., Hubbell, S.P., Wang, Y. & Yu, S. (2016) Positive effects of neighborhood complementarity on tree growth in a Neotropical forest. *Ecology*, **97**, 776–785.
- Coley, P.D. & Barone, J.A. (1996) Herbivory and plant defenses in tropical forests. *Annual Review of Ecology and Systematics*, **27**, 305–335.
- Comita, L.S., Queenborough, S.A., Murphy, S.J., Eck, J.L., Xu, K., Krishnadas, M., Beckman, N. & Zhu, Y. (2014) Testing predictions of the Janzen-Connell hypothesis: a meta-analysis of experimental evidence for distance- and density-dependent seed and seedling survival (ed L Gómez-Aparicio). *Journal of Ecology*, **102**, 845–856.
- Cornelissen, T., Wilson Fernandes, G. & Vasconcellos-Neto, J. (2008) Size does matter:

- variation in herbivory between and within plants and the plant vigor hypothesis. *Oikos*, **117**, 1121–1130.
- Dantas, V. de L. & Batalha, M.A. (2012) Can antiherbivory resistance explain the abundance of woody species in a Neotropical savanna? *Botany*, **90**, 93–99.
- Dantas, V.L., Batalha, M.A. & Pausas, J.G. (2013) Fire drives functional thresholds on the savanna–forest transition. *Ecology*, **94**, 2454–2463.
- Diniz, I.R. & Morais, H.C. (1997) Lepidopteran caterpillar fauna of cerrado host plants. *Biodiversity and Conservation*, **6**, 817–836.
- Furley, P.A. & Ratter, J.A. (1988) Soil resources and plant communities of the central Brazilian Cerrado and their development. *Journal of Biogeography*, **15**, 97–108.
- Futuyma, D.J. & Agrawal, A.A. (2009) Macroevolution and the biological diversity of plants and herbivores. *Proceedings of the National Academy of Sciences*, **106**, 18054–18061.
- Gilbert, G.S., Magarey, R., Suiter, K. & Webb, C.O. (2012) Evolutionary tools for phytosanitary risk analysis: phylogenetic signal as a predictor of host range of plant pests and pathogens. *Evolutionary Applications*, **5**, 869–878.
- Gilbert, G.S. & Webb, C.O. (2007) Phylogenetic signal in plant pathogen–host range. *Proceedings of the National Academy of Sciences*, **104**, 4979–4983.
- Hakes, A.S. & Cronin, J.T. (2011) Environmental heterogeneity and spatiotemporal variability in plant defense traits. *Oikos*, **120**, 452–462.
- Hambäck, P.A. & Englund, G. (2005) Patch area, population density and the scaling of migration rates: the resource concentration hypothesis revisited. *Ecology Letters*, **8**, 1057–1065.
- Hanley, M.E., Lamont, B.B., Fairbanks, M.M. & Rafferty, C.M. (2007) Plant structural traits and their role in anti-herbivore defence. *Perspectives in Plant Ecology*,

- Evolution and Systematics*, **8**, 157–178.
- Janzen, D.H. (1970) Herbivores and the Number of Tree Species in Tropical Forests. *The American Naturalist*, **104**, 501–528.
- Kim, T.N. & Underwood, N. (2015) Plant neighborhood effects on herbivory: damage is both density and frequency dependent. *Ecology*, **96**, 1431–1437.
- Klink, C.A. & Machado, R.B. (2005) Conservation of the Brazilian Cerrado. *Conservation Biology*, **19**, 707–713.
- Lebrija-Trejos, E., Wright, S.J., Hernández, A. & Reich, P.B. (2014) Does relatedness matter? Phylogenetic density-dependent survival of seedlings in a tropical forest. *Ecology*, **95**, 940–951.
- Lewinsohn, T.M., Freitas, A.V.L. & Prado, P.I. (2005) Conservation of terrestrial invertebrates and their habitats in Brazil. *Conservation Biology*, **19**, 640–645.
- Loranger, J., Meyer, S.T., Shipley, B., Kattge, J., Loranger, H., Roscher, C. & Weisser, W.W. (2012) Predicting invertebrate herbivory from plant traits: evidence from 51 grassland species in experimental monocultures. *Ecology*, **93**, 2674–2682.
- Marquis, R.J., Diniz, I.R. & Morais, H.C. (2001) Patterns and correlates of interspecific variation in foliar insect herbivory and pathogen attack in Brazilian cerrado. *Journal of Tropical Ecology*, **17**, 127–148.
- Meijden, E. (1996) Plant defence, an evolutionary dilemma: contrasting effects of (specialist and generalist) herbivores and natural enemies. *Entomologia Experimentalis et Applicata*, **80**, 307–310.
- Mundim, F.M., Costa, A.N. & Vasconcelos, H.L. (2009) Leaf nutrient content and host plant selection by leaf-cutter ants, *Atta laevigata*, in a Neotropical savanna. *Entomologia Experimentalis et Applicata*, **130**, 47–54.
- Murphy, S.J., Xu, K. & Comita, L.S. (2016) Tree seedling richness, but not

- neighborhood composition, influences insect herbivory in a temperate deciduous forest community. *Ecology and Evolution*, **6**, 6310–6319.
- Pearse, I.S. & Hipp, A.L. (2009) Phylogenetic and trait similarity to a native species predict herbivory on non-native oaks. *Proceedings of the National Academy of Sciences*, **106**, 18097–18102.
- Peel, M.C., Finlayson, B.L. & McMahon, T.A. (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences*, **11**, 1633–1644.
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M.S., Cornwell, W.K., Craine, J.M., Gurvich, D.E., Urcelay, C., Veneklaas, E.J., Reich, P.B., Poorter, L., Wright, I.J., Ray, P., Enrico, L., Pausas, J.G., de Vos, A.C., Buchmann, N., Funes, G., Quétier, F., Hodgson, J.G., Thompson, K., Morgan, H.D., ter Steege, H., Sack, L., Blonder, B., Poschlod, P., Vaieretti, M. V., Conti, G., Staver, A.C., Aquino, S. & Cornelissen, J.H.C. (2013) New handbook. *Australian Journal of Botany*, **61**, 167.
- Queenborough, S.A., Burslem, D.F.R.P., Garwood, N.C. & Valencia, R. (2007) Neighborhood and community interactions determine the spatial patterns of tropical tree seedling survival. *Ecology*, **88**, 2248–2258.
- R Core Team, R. (2016) R: A language and environment for statistical computing. R foundation for statistical computing.
- Ramos-Neto, M.B. & Pivello, V.R. (2000) Lightning fires in a Brazilian savanna national park: rethinking management strategies. *Environmental Management*, **26**, 675–684.
- Ratter, J.A., Ribeiro, J.F. & Bridgewater, S. (1997) The Brazilian Cerrado vegetation and threats to its biodiversity. *Annals of Botany*, **80**, 223–230.

- Ribeiro, S.P., Carneiro, M.A.A. & Fernandes, G.W. (1998) Free-feeding insect herbivores along environmental gradients in Serra do Cipó: basis for a management plan. *Journal of Insect Conservation*, **2**, 107–118.
- Ribeiro, J.F. & Walter, B.M.T. (2008) As principais fitofisionomias do bioma Cerrado. *Cerrado: ecologia e flora* (eds S.M. Sano, S.P. Almeida & J.F. Ribeiro), pp. 151–212. Embrapa Cerrados, Brasília.
- Root, R.B. (1973) Organization of a plant-arthropod association in simple and diverse habitats: the fauna of Collards (*Brassica oleracea*). *Ecological Monographs*, **43**, 95–124.
- Satterthwaite, F.E. (1946) An approximate distribution of estimates of variance components. *Biometrics Bulletin*, **2**, 110.
- Scherrer, S., Lepesqueur, C., Vieira, M.C., Almeida-Neto, M., Dyer, L., Forister, M. & Diniz, I.R. (2016) Seasonal variation in diet breadth of folivorous Lepidoptera in the Brazilian cerrado. *Biotropica*, **48**, 491–498.
- Schuldt, A., Bruelheide, H., Durka, W., Eichenberg, D., Fischer, M., Kröber, W., Härdtle, W., Ma, K., Michalski, S.G., Palm, W.-U., Schmid, B., Welk, E., Zhou, H. & Assmann, T. (2012) Plant traits affecting herbivory on tree recruits in highly diverse subtropical forests (ed V Novotny). *Ecology Letters*, **15**, 732–739.
- Silva, D.M. da & Batalha, M.A. (2008) Soil-vegetation relationships in cerrados under different fire frequencies. *Plant and Soil*, **311**, 87–96.
- Silva, D.M. da & Batalha, M.A. (2011) Defense syndromes against herbivory in a cerrado plant community. *Plant Ecology*, **212**, 181–193.
- Simon, M.F., Grether, R., de Queiroz, L.P., Skema, C., Pennington, R.T. & Hughes, C.E. (2009) Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by in situ evolution of adaptations to fire. *Proceedings of the National Academy of*

- Sciences of the United States of America*, **106**, 20359–64.
- Twigg, L.E. & Socha, L. V. (1996) Physical versus chemical defence mechanisms in toxic *Gastrolobium*. *Oecologia*, **108**, 21–28.
- Underwood, N., Inouye, B.D. & Hambäck, P.A. (2014) A conceptual framework for associational effects: when do neighbors matter and how would we know? *The Quarterly Review of Biology*, **89**, 1–19.
- Uriarte, M., Swenson, N.G., Chazdon, R.L., Comita, L.S., John Kress, W., Erickson, D., Forero-Montaña, J., Zimmerman, J.K. & Thompson, J. (2010) Trait similarity, shared ancestry and the structure of neighbourhood interactions in a subtropical wet forest: implications for community assembly. *Ecology Letters*, **13**, 1503–1514.
- Zanne, A.E., Tank, D.C., Cornwell, W.K., Eastman, J.M., Smith, S.A., FitzJohn, R.G., McGlinn, D.J., O'Meara, B.C., Moles, A.T., Reich, P.B., Royer, D.L., Soltis, D.E., Stevens, P.F., Westoby, M., Wright, I.J., Aarssen, L., Bertin, R.I., Calaminus, A., Govaerts, R., Hemmings, F., Leishman, M.R., Oleksyn, J., Soltis, P.S., Swenson, N.G., Warman, L. & Beaulieu, J.M. (2014) Three keys to the radiation of angiosperms into freezing environments. *Nature*, **506**, 89–92.
- Zava, P.C. & Cianciaruso, M. V. (2014) Can we use plant traits and soil characteristics to predict leaf damage in savanna woody species? *Plant Ecology*, **215**, 625–637.

Supporting information

Table S1. List of families, species and number of individuals recorded in this study. The species were identified and organized following the last Angiosperm Phylogeny Group classification (APG-IV 2016). Bold type indicates the 27 cerrado most abundant and frequent species utilized in the analyzes

Family	Species	Individuals
Annonaceae	<i>Annona crassifolia</i> Mart.	4
	<i>Annona coriacea</i> Mart.	1
	<i>Bocageopsis mattogrossensis</i> (R.E.Fr.) R.R.Fr.	1
Apocynaceae	<i>Aspidosperma macrocarpon</i> Mart.	1
	<i>Aspidosperma nobile</i> Müll.Arg	1
	<i>Hancornia speciosa</i> Gomes	1
Araliaceae	<i>Schefflera macrocarpa</i> (Cham. & Schltdl.) Frodin	11
	<i>Schefflera vinosa</i> (Cham. & Schltdl.) Frodin & Fiaschi	13
	<i>Eremanthus erythropappus</i> (DC.) MacLeish	24
	<i>Piptocarpha rotundifolia</i> (Less.) Baker	25
Bignoniaceae	<i>Handroanthus ochraceus</i> (Cham.) Mattos	16
	<i>Tabebuia aurea</i> (Silva Manso) Benth. & Hook.f. ex S.Moore	2
Calophyllaceae	<i>Kielmeyera coriacea</i> Mart. & Zucc.	6
	<i>Kielmeyera grandiflora</i> (Wawra) Saddi	7
	<i>Kielmeyera rubriflora</i> Cambess.	3
Caryocaraceae	<i>Caryocar brasiliense</i> Cambess.	14
Celastraceae	<i>Plenckia populnea</i> Reissek	1
Chrysobalanaceae	<i>Licania humilis</i> Cham. & Schltdl.	5
Connaraceae	<i>Connarus suberosus</i> Planch.	22
	<i>Rourea induta</i> Planch.	5
Dilleniaceae	<i>Davilla elliptica</i> A.St.-Hil.	13
Ebenaceae	<i>Diospyros hispida</i> A.DC.	35
Erythroxylaceae	<i>Erythroxylum engleri</i> O.E.Schulz	23
	<i>Erythroxylum suberosum</i> A.St.-Hil.	27
Fabaceae	<i>Anadenanthera peregrina</i> (L.) Spe.g.,	10
	<i>Andira cujabensis</i> Benth.	6
	<i>Bowdichia virgilioides</i> Kunth	7

Family	Species	Individuals
	<i>Dimorphandra mollis</i> Benth.	10
	<i>Hymenaea stigonocarpa</i> Mart. ex Hayne	9
	<i>Leptolobium dasycarpum</i> Vogel	11
	<i>Mimosa amnis-atric</i> Barneby	1
	<i>Tachigali aurea</i> Tul.	2
	<i>Tachigali vulgaris</i> L.G.Silva & H.C.Lima	2
	<i>Stryphnodendron adstringens</i> (Mart.) Coville	22
	<i>Stryphnodendron rotundifolium</i> Mart.	3
Malpighiaceae	<i>Byrsonima basiloba</i> A.Juss.	3
	<i>Byrsonima coccolobifolia</i> Kunth	26
	<i>Byrsonima pachyphylla</i> A.Juss.	4
	<i>Byrsonima verbascifolia</i> (L.) DC.	4
Malvaceae	<i>Eriotheca gracilipes</i> (K.Schum.) A.Robyns	6
	<i>Eriotheca pubescens</i> (Mart. & Zucc.) Schott & Endl.	13
Melastomataceae	<i>Miconia albicans</i> (Sw.) Triana	91
	<i>Miconia ferruginata</i> DC.	2
	<i>Miconia ligustroides</i> (DC.) Naudin	4
	<i>Mouriri elliptica</i> Mart.	1
Myrtaceae	<i>Campomanesia adamantium</i> (Cambess.) O.Berg	2
	<i>Eugenia aurata</i> O.Berg	2
	<i>Eugenia puniceifolia</i> (Kunth) DC.	7
	<i>Eugenia ternatifolia</i> Cambess.	2
	<i>Eugenia</i> sp.	1
	<i>Myrcia bella</i> Cambess.	73
	<i>Myrcia camapuanensis</i> N.Silveira	11
	<i>Myrcia guianensis</i> (Aubl.) DC.	15
	<i>Myrcia multiflora</i> (Lam.) DC.	1
	<i>Myrcia variabilis</i> DC.	1
	<i>Myrcia vestita</i> DC.	3
	<i>Myrcia</i> sp.	2
	<i>Psidium laruotteanum</i> Cambess.	54
Nyctaginaceae	<i>Guapira noxia</i> (Netto) Lundell	6
Ochnaceae	<i>Ouratea hexasperma</i> (A.St.-Hil.) Baill.	65
	<i>Ouratea spectabilis</i> (Mart.) Engl.	21
Proteaceae	<i>Roupala montana</i> Aubl.	24

Family	Species	Individuals
Rubiaceae	<i>Palicourea rigida</i> Kunth	7
Salicaceae	<i>Casearia sylvestris</i> Sw.	9
Sapotaceae	<i>Pouteria ramiflora</i> (Mart.) Radlk.	91
	<i>Pouteria torta</i> (Mart.) Radlk.	51
Styracaceae	<i>Styrax ferrugineus</i> Nees & Mart.	4
Vochysiaceae	<i>Qualea grandiflora</i> Mart.	12
	<i>Qualea multiflora</i> Mart.	1
	<i>Qualea parviflora</i> Mart.	12
	<i>Vochysia cinnamomea</i> Pohl	5

Table S2. Predictor variables used in the model and their definition.

Predictor variables	Definition
Conspecific abundance (AB)	The number of individuals of the same species in the same plot.
Phylogenetic isolation (PI)	The total sum of branches separating each individual and all other heterospecific individuals in the same plot.
Relative SLA (SLA)	The difference between the specific leaf area (SLA) of the individual and the mean SLA of the other individuals in the plot.
Relative height	The difference between the plant height of the individual and the mean height of the other individuals in the plot.
Relative leaf toughness (LT)	The difference between the leaf toughness (TO) of the individual and the mean TO of the other individuals in the plot.




Figure S1. Relationship between the relative traits and absolute trait values focal plant and the herbivore leaf damage on 27 cerrado species in Emas National Park, Brazil. The shaded bands indicate the 95% confidence interval of the response.

Conclusão Geral

Neste trabalho demonstramos que:

- Para ambientes contrastantes que ocorrem lado a lado como florestas estacionais e cerrado sentido restrito, as estratégias ecológicas das espécies estão de acordo com os fatores limitantes para a ocorrência de espécies em cada um destes habitats. Atributos aquisitivos que representam habilidade competitiva e rápida aquisição de recursos foram relacionados para espécies especialistas de floresta, enquanto atributos conservativos que promove resistência contra o fogo e estresse ambiental foram associados a espécies especialistas de savanas. Além disso, a plasticidade fenotípica pode determinar a habilidade das espécies em habitats contrastantes como florestas estacionais e cerrado sentido restrito.
- Espécies de comunidades de florestas estacionais e cerrado sentido restrito de fato adotam diferentes estratégias relacionada ao uso de recursos. Nós demonstramos uma forte relação negativa de densidade de indivíduos com o gradiente de fertilidade e toxicidade do solo. Para os valores médios dos atributos, nós registramos que em solos mais férteis e menos tóxicos tem espécies com atributos relacionados à estratégias de aquisição de recursos, enquanto solos com baixa fertilidade e mais tóxicos tem espécies com atributos associados à estratégias de conservação de recursos.
- Em ambientes savânicos, a distância ecológica e evolutiva de uma espécie focal para a sua vizinhança não é fundamental para determinar o nível de dano foliar por herbívoros. Sugerimos que outros fatores, tais como dominância de herbívoros

generalistas, co-evolução de plantas e herbívoros e consumo de folhas jovens são mais importantes para determinar o dano foliar do que o contexto da vizinhança.