

Short communication

Pandora bullata (Entomophthoromycota: Entomophthorales) affecting calliphorid flies in central Brazil

Cristian Montalva^a, Karin Collier^b, Christian Luz^{a,*}, Richard A. Humber^c

^a Laboratório de Patologia de Invertebrados (LPI), Instituto de Patologia Tropical e Saúde Pública (IPTSP), Universidade Federal de Goiás (UFG), Goiânia, Goiás, Brazil

^b Centro Universitário de Gurupí (UnirG), Gurupí, Tocantins, Brazil

^c USDA-ARS Emerging Pests and Pathogens Research Unit, Robert W. Holley Center for Agriculture and Health, Ithaca, NY, USA

ARTICLE INFO

Article history:

Received 31 December 2015
Received in revised form 28 February 2016
Accepted 5 March 2016
Available online 8 March 2016

Keywords:

Diptera
Natural enemy
Entomopathogenic fungi
Entomophthoromycota
Chrysomya

ABSTRACT

Fungi are where one finds them, and if one seeks fungal pathogens affecting flies, then a garbage dump may be an ideal place to find both persistent, abundant fly populations and their fungal pathogens. An obvious fungal epizootic affecting the oriental latrine fly, *Chrysomya megacephala* (Diptera: Calliphoridae), was observed over several days in mid-February 2015 at the local garbage dump adjacent to the city of Cavalcante, northern Goiás. This site harbored large populations of both *C. megacephala* and a *Musca* sp. (Diptera: Muscidae) but only the population of oriental latrine fly was affected by any fungal pathogen and presented unusually dense populations of fresh cadavers. The fungus was identifiable as *Pandora bullata* (Entomophthorales: Entomophthoraceae) only after a very small number of characteristically decorated resting spores were found in these flies two months later; this represents the first Brazilian (and South American) record of this species. *P. bullata* is known previously from a small number of North American, European and Australian collections, all of which have included relatively abundant production of resting spores. We cannot dismiss the possibility that the extremely sparse formation of resting spores at this Brazilian site may be due to abiotic factors such as latitude (13°46'40.53"S), day length, ambient temperatures, or even the precipitation patterns in this mid-tropical montaine site. Epizootic events affecting calliphorids in Brazil strengthen the interest in entomophthoran pathogens for biological control of flies.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Blowflies (Diptera: Calliphoridae) play a crucial role in the balance of land ecosystems, however they are also responsible for great damage through the traumatic myiasis that their larvae cause when burrowing the skin of domestic and wild animals, and even humans (Guimarães et al., 1982; Sukontason et al., 2005; Fernandes et al., 2009; Mello et al., 2009; Kuria et al., 2015). Among the many calliphorids affecting human and animal populations, the oriental latrine fly, *Chrysomya megacephala*, feeds and breeds in carrion, feces, and is an important indicator species in forensic entomology (Goff, 2001) because of its ready infestation of animal cadavers and carrion. Maggots of this species can, however, also serve positively in the treatment of cuticular wounds and

lesions (Pineiro et al., 2015). *C. megacephala* can carry and transfer the poliovirus, salmonellae, and other enteric pathogens among humans living under substandard conditions of sanitation and nutrition (Baumgartner and Greenberg, 1984; Olsen et al., 1993; Sukontason et al., 2007), as well as transferring helminth parasites to humans and poultry (Sulaiman et al., 1989; Avancini and Ueta, 1990; Monzon et al., 1991).

What little is yet known on the biodiversity of entomopathogenic fungi in Brazil (Sosa-Gómez et al., 2010) comes primarily from agricultural sites. The control of free-flying insects with entomopathogenic fungi has not been very widely considered because of the difficulty of applying the pathogen to the insects (Wright et al., 2004). Nevertheless, the development of trap-and-target technology, in which visual and semiochemical attractants are used to draw insects to a point source, makes such applications a realistic possibility (Muirhead-Thomson, 1991), and there can be no doubt that the populations of noxious and deleterious flies is a major concern that has received too little scientific scrutiny.

* Corresponding author at: Instituto de Patologia Tropical e Saúde Pública, UFG, Avenida Esperança s/n, Campus Samambaia, 74690-900 Goiânia, Goiás, Brazil.
E-mail address: wchrisluz@hotmail.com (C. Luz).

The diversity of entomophthoran fungi in Brazil is vastly less well documented than the more numerous and taxonomically diverse entomopathogenic conidial and teleomorphic ascomycetes from the order Hypocreales. Not surprisingly, extremely little is known about the diversity of entomophthoran fungi from hosts in nonagricultural sites in Brazil (or most other parts of the world).

As a result of our seeking fungal pathogens of dipterans at a distinctly nonagricultural, fly-infested garbage dump in central Brazil, we are able here to report the first South American record of the entomophthoran fungus *Pandora bullata* (Thaxter & MacLeod ex Humber) Humber occurring in a natural epizootic affecting adults of the oriental latrine fly, *C. megacephala* (Diptera: Calliphoridae).

2. Materials and methods

Surveys of entomopathogenic fungi on dipterans were done in a garbage dump adjacent to the city of Cavalcante, State of Goiás, in Central Brasil (13°46'40.53"S; 47°27'08.71"W) on 12–14 February and 3–5 April 2015. Twigs and leaves bearing mycotized cadavers were carefully removed, put into paper bags, and transported in a polystyrene cooler at 20°C (Benjamin et al., 2004) for later examination and laboratory processing. In the laboratory, all specimens were examined immediately or stored at 4°C. Cadavers filled with vegetative cells were incubated at 25°C in a moist chamber (a plastic bag with moistened filter paper) to induce external conidiogenesis. Semi-permanent slide mounts were prepared in lactophenol-aceto-orcein (LPAO) as described by Keller (1987). Fungal microstructures were examined by brightfield or phase contrast microscopy (Leica DMLS 020-518.500), measured microscopically (Nova 180i-T; Toupview), and documented with a digital camera (Leica UCMOS01300KPA). The micrographs of resting spores of this fungus are photomontages of four to eight focal planes of these structures taken with a compound microscope (Olympus BX-51, Olympus Corporation of America) with planapochromatic objectives to increase the apparent depth of focus, and were made using Helicon Focus Pro software (<http://www.heliconsoft.com>). Fifty measurements for any given structure usually came from fungus on several individual specimens.

3. Results and discussion

In February, it was impossible to overlook a very active fungal epizootic affecting oriental latrine flies, *C. megacephala*, when huge numbers of cadavers were visible on plants, fence wires and other surfaces in the local garbage dump (Fig. 1). The outbreak was observed over three consecutive days. This site also included very large populations of both *C. megacephala* and a *Musca* species (Diptera: Muscidae), but only the calliphorids were found to be mycotized and their cadavers were present in unusually dense populations (Fig. 1). During the survey in April, very few calliphorid individuals were found at the site; similarly small populations of the muscid flies were present. We are unable to account for the decreases in observable populations of both fly species as a matter of seasonality or some other physical factor although it appears that populations of this host may be positively correlated with temperature (Seolin Dias et al., 2009; Ngoen-klan et al., 2011). We do not know whether the calliphorid population was largely eliminated by the action of the *Pandora* species, but no mycotized individuals of *Musca* were found on this site during any of the surveys. A total of 261 mycotized calliphorid cadavers were collected in February, but only five individuals in April.

When the cadavers were put into high relative humidity, the further development of the fungus was stimulated, and characteristic light brown to white bands and clusters of actively sporulating conidiophores emerged through abdominal and other


intersegmental membranes (Fig. 2). The bitunicate, uninucleate primary conidia of this pathogen, $29.6 \pm 0.4 \times 13.8 \pm 0.2 \mu\text{m}$ [$23.2\text{--}34.8 \times 11.5\text{--}16.3 \mu\text{m}$] (Fig. 3), were formed singly and forcibly discharged from conidiogenous cells formed singly at the apices of the digitately branched conidiophores typical of *Pandora* species (Humber, 2012a). Rhizoids with large, discoid holdfasts emerged singly and in small bundles from the host's abdomen and served to anchor the cadavers to twigs, leaves, and wires on which the host died (Fig. 4). *Chrysomya* cadavers collected in early April included extremely small numbers of resting spores, $43.1 \pm 1.5 \mu\text{m}$ [$37.3\text{--}53.5 \mu\text{m}$] overall diameter (including the superficial decorations), with prominent bullate (rounded, bump-like) projections of the epispore's surface (Fig. 5). The combination of the host identity and the broadly rounded (bullate) decorations of these resting spores are unambiguously diagnostic for *P. bullata*. Humber (1981) described this fungus as *Erynia bullata* Thaxter & MacLeod by combining two earlier but nomenclaturally invalid descriptions of *Entomophthora bullata* by Povah (1935) who transcribed Roland Thaxter's herbarium notes about a set of infected flies, and by MacLeod et al. (1973) who provided a thorough characterization of this fungus. The species was later transferred to the newly recognized genus *Pandora* (Humber, 1989). The infected calliphorids in these collections were deposited in the Herbarium of the Federal University of Goiás, Goiânia, Brazil (UFG 50067, UFG 50069 and UFG 50071).

P. bullata was known previously from few collections in the northern states of the United States and adjacent areas of Canada (MacLeod et al., 1973) and, less prominently, from Australia, the United Kingdom, and Spain (Glare and Milner, 1987; Balazy, 1993; Niell and Santamaria, 2001). This is also the first record of *P. bullata* from any tropical location. The ARSEF culture collection (Ithaca, NY) lost its only isolate of *P. bullata* long ago before ARSEF acquired more specialized equipment for controlled cryopreservation of its isolates; the only available culture of *P. bullata* might be the American Type Culture Collection's 24298 (which was derived from the now defunct ARSEF 116).

As with the great majority of entomophthoran taxa, few gene sequence data are yet available to suggest or to verify individual species identifications, and since we were not able to isolate *P. bullata* in culture, we have not attempted to add any sequences for the Cavalcante fungus. For most of these fungi there is no option except to use traditional morphological, developmental and pathobiological characters as the bases for species identifications. In the case of *P. bullata*, the morphology of the resting spores coupled with the host identity is unambiguously diagnostic.

We attempted to culture this fungus in the field on quarter-strength Sabouraud dextrose agar amended with 0.2% yeast (SDAY/4) (Inglis et al., 2012). None of these attempts was successful using the media available to us at that place and time. The greatest likelihood of isolating a culture of this fungus would have been to use vegetative (hyphal or conidiophore) inoculum and any of a wide range of liquid culture media. Because the vegetative cells of many entomophthoraceous fungi such as this grow in the hosts as wall-less protoplasts, culture isolations are often more successful in liquid media than on solid culture media (Humber, 2012b), and then many (but not all) can be transferred to grow successfully as walled hyphae on solid media.

The resting spores of entomophthoran fungi often do not form until the end of a disease outbreak, and when these spores are formed the host cadavers are almost always substantially filled with resting spores in temperate regions. That so few resting spores were recovered from this site literally in the middle of the southern tropic zone raises interesting questions that, while unanswered, need to be asked: Were these few flies with resting spores collected too late in the disease cycle? Might some cadavers with resting spores have fallen off the substrates or been eaten by birds, rodents, or



Figs. 1–5. *Pandora bullata* affecting *Chrysomya megacephala*. (1) Dense clustering of mycotized fly cadavers on vegetation in Cavalcante garbage dump. (2) Clearly defined abdominal bands of conidiophores. (3) Bitunicate primary conidia in which the outer layer of the cell wall partially detaches from the inner wall layer; these conidia are uninucleate but the nuclei are not visible in this image. (4) Two superimposed holdfasts at the apices of rhizoids; these disks attach the fly to the substrate. (5) Three thick-walled resting spores showing characteristic raised, rounded bumps (bullations) on the outer wall layer. Bars (Figs. 3–5) = 20 μ m.

insectivores? Might resting spores have been formed more prominently at some point between the February and April collections of these flies on this site but the cadavers containing them no longer in the same substrates in April from which the earlier, conidial material was collected? Most intriguingly, however, we cannot know whether abiotic factors such as the prevalent temperature, day length (i.e., the lack of large fluctuations in day length through the season), precipitation patterns, or even more factors that characterize such a tropical site may not be favorable to the formation of entomophthoran resting spores. Very little is known about how pathogens such as *P. bullata* behave in tropical collection sites in comparison to their better studied biology and greater apparent presence in temperate zones, but the fundamental absence of a resistant spore type for an entomopathogenic fungus (that routinely forms these spores from temperate collection sites) has also been observed with a conspicuous absence of the formation of diagnostically ornamented oogonia for a wide range of central Brazilian collections of *Leptolegnia chapmanii* (Peronosporomycetes: Saprolegniales) (Montalva et al., 2016).

C. megacephala is known to be a prominent nuisance fly in garbage dumps in Brazil (Ferreira and Lacerda, 1993; Seolin Dias et al., 2009). The populations of this fly have been shown to be directly proportional to overall ambient temperatures but to be comparatively unaffected by changing rainfall (Seolin Dias et al., 2009; Ngoen-klan et al., 2011). The decline of populations of *C. megacephala* observed in Cavalcante between February and April does appear to reflect the much cooler (mid-autumn) weather in that city in April than in the (summer) heat of February. Curiously, in a distinctly different region of Brazil, it was found that the populations of *C. megacephala* appeared to be positively correlated with overall rainfall patterns but were little affected by temperature (Mello et al., 2007). Independent of any effects of the changing seasonal weather, there is a possibility that the heavy infestation of the oriental latrine flies observed in February might have contributed a direct and significant decline in the numbers of these flies on this site by the time of the April collection.

We are not aware of any intentional applications of entomopathogenic fungi to attempt to control flies on sites of this type, but the epizootic reported here suggests that garbage dumps and their high densities of nuisance fly species might be ideal and novel venues for inoculative or inundative applications of fungal biocontrol agents to help to limit the populations of these pests.

Acknowledgment

This study was supported by CAPES (Coordination of the Improvement of Higher Education) Science Without Borders project PVE 71/2013, 149988.

References

- Avancini, R.M.P., Ueta, M.T., 1990. Manure breeding insects (Diptera and Coleoptera) responsible for cestidiosis in caged layer hens. *J. Appl. Entomol.* 110, 307–312.
- Balazy, S., 1993. Flora of Poland (Flora Polska). Fungi (Mycota): Entomophthorales. Polish Academy of Science, vol. 24, W. Szafer Institute of Botany, Kraków.
- Baumgartner, D., Greenberg, B., 1984. The genus *Chrysomya* (Diptera: Calliphoridae) in the new world. *J. Med. Entomol.* 21, 105–113.
- Benjamin, R.K., Blackwell, M., Chapela, I.H., Humber, R.A., Jones, K.G., Klepzig, K.D., Lichtwardt, R.W., Malloch, D., Noda, H., Roeper, R.A., Spatafora, J.W., Weir, A., 2004. Insect- and other arthropod-associated fungi. In: Mueller, G.M., Bills, G.F., Foster, M. (Eds.), *Biodiversity of Fungi: Inventory and Monitoring Methods*. Elsevier Academic Press Amsterdam, pp. 395–433.
- Fernandes, L.F., Pimenta, F.C., Fernandes, F.F., 2009. First report of human myiasis in Goiás state, Brazil: frequency of different types of myiasis, their various etiological agents, and associated factors. *J. Parasitol.* 95, 32–38.
- Ferreira, M.J.M., Lacerda, P.V., 1993. Muscoides sinantrópicos associados ao lixo urbano em Goiânia, Goiás. *Rev. Bras. Zool.* 10, 185–195.
- Glare, T.R., Milner, R.J., 1987. New records of entomophthoran fungi from insects in Australia. *Aust. J. Bot.* 35, 69–78.
- Goff, M.L., 2001. *A Fly for the Prosecution: How Insect Evidence Helps Solve Crimes*. Harvard University Press, Boston.
- Guimarães, J.H., Papavero, N., Prado, A., 1982. As míases na região neotropical (identificação biológica, bibliografia). *Rev. Bras. Zool.* 1, 239–416.
- Humber, R.A., 1981. *Erynia* (Zygomycetes: Entomophthorales): validations and new species. *Mycotaxon* 13, 471–480.
- Humber, R.A., 1989. Synopsis of a revised classification for the Entomophthorales (Zygomycotina). *Mycotaxon* 31, 441–460.
- Humber, R.A., 2012a. Identification of entomopathogenic fungi. In: Lacey, L.A. (Ed.), *Manual of Techniques in Invertebrate Pathology*. Academic Press, London, UK, pp. 151–187.
- Humber, R.A., 2012b. Entomogenous fungi. In: Schaechter, M. (Ed.), *Eukaryotic Microbes*. Academic Press, San Diego, CA, pp. 127–141.
- Inglis, G.D., Enkerli, J., Goettel, M.S., 2012. Laboratory techniques used for entomopathogenic fungi: Hypocreales. In: Lacey, L.A. (Ed.), *Manual of Techniques in Invertebrate Pathology*. London, UK, Academic Press, pp. 189–253.
- Keller, S., 1987. Arthropod-pathogenic Entomophthorales of Switzerland. I. *Conidiobolus*, *Entomophaga* and *Entomophthora*. *Sydowia* 40, 122–167.
- Kuria, S.K., Kingu, H.J.C., Villet, M.H., Dhaffala, A., 2015. Human myiasis in rural South Africa is under-reported. *S. Afr. Med. J.* 105, 129–133.
- MacLeod, D.M., Tyrrell, D., Soper, R.S., De Lyzer, A.J., 1973. *Entomophthora bullata* as a pathogen of *Sarcophaga aldrichi*. *J. Invertebr. Pathol.* 22, 75–79.
- Mello, R.S., Quiroz, M.M.C., Aguiar-Coelho, V.M., 2007. Population fluctuations of calliphorid species (Diptera Calliphoridae) in the Biological Reserve of Tingua, state of Rio de Janeiro, Brazil. *Iheringia Ser. Zool.* 97, 481–485.
- Mello, R.S., Queiroz, M.M.C., Nunes-Freitas, A.F., Aguiar-Coelho, V.M., 2009. Calliphorid fly (Diptera Calliphoridae) attraction to different colored traps in the Tingua Biological Reserve, Rio de Janeiro, Brazil. *Iheringia Sér. Zool.* 99, 426–430.
- Montalva, C., dos Santos, K., Collier, K., Fernandes, É.K.K., Luz, C., Humber, R.A., 2016. First report of *Leptolegnia chapmanii* (Peronosporomycetes: Saprolegniales) affecting mosquitoes in central Brazil. *J. Invertebr. Pathol.* (in revision).
- Monzon, R.B., Sanchez, A.R., Tadiaman, B.M., Najos, O.A., Valencia, E.G., Rueda, R.R., Ventura, J.V.M., 1991. A comparison of the role of *Musca domestica* (Linnaeus) and *Chrysomya megacephala* (Fabricius) as mechanical vectors of helminth parasites in a typical slum area of metropolitan Manila. *Southeast Asian J. Trop. Med. Public Health* 22, 222–228.
- Muirhead-Thomson, R.C., 1991. *Trap Responses of Flying Insects*. Academic Press, San Diego.
- Ngoen-klan, R., Moophayak, K., Klong-klaw, T., Irvine, K.N., Sukantason, K., Prangkio, C., Somoon, P., 2011. Do climatic and physical factors affect populations of the blow fly, *Chrysomya megacephala*, and house fly, *Musca domestica*? *Parasitol. Res.* 109, 1279–1292.
- Niell, M., Santamaria, S., 2001. Additions to the knowledge of entomopathogenic Entomophthorales (Fungi, Zygomycota) from Spain. *Nova Hedwigia* 73, 167–197.
- Olsen, A.R., Sidebottom, T.H., Bennett, S.G., 1993. The Oriental latrine fly, *Chrysomya megacephala* (Fabricius 1794) (Diptera Calliphoridae), as an invading blow fly of public health importance. *Bull. Soc. Vector Ecol.* 18, 133–146.
- Pinheiro, M.A.R.Q., Ferraz, J.B., Miguel Junior, A.A., Moura, A.D., Costa, M.E.S.M., Costa, F.J.M.D., Neto, V.F.A., Neto, R.M., Gama, R.A., 2015. Use of maggot therapy for treating a diabetic foot ulcer colonized by multidrug resistant bacteria in Brazil. *Indian J. Med. Res.* 141, 340–342.
- Povah, A.H.W., 1935. The fungi of Isle Royale: Lake Superior. *Papers Michigan Acad. Sci. Arts Let.* 20, 113–156.
- Seolin Dias, L., Santarem, V.A., Almeida, M.S.R., Medina, A.O., Silva, A.V., 2009. Biodiversidade de moscas calliphoridae no lixo urbano de Presidente Prudente São Paulo, Brazil. *Arq. Inst. Biol. São Paulo* 76, 659–663.
- Sosa-Gómez, D.R., López-Lastra, C.C., Humber, R.A., 2010. An overview of arthropod-associated fungi from Argentina and Brazil. *Mycopathologia* 170, 61–76.
- Sukontason, K.L., Narongchai, P., Sripakdee, D., Boonchu, N., Chaiwong, T., Ngeru-Klun, R., Piangjai, S., Sukontason, K., 2005. First report of human myiasis caused by *Chrysomya megacephala* and *Chrysomya rufifacies* (Diptera: Calliphoridae) in Thailand, and its implication in forensic entomology. *J. Med. Entomol.* 42, 702–704.
- Sukontason, K.L., Bunchoo, M., Khantawa, B., Piangjai, S., Rongsriyam, Y., Sukontason, K., 2007. Comparison between *Musca domestica* and *Chrysomya megacephala* as carriers of bacteria in northern Thailand. *Southeast Asian J. Trop. Med. Public Health* 38, 38–44.
- Sulaiman, S., Sohadi, A.R., Jeffery, J., 1989. Human helminth parasite burdens on clyorrhaphan flies (Diptera) trapped at an aboriginal settlement in Malaysia. *Bull. Entomol. Res.* 79, 625–629.
- Wright, C., Brooks, A., Wall, R., 2004. Toxicity of the entomopathogenic fungus, *Metarhizium anisopliae* (Deuteromycotina: Hyphomycetes) to adult females of the blowfly *Lucilia sericata* (Diptera: Calliphoridae). *Pest Manag. Sci.* 60, 639–644.