

Review



Cite this article: Carvalheiro LG, Bartomeus I, Rollin O, Timóteo S, Tinoco CF. 2021 The role of soils on pollination and seed dispersal. *Phil. Trans. R. Soc. B* **376**: 20200171. <https://doi.org/10.1098/rstb.2020.0171>

Accepted: 21 December 2020

One contribution of 17 to a theme issue ‘The role of soils in delivering Nature’s Contributions to People’.

Subject Areas:
ecology

Keywords:
flower visitors, pollination, seed dispersal, frugivory, soil chemical and physical properties, soil microbiota

Author for correspondence:
Luísa G. Carvalheiro
e-mail: lgcarvalheiro@gmail.com

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5490651>.

The role of soils on pollination and seed dispersal

Luísa G. Carvalheiro^{1,2}, Ignasi Bartomeus³, Orianne Rollin², Sérgio Timóteo⁴ and Carla Faleiro Tinoco¹

¹Departamento de Ecologia, Universidade Federal de Goiás, 74001-970 Goiânia, Brasil

²Centre for Ecology, Evolution and Environmental Changes (CE3c), Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

³Estación Biológica de Doñana (EBD-CSIC), 41092 Sevilla, Spain

⁴Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, Calçada Martim de Freitas, 3000-456 Coimbra, Portugal

ORCID LGC, 0000-0001-7655-979X; IB, 0000-0001-7893-4389; OR, 0000-0002-1455-9302; ST, 0000-0003-2417-3259

Ongoing environmental changes are affecting physical, chemical and biological soil components. Evidence of impacts of soil changes on pollinators’ and seed dispersers’ behaviour, fitness and density is scarce, but growing. Here, we reviewed information on such impacts and on a number of mechanisms that may explain its propagation, taking into account the full range of resources required by the large and diverse number of species of these two important functional groups. We show that while there is substantial evidence on the effects of soil nitrogen enrichment and changes in soil water content on the quality and quantity of floral and fruit resources, little is known on the effects of changes of other soil properties (e.g. soil pH, soil structure, other nutrients). Also, the few studies showing correlations between soil changes and pollinator and seed disperser foraging behaviour or fitness do not clearly identify the mechanisms that explain such correlation. Finally, most studies (including those with nitrogen and water) are local and limited to a small number of species, and it remains unclear how variable such effects are across time and geographical regions, and the strength of interactive effects between soil properties. Increasing research on this topic, taking into consideration how impacts propagate through species interaction networks, will provide essential information to predict impacts of ongoing environmental changes and help guide conservation plans that aim to minimize impacts on ecosystem functioning.

This article is part of the theme issue ‘The role of soils in delivering Nature’s Contributions to People’.

1. Introduction

Pollination and seed dispersal are essential ecosystem functions upon which the reproduction of most plants depends [1,2] both being performed by a vast group of animals including a diverse set of invertebrates and vertebrates (figure 1). These functions make an important contribution to humankind, increasing production and quality of the vast majority of crops [3,4] and maintaining the populations of many plants [2] essential for several ecosystem services. Human activities have changed immensely chemical [5,6], physical [7,8] and biological [9] properties of the soil affecting plants at the species [10,11] and community level [12]. Yet, compared to other global changes (e.g. biological invasions, pollution, climate and land use changes [13,14]), little is known about how impacts of such changes in soil properties propagate through trophic levels and even less about the direct impacts on pollinators and seed dispersers [15,16]. In addition, the scarce existing information shows that impacts of soil changes vary in strength and direction depending on the species of the consumer evaluated. For example, some flower visitor species increase their visitation rates with nitrogen enrichment, while other species have their visitation rates maintained




Figure 1. Diversity of feeding resources used by pollinators, during larval or adult stages, that can be affected by soil properties. Dashed lines represent resources used during larval stages. References used for the construction of this figure are listed in the electronic supplementary material.

or even reduced [17,18]. Similarly, some vertebrate dispersers benefit from soil nutrient enrichment, while ant dispersers seem more frequent on infertile soils [16,19]. Diet preferences may partly explain such variability in species responses [20,21], which effects scale up to shape communities (e.g. increased dominance [18], reduced density and richness [21–23] and changes in overall interaction network patterns [24]). Understanding the mechanisms that mediate the propagation of soil changes to pollinators and seed dispersers is essential to improve our ability to predict impacts of ongoing environmental changes and define adequate conservation plans. Here, we explore potential mechanisms mediating impacts on pollinators and seed dispersers by summarizing the existing information on the impacts of changes of several soil properties on the great diversity of dietary (figure 1) and nesting resources of these important functional groups and how such changes can explain the reported effects on them (figure 2).

2. Effects mediated by changes on abundance and diversity of floral and fruit resources

Soil abiotic and biotic characteristics are important regulators of plant physiology and affect the spatial abundance and diversity of flowers and fruits [25–28]. While the effects of increasing

water availability on flower and fruit production tend to be generally positive, most plant species are better adapted to soils with low levels of nutrients, with nitrogen enrichment leading to local losses of plant diversity and promoting the expansion of invasive plants [25,26]. Similarly, soil phosphorous enrichment, which facilitates nitrogen uptake by plants [6], can lead to losses of plant richness [29]. Even in regions where plant richness is recovering (e.g. northwest Europe), increases of species are dominated by nitrophilous plants, benefitting only those pollinators that are able to make use of such species [21]. Other important macronutrients (phosphorus, potassium, calcium) may also affect flower availability, but are still poorly studied. On the other hand, whereas fertile soils have a higher percentage of fleshy-fruit species, infertile soils are more likely to have a community of plants that develop elaiosomes and arils (important rewards for many dispersers, including ants, birds and mammals [16,30,31]). Indeed, soils rich in potassium, calcium and phosphorus, are associated with vertebrate-attracting fleshy fruits, while in the nutrient-poor soil, seed dispersal communities may have a greater proportion of ants [19].

Effects of changes in soil microbiota on individual flower production can also vary among species. A reduction in the activity of soil microbiota may hence induce accentuated changes in plant community composition [32], the flower




Figure 2. Pathways through which changes in soil properties (chemical, physical and biotic) may affect pollinators and seed dispersers. Blue dashed arrows represent effects on nesting and dietary resources used mostly by immature stages, orange solid arrows represent effects on dietary resources collected by adult individuals and orange thin dashed line represents direct consumption of soil by adults.

abundance of some species being positively affected by mycorrhizal fungi colonization ([33,34]), while others exhibit a negative or null response [33].

Phenology can also be affected by soil changes (e.g. nutrients [35], soil microclimate [36–38]), affecting the abundance of flowers and fruits through time, potentially creating temporal mismatches between plants and their pollinators and seed dispersers [39,40]. Some species have their flowering periods anticipated by increased nitrogen availability [41,42], while in others, flowering may be delayed or shortened [43,44]. Drought may delay the flowering period of certain plant species (e.g. [36–38]).

The changes in plant distribution, flower production and phenology described above may affect the availability of dietary resources with adequate nutritional content for pollinators and seed dispersers. Moreover, even when flower resource availability and assemblage composition are not affected, soil-driven changes in plant morphology, physiology and chemistry can have strong impacts on their consumers [45]. Below we describe in more detail effects of soil changes on such properties.

3. Effects mediated by changes on fruit and flower morphology

Alteration of soil nutrient levels [15,46], pH [47] and humidity [36,48] may change flower and fruit morphology. For example, under water restriction, certain plant species produce smaller flowers [36,48], while under lower soil pH, some species produce smaller inflorescences [47]. Nutrient increases have been related both to increases [46,49] and decreases [50] in fruit and flower size (e.g. corolla length, petal width), and soil changes may also affect flower and fruit colour [51,52]. Such changes can affect attractiveness to pollinators [53,54] and seed dispersers [16], or access of pollinators to floral resources [36,48]. Other morphological changes caused by soil changes (e.g. nutrient level, mycorrhizal fungi) may involve pollen

sculpture [55] and size [33,56], which can affect pollen adherence to pollinator body and their performance as pollinators [57,58].

4. Effects mediated by changes on nutritional value of resources

Similar to other guilds of primary consumers, to satiate the dietary needs of pollinators and dispersers, the nutritional composition of plant resources consumed must match, at least partially, the requirements of those animals, which can greatly vary across species [21,59–62]. Nutrient enrichment may affect nectar [49,63,64] and pollen [56] production per flower, negative effects on nectar volume being more likely when nitrogen input is high [49,50]. Nectar and pollen quality [65–67] (i.e. amino acid (AA) content, sugar or secondary compounds levels) may also be affected by nutrient enrichment. For example, increasing nitrogen availability can increase nectar AAs and sugar content, some AAs being more affected than others [42,64,68–71], with climate mediating the strength of such effects [42]. Indeed, Gardener & Gillman [68] showed that glutamine and proline, key AAs in pollinator nectar selection, exhibit a large and significant increase under fertilizer treatment. Soil fertility (especially nitrogen and phosphorus) and irrigation may also affect the production of essential oils consumed by multiple flower visitor species [72–74]. For example, soil phosphorus increased the yield of essential oil in *Achillea millefolium* L. [75]. Yet, it is unclear if similar effects can be detected in non-volatile (fatty) floral oils, and if such effects depend on soil nutrients and water levels. Changes on mycorrhizal fungi root colonization can also affect the quality of nectar, affecting sugar content [33,34,76], flavonoids and other secondary compounds (e.g. [77–80]) found in floral resources.

Fruit quality may also be affected by soil nutrient enrichment [30,31]. An increase in potassium level may lead to more nutritious fruits [46], rich in sugars and AAs [30,81], and overall

benefits for fleshy-fruited plants [19,82]. Increases in secondary compounds caused by nitrogen enrichment can be toxic when ingested [83,84], which many dispersers can counter by ingesting clay [83]. Soil ingestion may also help to compensate for fruit deficits in nitrogen and calcium [85,86]. In addition, soils with low nutrient levels tend to have plants with drier and harder fruits [30,31] and elaiosomes and arils are more common [19]. The production of edible fruits can also be limited by soil water stress, reducing fruit fleshiness and energetic content available to animals [87,88].

While many species of these two groups of animals have a generalized diet, resource quality changes mentioned above can have a strong influence on the foraging behaviour and fitness of pollinators [89,90] and seed dispersers [19,68,90–92], and potentially lead to local losses of diversity [93]. However, these effects may greatly vary among species. Low levels of nitrogen enrichment may be beneficial to some pollinator species (e.g. by reducing parasite loads [94,95]), but repel pollinators and negatively affect their physiology at high concentrations [96]. Finally, plants growing in heavy-metal-rich soils can accumulate metals into their nectar, shortening foraging time of pollinators and nectar robbers, leading to an overall positive effect on fitness of some plant species (e.g. [97]), possibly owing to increased cross-pollination.

5. Effects mediated by changes on floral and fruit scent

While vastly understudied, there is evidence that soil property changes can affect secondary compounds related to floral volatile compounds. Nitrogen enrichment can increase a phenylpropanoid floral volatile (eugenol) that attracts pollinators [17], while the number of floral volatiles and total fragrance emission can decrease with mycorrhizal fungi colonization rate [76]. Earthworm-mediated changes in soil properties can also affect the production of defence-related phytohormone jasmonic acid and of phenolic compounds [98]. Reduced soil humidity is also thought to limit the emission of olfactory cues used by dispersers to find seeds [59]. Given the importance of floral and fruit scent for detecting the presence of resources [60–62,84], any change in chemical composition of olfactory cues can affect pollinator and seed disperser foraging activity. Indeed, previous studies have suggested nitrogen-induced changes in floral volatiles increase pollinator visitation rates [17] and overall plant–pollinator communication [99].

6. Effects mediated by changes on dietary non-floral and non-fruit resources

Recognizing the diversity of resources used by consumers is essential to understand the mechanisms by which soil changes can affect them. While tight coevolutionary processes have resulted in highly specialized relationships between animals and plants [100], many pollinators feed on multiple species across multiple families [101,102], some including a variety of non-floral resources, especially during immature life stages (figure 1). Among seed dispersers, there is also variation in specialization levels, some being almost exclusive frugivorous species [1], while others include multiple alternative resources (figure 1). For pollinators that feed on leaves, decaying material, soil fungi and plant roots during immature stages (e.g.

Lepidoptera and Syrphidae [103–107]), changes in leaf biomass, nutrient content and chemical clues caused by increased nutrient level or other soil changes affect the behaviour (e.g. oviposition, consumption patterns) and physiology of those insects [107–110]. Some species of pollinators also act as predators during larval stages, hence, depending on the populations of their prey. Such prey are frequently herbivores (e.g. aphids [110]) and, consequently are highly susceptible to plant chemical properties which are affected by soil properties [111]. Moreover, for pollinators highly specialized in their oviposition locations (e.g. on nitrophobous plant species [21,112]), any change in plant community composition will change oviposition opportunities. However, little is known on how changes in soil chemical and physical conditions affect larval stages of pollinators and seed dispersers and their role in ecosystem functioning.

7. Effects mediated by changes on nesting resources

Several pollinators and seed dispersers have a central place foraging pattern around a nesting location (e.g. bees, ants and vertebrates). For species constructing nests aboveground, nesting requirements are related to habitat structural complexity [28,113], which can be affected by soil properties (e.g. [114,115]). For those directly using soil as nesting substrate (e.g. the vast majority of non-parasitic bees [116,117] and several ant species) and for those that use soil to construct their cells above ground (e.g. Megachillidae [118]), the impacts of altered soil properties can be profound [119].

Although ground-nesting species can adapt to distinct soil types [120–122], some studies suggest soil texture is important. For example, many bee species tend to prefer sandy soils [122,123], deserts and dunes hosting a large diversity of bees, while clay or silt soils are less favourable. Slope [124] and soil compaction [125] also influence choice of nesting sites. While some bee communities prefer flat areas with little compaction [126], others require steep and sloping ground [127], or compacted soil in irregular surfaces [123,125]. Soil temperature and humidity can affect oviposition and the availability of nesting area for pollinators and seed dispersers (e.g. [128–131]). Central place foragers tend to minimize the difference between nest microclimate and optimal environmental foraging conditions [132], and exposed bare ground [126,127], litter cover [133], or direct sunlight and warmth [125] are documented to be preferred by some species.

Anthropogenic activities (e.g. agriculture, pasture or mining) can constrain nest availability [130,134] owing to intense soil disturbance and changes in the physical characteristics of soils and habitat structure [134]. Agricultural soils are known nesting sites of pollinators (wild bees) and seed dispersers (ants, birds) and, despite the limited knowledge, agricultural practices such as tillage or pesticide use may directly harm nest sites. Anecdotal evidence suggests that tillage can directly impact bee survival and delay emergence time [125,135]. For example, for squash bees, *Peponapis pruinose*, it has been reported that tilling can halve offspring survival owing to direct destruction of nests which, if males and females are at different depths, may also affect sex ratios [136]. However, tillage may also have positive effects on soil properties for ground-nesting by creating open bare ground, loosening compacted soils or changing the predator community [137]. In addition, in agricultural land, ground-nesting pollinators

and seed dispersers can also be exposed to pesticides that accumulate in the soil (e.g. [138]). Substantial knowledge gaps remain around the toxicity and effects of neonicotinoids to arthropods, including ground-nesting bees. Further research on pollinator and seed disperser soil preferences is needed, and citizen science approaches integrating large-scale nesting occurrences documented by volunteers [139] with soil texture maps is a promising avenue to advance in this regard.

8. Effects of pollinators and seed dispersers on soil properties

Pollinators and seed dispersers are important in soil formation and maintenance. Soil nesting species impact soil characteristics owing to their excavation activities promoting vertical and horizontal mobilization of soil [116,140,141]. For example, a single bee species (*Nomia meander*) can move around 95 t yr⁻¹ in the Touchet Valley of southeastern Washington [142]. Such activity may affect soil physical, chemical and hydrological profiles [141], as demonstrated in soils surrounding ants' nests [143]. This can be particularly relevant in arid and semi-arid ecosystems, where soil changes promoted by ants can generate islands of fertility [144]. Moreover, the ramified nests of communal nesters (i.e. bees nesting in aggregations) create an extensive network of tunnels that contribute to soil aeration. In addition, bees are known to bring in external soil into nests to complement its construction, contributing to soil mixing [121]. Overall, like earthworms, bees and ants contribute largely to soil aeration and rejuvenation.

9. Pathways for future research

Ongoing environmental changes are affecting soil in all its components. Evidence of impacts of soil changes on pollinators' and seed dispersers' behaviour, fitness and density is growing. While we have reviewed a number of mechanisms that may explain how such impacts propagate to these important groups of ecosystem service providers, most studies

focus only on a subset of the pathways (e.g. effect of soil changes on resources, effects of resource changes on consumers). More experimental studies testing the full chain of effects from soil changes to both resources and consumers are needed. Moreover, it is likely that there is plenty of geographical variation in such impacts. Indeed, Zhong *et al.* [145] found differences in effects of nitrogen enrichment on soil respiration across biomes, with stronger negative effects in forests than in deserts. Global studies comparing strength and type of effects across climates and biomes are essential to better understand temporal and spatial variations of such effects. Moreover, much of the evidence on potential mechanisms presented here comes from a limited number of studies, and substantial knowledge gaps exist on the effects of most soil characteristics. Also, most studies focus on changes in nitrogen and water content, with more limited evidence on the effects of soil structure, pH, biota and other soil nutrients and pollutants. Experimental manipulation of other soil variables would help to disentangle interactive effects between multiple soil characteristics. Finally, multi-disciplinary studies involving different guilds would help understand the indirect impacts of soil changes on pollinators and seed dispersers. For example, as pollinators have a strong influence on fruit abundance and quality, it is likely that any impact of soil changes on pollinators propagate to frugivores, affecting seed dispersal.

Data accessibility. This article has no additional data.

Authors' contributions. L.G.C. and I.B. conceived the initial structure of the manuscript. All authors contributed to the literature review and manuscript writing. L.G.C. coordinated and integrated all contributions.

Competing interests. We declare we have no competing interests.

Funding. L.G.C. and O.R. were funded by Fundação para Ciência e Tecnologia (FCT) and European Union via the programa operacional regional de Lisboa 2014/2020 (project no. EUCLIPO-028360). L.G.C. was also funded by the Brazilian National Council for Scientific and Technological Development (CNPq. Universal 421668/2018-0; PQ 305157/2018-3). S.T. was funded by Fundação para Ciência e Tecnologia (FCT: grant no. UID/BIA/04004/2020 and contract CEECIND/00135/2017). I.B. was funded by H2020 project SHOW-CASE. C.F.T. was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

References

- Jordano P. 2000 Fruits and frugivory. In *Seeds: the ecology of regeneration in plant communities* (ed. M Fenner), pp. 125–165. Wallingford, UK: CABI.
- Ollerton J, Winfree R, Tarrant S. 2011 How many flowering plants are pollinated by animals? *Oikos* **120**, 321–326. (doi:10.1111/j.1600-0706.2010.18644.x)
- Garibaldi LA *et al.* 2013 Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science* **339**, 1608–1611. (doi:10.1126/science.1230200)
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) 2016 The assessment report on pollinators, pollination and food production. (doi:10.5281/ZENODO.3402857)
- Fowler D *et al.* 2013 The global nitrogen cycle in the twenty-first century. *Phil. Trans. R. Soc. B* **368**, 20130164. (doi:10.1098/rstb.2013.0164)
- Smil V. 2000 Phosphorus in the environment: natural flows and human interferences. *Annu. Rev. Energy Environ.* **25**, 53–88. (doi:10.1146/annurev.energy.25.1.53)
- Bai Z *et al.* 2018 Effects of agricultural management practices on soil quality: a review of long-term experiments for Europe and China. *Agric. Ecosyst. Environ.* **265**, 1–7. (doi:10.1016/j.agee.2018.05.028)
- Fageria NK. 2002 Soil quality vs. environmentally-based agricultural management practices. *Commun. Soil Sci. Plant Anal.* **33**, 2301–2329. (doi:10.1081/CSS-120005764)
- Zogg GP, Zak DR, Ringelberg DB, White DC, MacDonald NW, Pregitzer KS. 1997 Compositional and functional shifts in microbial communities due to soil warming. *Soil Sci. Soc. Am. J.* **61**, 475–481. (doi:10.2136/sssaj1997.03615995006100020015x)
- Hageman RH. 1980 Effect of form of nitrogen on plant growth. In *Nitrification inhibitors—potentials and limitations* (eds JJ Meisinger, GW Randall, ML Vitosh), pp. 47–62. Maddison, WI: American Society of Agronomy and Soil Science Society of America.
- Vitousek PM, Howarth RW. 1991 Nitrogen limitation on land and in the sea: how can it occur? *Biogeochemistry* **13**, 87–115. (doi:10.1007/BF00002772)
- Tilman D, Lehman C. 2001 Human-caused environmental change: impacts on plant diversity and evolution. *Proc. Natl Acad. Sci. USA* **98**, 5433–5440. (doi:10.1073/pnas.091093198)
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010 Global pollinator declines: trends, impacts and drivers. *Trends Ecol. Evol.* **25**, 345–353. (doi:10.1016/j.tree.2010.01.007)

14. González-Varo JP *et al.* 2013 Combined effects of global change pressures on animal-mediated pollination. *Trends Ecol. Evol.* **28**, 524–530. (doi:10.1016/j.tree.2013.05.008)
15. David TI, Storkey J, Stevens CJ. 2019 Understanding how changing soil nitrogen affects plant–pollinator interactions. *Arthropod. Plant. Interact.* **13**, 671–684. (doi:10.1007/s11829-019-09714-y)
16. Westoby M, French K, Hughes L, Rice B, Rodgeron L. 1991 Why do more plant species use ants for dispersal on infertile compared with fertile soils? *Aust. J. Ecol.* **16**, 445–455. (doi:10.1111/j.1442-9993.1991.tb01074.x)
17. Majetic CJ, Fetters AM, Beck OM, Stachnik EF, Beam KM. 2017 *Petunia* floral trait plasticity in response to soil nitrogen content and subsequent impacts on insect visitation. *Flora Morphol. Distrib. Funct. Ecol. Plants* **232**, 183–193. (doi:10.1016/j.flora.2016.08.002)
18. de Ramos DL, Bustamante MMC, Da Silva E, Silva FD, Carvalheiro LG. 2018 Crop fertilization affects pollination service provision—common bean as a case study. *PLoS ONE* **13**, 1–16. (doi:10.1371/journal.pone.0204460)
19. Milewski AV. 1982 The occurrence of seeds and fruits taken by ants versus birds in Mediterranean Australia and Southern Africa, in relation to the availability of soil potassium. *J. Biogeogr.* **9**, 505–516. (doi:10.2307/2844617)
20. Pöyry J, Carvalheiro LG, Heikkinen RK, Kühn I, Kuussaari M, Schweiger O, Valtonen A, van Bodegom PM, Franzén M. 2017 The effects of soil eutrophication propagate to higher trophic levels. *Glob. Ecol. Biogeogr.* **26**, 18–30. (doi:10.1111/geb.12521)
21. Carvalheiro LG *et al.* 2020 Soil eutrophication shaped the composition of pollinator assemblages during the past century. *Ecography* **43**, 209–221. (doi:10.1111/ecog.04656)
22. Öckinger E, Hammarstedt O, Nilsson SG, Smith HG. 2006 The relationship between local extinctions of grassland butterflies and increased soil nitrogen levels. *Biol. Conserv.* **128**, 564–573. (doi:10.1016/j.biocon.2005.10.024)
23. Van Dyck H, Van Strien AJ, Maes D, Van Swaay CAM. 2009 Declines in common, widespread butterflies in a landscape under intense human use. *Conserv. Biol.* **23**, 957–965. (doi:10.1111/j.1523-1739.2009.01175.x)
24. Burkle L, Irwin R. 2009 The importance of interannual variation and bottom-up nitrogen enrichment for plant–pollinator networks. *Oikos* **118**, 1816–1829. (doi:10.1111/j.1600-0706.2009.17740.x)
25. Tamis WLM, Van't Zelfde M, Van Der Meijden R, De Haes HAU. 2005 Changes in vascular plant biodiversity in the Netherlands in the 20th century explained by their climatic and other environmental characteristics. *Clim. Change* **72**, 37–56. (doi:10.1007/s10584-005-5287-7)
26. Bobbink R *et al.* 2010 Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecol. Appl.* **20**, 30–59. (doi:10.1890/08-1140.1)
27. McConkey KR, Prasad S, Corlett RT, Campos-Arceiz A, Brodie JF, Rogers H, Santamaria L. 2012 Seed dispersal in changing landscapes. *Biol. Conserv.* **146**, 1–13. (doi:10.1016/j.biocon.2011.09.018)
28. Plein M, Längsfeld L, Neuschulz EL, Schultheiß C, Ingmann L, Töpfer T, Böhning-Gaese K, Schleuning M. 2013 Constant properties of plant–frugivore networks despite fluctuations in fruit and bird communities in space and time. *Ecology* **94**, 1296–1306. (doi:10.1890/12-1213.1)
29. Wassen MJ, Schrader J, van Dijk J, Eppinga MB. 2021 Phosphorus fertilization is eradicating the niche of northern Eurasia's threatened plant species. *Nat. Ecol. Evol.* **5**, 67–73. (doi:10.1038/s41559-020-01323-w)
30. Hughes L, Westoby M, Johnson AD. 1993 Nutrient costs of vertebrate- and ant-dispersed fruits. *Funct. Ecol.* **7**, 54–62. (doi:10.2307/2389867)
31. Herrera CM. 1987 Vertebrate-dispersed plants of the Iberian Peninsula: a study of fruit characteristics. *Ecol. Monogr.* **57**, 305–331. (doi:10.2307/2937089)
32. Van Der Heijden MGA, Bardgett RD, Van Straalen NM. 2008 The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol. Lett.* **11**, 296–310. (doi:10.1111/j.1461-0248.2007.01139.x)
33. Barber NA, Soper Gorden NL. 2014 How do belowground organisms influence plant–pollinator interactions? *J. Plant Ecol.* **8**, 1–11. (doi:10.1093/jpe/rtu012)
34. Gange AC, Smith AK. 2005 Arbuscular mycorrhizal fungi influence visitation rates of pollinating insects. *Ecol. Entomol.* **30**, 600–606. (doi:10.1111/j.0307-6946.2005.00732.x)
35. Silvertown J, Poulton P, Johnston E, Edwards G, Heard M, Biss PM. 2006 The park grass experiment 1856–2006: its contribution to ecology. *J. Ecol.* **94**, 801–814. (doi:10.1111/j.1365-2745.2006.01145.x)
36. Caruso CM. 2006 Plasticity of inflorescence traits in *Lobelia siphilitica* (Lobeliaceae) in response to soil water availability. *Am. J. Bot.* **93**, 531–538. (doi:10.3732/ajb.93.4.531)
37. Desclaux D, Romet P. 1996 Impact of drought stress on the phenology of two soybean (*Glycine max* L. Merr) cultivars. *F. Crop Res.* **46**, 61–70. (doi:10.1016/0378-4290(95)00086-0)
38. McMaster GS, White JW, Weiss A, Stephen Baenziger P, Wilhelm WW, Porter JR, Jamieson PD. 2015 Simulating crop phenological responses to water deficits. In *Response of crops to limited water* (eds L Ahuja, V Reddy, S Saseendran, Q Yu), pp. 277–300. Hoboken, NJ: John Wiley & Sons, Ltd.
39. Schenk M, Krauss J, Holzschuh A. 2018 Desynchronizations in bee–plant interactions cause severe fitness losses in solitary bees. *J. Anim. Ecol.* **87**, 139–149. (doi:10.1111/1365-2656.12694)
40. Warren RJ, Bahn V, Bradford MA. 2011 Temperature cues phenological synchrony in ant-mediated seed dispersal. *Glob. Chang. Biol.* **17**, 2444–2454. (doi:10.1111/j.1365-2486.2010.02386.x)
41. Cleland EE, Chiariello NR, Loarie SR, Mooney HA, Field CB. 2006 Diverse responses of phenology to global changes in a grassland ecosystem. *Proc. Natl Acad. Sci. USA* **103**, 13 740–13 744. (doi:10.1073/pnas.0600815103)
42. Hoover SER, Ladley JJ, Shchetkina AA, Tisch M, Giese SP, Tylisanakis JM. 2012 Warming, CO₂, and nitrogen deposition interactively affect a plant–pollinator mutualism. *Ecol. Lett.* **15**, 227–234. (doi:10.1111/j.1461-0248.2011.01729.x)
43. Xia J, Wan S. 2013 Independent effects of warming and nitrogen addition on plant phenology in the Inner Mongolian steppe. *Ann. Bot.* **111**, 1207–1217. (doi:10.1093/aob/mct079)
44. Liu Y, Miao R, Chen A, Miao Y, Liu Y, Wu X. 2017 Effects of nitrogen addition and mowing on reproductive phenology of three early-flowering forb species in a Tibetan alpine meadow. *Ecol. Eng.* **99**, 119–125. (doi:10.1016/j.ecoleng.2016.11.033)
45. Muñoz AA, Celedon-Neghme C, Cavieres LA, Arroyo MTK. 2005 Bottom-up effects of nutrient availability on flower production, pollinator visitation, and seed output in a high-Andean shrub. *Oecologia* **143**, 126–135. (doi:10.1007/s00442-004-1780-3)
46. Reuther W, Embleton TW, Jones WW. 1958 Mineral nutrition of tree crops. *Annu. Rev. Plant Physiol.* **9**, 175–206. (doi:10.3168/lhdm.0849)
47. Gentili R, Ambrosini R, Montagnani C, Caronni S, Citterio S. 2018 Effect of soil pH on the growth, reproductive investment and pollen allergenicity of *Ambrosia artemisiifolia* L. *Front. Plant Sci.* **9**, 1335. (doi:10.3389/fpls.2018.01335)
48. Galen C. 2000 High and dry: drought stress, sex-allocation trade-offs, and selection on flower size in the alpine wildflower *Polemonium viscosum* (Polemoniaceae). *Am. Nat.* **156**, 72–83. (doi:10.1086/303373)
49. Burkle LA, Irwin RE. 2009 The effects of nutrient addition on floral characters and pollination in two subalpine plants, *Ipomopsis aggregata* and *Linum lewisii*. *Plant Ecol.* **203**, 83–98. (doi:10.1007/s11258-008-9512-0)
50. Burkle LA, Irwin RE. 2010 Beyond biomass: measuring the effects of community-level nitrogen enrichment on floral traits, pollinator visitation and plant reproduction. *J. Ecol.* **98**, 705–717. (doi:10.1111/j.1365-2745.2010.01648.x)
51. Miller R, Owens SJ, Rørslett B. 2011 Plants and colour: flowers and pollination. *Optics Laser Technol.* **43**, 282–294. (doi:10.1016/i.optlastec.2008.12.018)
52. Schaefer HM, Valido A, Jordano P. 2014 Birds see the true colours of fruits to live off the fat of the land. *Proc. R. Soc. B* **281**, 20132516. (doi:10.1098/rspb.2013.2516)
53. Kaczorowski RL, Seliger AR, Gaskett AC, Wigsten SK, Raguso RA. 2012 Corolla shape vs. size in flower choice by a nocturnal hawkmoth pollinator. *Funct. Ecol.* **26**, 577–587. (doi:10.1111/j.1365-2435.2012.01982.x)
54. Macukanovic-Jocic M, Stevanović ZD, Mladenović M, Jocić G. 2011 Flower morphophysiology of selected Lamiaceae species in relation to pollinator

- attraction. *J. Apic. Res.* **50**, 89–101. (doi:10.3896/IBRA.1.50.2.01)
55. Reshetova SA, Sumarokova IE. 2019 Pollen of plants in urban areas as a bioindicator of industrial pollution (Transbaikalia). *Geosfernye Issled.* **4**, 15–23. (doi:10.17223/25421379/13/2)
56. Lau T-C, Stephenson AG. 1993 Effects of soil nitrogen on pollen production, pollen grain size, and pollen performance in *Cucurbita pepo* (Cucurbitaceae). *Am. J. Bot.* **80**, 763–768. (doi:10.2307/2445596)
57. Hesse M. 2000 Pollen wall stratification and pollination. *Plant Syst. Evol.* **222**, 1–17. (doi:10.1007/BF00984093)
58. Tanaka N, Uehara K, Murata J. 2004 Correlation between pollen morphology and pollination mechanisms in the Hydrocharitaceae. *J. Plant Res.* **117**, 265–276. (doi:10.1007/s10265-004-0155-5)
59. Wall SBV. 2002 Secondary dispersal of Jeffrey pine seeds by rodent scatter-hoarders: the roles of pilfering, recaching and a variable environment. In *Seed dispersal and frugivory: ecology, evolution and conservation. Third International Symposium-Workshop on Frugivores and Seed Dispersal, São Pedro, BR.* (eds DJ Levey, WR Silva, M Galetti), pp. 193–208. Wallingford, UK: CABI Publishing.
60. Dötterl S, Schäffler I. 2007 Flower scent of floral oil-producing *Lysimachia punctata* as attractant for the oil-bee *Macropis fulvipipes*. *J. Chem. Ecol.* **33**, 441–445. (doi:10.1007/s10886-006-9237-2)
61. Wright GA, Schiestl FP. 2009 The evolution of floral scent: the influence of olfactory learning by insect pollinators on the honest signalling of floral rewards. *Funct. Ecol.* **23**, 841–851. (doi:10.1111/j.1365-2435.2009.01627.x)
62. Nevo O, Ayasse M. 2020 Fruit scent: biochemistry, ecological function, and evolution. In *Co-evolution of secondary metabolites* (eds JM Méridon, K Ramawat), pp. 403–425. Berlin, Germany: Springer Nature Switzerland.
63. Campbell DR, Halama KJ. 1993 Resource and pollen limitations to lifetime seed production in a natural plant population. *Ecology* **74**, 1043–1051. (doi:10.2307/1940474)
64. Gardener MC, Gillman MP. 2001 The effects of soil fertilizer on amino acids in the floral nectar of corncockle, *Agrostemma githago* (Caryophyllaceae). *Oikos* **92**, 101–106. (doi:10.1034/j.1600-0706.2001.920112.x)
65. Ceulemans T, Hulsmans E, Vanden Ende W, Honnay O. 2017 Nutrient enrichment is associated with altered nectar and pollen chemical composition in *Succisa pratensis* Moench and increased larval mortality of its pollinator *Bombus terrestris* L. *PLoS ONE* **12**, 1–15. (doi:10.1371/journal.pone.0175160)
66. Audusseau H, Kolb G, Janz N. 2015 Plant fertilization interacts with life history: variation in stoichiometry and performance in nettle-feeding butterflies. *PLoS ONE* **10**, 1–15. (doi:10.1371/journal.pone.0124616)
67. Kurze S, Heinken T, Fartmann T. 2017 Nitrogen enrichment of host plants has mostly beneficial effects on the life-history traits of nettle-feeding butterflies. *Acta Oecol.* **85**, 157–164. (doi:10.1016/j.actao.2017.11.005)
68. Gardener MC, Gillman MP. 2002 The taste of nectar—a neglected area of pollination ecology. *Oikos* **98**, 552–557. (doi:10.1034/j.1600-0706.2002.980322.x)
69. Gijbels P, Van den Ende W, Honnay O. 2014 Landscape scale variation in nectar amino acid and sugar composition in a Lepidoptera pollinated orchid species and its relation with fruit set. *J. Ecol.* **102**, 136–144. (doi:10.1111/1365-2745.12183)
70. Gijbels P, Ceulemans T, Van den Ende W, Honnay O. 2015 Experimental fertilization increases amino acid content in floral nectar, fruit set and degree of selfing in the orchid *Gymnadenia conopsea*. *Oecologia* **179**, 785–795. (doi:10.1007/s00442-015-3381-8)
71. Ceulemans T *et al.* 2017 Phosphorus resource partitioning shapes phosphorus acquisition and plant species abundance in grasslands. *Nat. Plants* **3**, 16224. (doi:10.1038/nplants.2016.224)
72. Machado IC. 2004 Oil-collecting bees and related plants: a review of the studies in the last twenty years and case histories of plants occurring in NE Brazil. In *Solitary bees, conservation, rearing and management for pollination.* (eds BM Freitas, JOP Pereira), pp. 255–280. Fortaleza, Brazil: Universidade federal do Ceará.
73. Rasmussen C, Olesen J. 2000 Oil flowers and oil-collecting bees. *Det. Nar. Videnskaps-akadem. I. Mat. Naturvidenskabelig Klasse, Skr. Ny Ser.* **39**, 23–31.
74. Buchmann SL. 1987 The ecology of oil flowers and their bees. *Annu. Rev. Ecol. Syst.* **18**, 343–369. (doi:10.1146/annurev.es.18.110187.002015)
75. Scheffer MC, Ronzelli JP, Koehler HS. 1993 Influence of organic fertilization on the biomass, yield and composition of the essential oil of *Achillea millefolium* L. *Acta Hortic.* **331**, 109–114. (doi:10.17660/actahortic.1993.331.14)
76. Becklin KM, Gamez G, Uelk B, Raguso RA, Galen C. 2011 Soil fungal effects on floral signals, rewards, and aboveground interactions in an alpine pollination web. *Am. J. Bot.* **98**, 1299–1308. (doi:10.3732/ajb.1000450)
77. Peipp H, Maier W, Schmidt J, Wray V, Strack D. 1997 Arbuscular mycorrhizal fungus-induced changes in the accumulation of secondary compounds in barley roots. *Phytochemistry* **44**, 581–587. (doi:10.1016/S0031-9422(96)00561-4)
78. Kapoor R, Anand G, Gupta P, Mandal S. 2017 Insight into the mechanisms of enhanced production of valuable terpenoids by arbuscular mycorrhiza. *Phytochem. Rev.* **16**, 677–692. (doi:10.1007/s11101-016-9486-9)
79. Araújo GC, Sousa NR, Castro PML. 2018 The effect of fungal-bacterial interaction on the phenolic profile of *Pinus pinea* L. *Plant Growth Regul.* **86**, 465–475. (doi:10.1007/s10725-018-0445-x)
80. Pedone-Bonfim MVL, da Silva DKA, da Silva-Batista AR, de Oliveira AP, da Silva Almeida JRG, Yano-Melo AM, Maia LC. 2018 Mycorrhizal inoculation as an alternative for the sustainable production of *Mimosa tenuiflora* seedlings with improved growth and secondary compounds content. *Fungal Biol.* **122**, 918–927. (doi:10.1016/j.funbio.2018.05.009)
81. Fischer RC, Richter A, Hadacek F, Mayer V. 2008 Chemical differences between seeds and elaiosomes indicate an adaptation to nutritional needs of ants. *Oecologia* **155**, 539–547. (doi:10.1007/s00442-007-0931-8)
82. Irion G. 1978 Soil infertility in the Amazonian rain forest. *Naturwissenschaften* **65**, 515–519. (doi:10.1007/BF00439791)
83. Cipollini ML, Levey DJ. 1997 Secondary metabolites of fleshy vertebrate-dispersed fruits: adaptive hypotheses and implications for seed dispersal. *Am. Nat.* **150**, 346–372. (doi:10.1086/286069)
84. Rodríguez A, Alquézar B, Peña L. 2013 Fruit aromas in mature fleshy fruits as signals of readiness for predation and seed dispersal. *New Phytol.* **197**, 36–48. (doi:10.1111/j.1469-8137.2012.04382.x)
85. Bravo A, Harms KE, Emmons LH. 2010 Puddles created by geophagous mammals are potential mineral sources for frugivorous bats (Stenodermatinae) in the Peruvian Amazon. *J. Trop. Ecol.* **26**, 173–184. (doi:10.1017/S0266467409990472)
86. Courts SE. 1998 Dietary strategies of old world fruit bats (Megachiroptera, Pteropodidae): how do they obtain sufficient protein? *Mamm. Rev.* **28**, 185–194. (doi:10.1046/j.1365-2907.1998.00033.x)
87. Bazzaz FA, Ackerly DD, Reekie EG. 2000 Reproductive allocation in plants. In *Seeds: the ecology of regeneration in plant communities* (ed. M Fenner), pp. 1–29. Wallingford, UK: CABI Publishing.
88. Willson MF, Irvine AK, Walsh NG. 1989 Vertebrate dispersal syndromes in some Australian and New Zealand plant communities, with geographic comparisons. *Biotropica* **21**, 133–147. (doi:10.2307/2388704)
89. Alm J, Ohnmeiss TE, Lanza J, Vriesenga L. 1990 Preference of cabbage white butterflies and honey bees for nectar that contains amino acids. *Oecologia* **84**, 53–57. (doi:10.1007/BF00665594)
90. Nepi M. 2014 Beyond nectar sweetness: the hidden ecological role of non-protein amino acids in nectar. *J. Ecol.* **102**, 108–115. (doi:10.1111/1365-2745.12170)
91. Givnish TJ. 1999 On the causes of gradients in tropical tree diversity. *J. Ecol.* **87**, 193–210. (doi:10.1046/j.1365-2745.1999.00333.x)
92. Vanderplanck M, Vereecken NJ, Grumiau L, Esposito F, Lognay G, Wattiez R, Michez D. 2017 The importance of pollen chemistry in evolutionary host shifts of bees. *Sci. Rep.* **7**, 1–10. (doi:10.1038/srep43058)
93. Elliott SE, Irwin RE, Adler LS, Williams NM. 2008 The nectar alkaloid, gelsemine, does not affect offspring performance of a native solitary bee, *Osmia lignaria* (Megachilidae). *Ecol. Entomol.* **33**, 298–304. (doi:10.1111/j.1365-2311.2007.00974.x)
94. Richardson LL, Adler LS, Leonard AS, Andicoechea J, Regan KH, Anthony WE, Manson JS, Irwin RE. 2015 Secondary metabolites in floral nectar reduce

- parasite infections in bumblebees. *Proc. R. Soc. B* **282**, 20142471. (doi:10.1098/rspb.2014.2471)
95. Richardson LL, Bowers MD, Irwin RE. 2016 Nectar chemistry mediates the behavior of parasitized bees: consequences for plant fitness. *Ecology* **97**, 325–337. (doi:10.1890/15-0263.1)
96. Manson JS, Cook D, Gardner DR, Irwin RE. 2013 Dose-dependent effects of nectar alkaloids in a montane plant-pollinator community. *J. Ecol.* **101**, 1604–1612. (doi:10.1111/1365-2745.12144)
97. Xun E, Zhang Y, Zhao J, Guo J. 2018 Heavy metals in nectar modify behaviors of pollinators and nectar robbers: consequences for plant fitness. *Environ. Pollut.* **242**, 1166–1175. (doi:10.1016/j.envpol.2018.07.128)
98. Xiao Z, Jiang L, Chen X, Zhang Y, Defossez E, Hu F, Liu M, Rasmann S. 2019 Earthworms suppress thrips attack on tomato plants by concomitantly modulating soil properties and plant chemistry. *Soil Biol. Biochem.* **130**, 23–32. (doi:10.1016/j.soilbio.2018.11.023)
99. Majetic CJ, Raguso RA, Ashman TL. 2009 Sources of floral scent variation. *Plant Signal. Behav.* **4**, 129–131. (doi:10.4161/psb.4.2.7628)
100. Anderson B, Johnson SD. 2008 The geographical mosaic of coevolution in a plant-pollinator mutualism. *Evolution* **62**, 220–225. (doi:10.1111/j.1558-5646.2007.00275.x)
101. Waser NM, Chittka L, Price MV, Williams NM, Ollerton J. 1996 Generalization in pollination systems, and why it matters. *Ecology* **77**, 1043–1060. (doi:10.2307/2265575)
102. Johnson SD, Steiner KE. 2000 Generalization versus specialization in plant pollination systems. *Trends Ecol. Evol.* **15**, 140–143. (doi:10.1016/S0169-5347(99)01811-X)
103. Ricarte A, Souba-Dols GJ, Hauser M, Marcos-García MÁ. 2017 A review of the early stages and host plants of the genera *Eumerus* and *Merodon* (Diptera: Syrphidae), with new data on four species. *PLoS ONE* **12**, e0189852. (doi:10.1371/journal.pone.0189852)
104. Skevington JH, Dang PT. 2002 Exploring the diversity of flies (Diptera). *Biodiversity* **3**, 3–27. (doi:10.1080/14888386.2002.9712613)
105. Szymank A, Kearns CA, Pape T, Thompson FC. 2008 Pollinating flies (diptera): a major contribution to plant diversity and agricultural production. *Biodiversity* **9**, 86–89. (doi:10.1080/14888386.2008.9712892)
106. Kristensen NP. 2003 *Band/Volume IV: Arthropoda: Insecta. Lepidoptera, moths and butterflies. Teilband/Part 36, vol. 2: morphology, physiology, and development*. Berlin, Germany, Boston, MA: De Gruyter. (doi:10.1515/9783110893724)
107. Prudic KL, Oliver JC, Bowers MD. 2005 Soil nutrient effects on oviposition preference, larval performance, and chemical defense of a specialist insect herbivore. *Oecologia* **143**, 578–587. (doi:10.1007/s00442-005-0008-5)
108. Butler J, Garratt MPD, Leather SR. 2012 Fertilisers and insect herbivores: a meta-analysis. *Ann. Appl. Biol.* **161**, 223–233. (doi:10.1111/j.1744-7348.2012.00567.x)
109. Li F, Dudley TL, Chen B, Chang X, Liang L, Peng S. 2016 Responses of tree and insect herbivores to elevated nitrogen inputs: a meta-analysis. *Acta Oecol.* **77**, 160–167. (doi:10.1016/j.actao.2016.10.008)
110. Verheggen FJ, Arnaud L, Bartram S, Gohy M, Haubruge E. 2008 Aphid and plant volatiles induce oviposition in an aphidophagous hoverfly. *J. Chem. Ecol.* **34**, 301–307. (doi:10.1007/s10886-008-9434-2)
111. Awmack CS, Leather SR. 2002 Host plant quality and fecundity in herbivorous insects. *Annu. Rev. Entomol.* **47**, 817–844. (doi:10.1146/annurev.ento.47.091201.145300)
112. WallisDeVries MF, van Swaay CAM. 2017 A nitrogen index to track changes in butterfly species assemblages under nitrogen deposition. *Biol. Conserv.* **212**, 448–453. (doi:10.1016/j.biocon.2016.11.029)
113. Breitbach N, Laube I, Steffan-Dewenter I, Böhning-Gaese K. 2010 Bird diversity and seed dispersal along a human land-use gradient: high seed removal in structurally simple farmland. *Oecologia* **162**, 965–976. (doi:10.1007/s00442-009-1547-y)
114. Roubik DW. 1989 *Ecology and natural history of tropical bees*. Cambridge, UK: Cambridge University Press.
115. Frankie GW, Newstrom L, Vinson SB, Barthell JF. 1993 Nesting-habitat preferences of selected *Centris* bee species in Costa Rican dry forest. *Biotropica* **25**, 322–333. (doi:10.2307/2388790)
116. Cane J, Neff J. 2011 Predicted fates of ground-nesting bees in soil heated by wildfire: thermal tolerances of life stages and a survey of nesting depths. *Biol. Conserv.* **144**, 2631–2636. (doi:10.1016/j.biocon.2011.07.019)
117. Kells AR, Goulson D. 2003 Preferred nesting sites of bumblebee queens (Hymenoptera: Apidae) in agroecosystems in the UK. *Biol. Conserv.* **109**, 165–174. (doi:10.1016/S0006-3207(02)00131-3)
118. Pinilla-Gallego MS, Crum J, Schaeztl R, Isaacs R. 2018 Soil textures of nest partitions made by the mason bees *Osmia lignaria* and *O. cornifrons* (Hymenoptera: Megachilidae). *Apidologie* **49**, 464–472. (doi:10.1007/s13592-018-0574-2)
119. Lybrand RA, Fedenko J, Tfaily M, Rao S. 2020 Soil properties and biochemical composition of ground-dwelling bee nests in agricultural settings. *Soil Sci. Soc. Am. J.* **84**, 1139–1152. (doi:10.1002/saj2.20085)
120. Orr MC, Griswold T, Pitts JP, Parker FD. 2016 A new bee species that excavates sandstone nests. *Curr. Biol.* **26**, R792–R793. (doi:10.1016/j.cub.2016.08.001)
121. Danforth BN, Minckley RL, Neff JL. 2019 *The solitary bees: biology, evolution, conservation*. Princeton, NJ, Oxford, UK: Princeton University Press.
122. Cane J. 1991 Soils of ground-nesting bees (Hymenoptera: Apoidea): texture, moisture, cell depth and climate. *J. Kansas Entomol. Soc.* **64**, 406–413.
123. Tsiolis K. 2018 Do bare soil landscapes encourage ground nesting bees? Masters thesis, Canterbury Christ Church University School of Human and Life Sciences, Canterbury, UK.
124. Burkle LA, Alarcón R. 2011 The future of plant-pollinator diversity: understanding interaction networks across time, space, and global change. *Am. J. Bot.* **98**, 528–538. (doi:10.3732/ajb.1000391)
125. Wuellner CT. 1999 Nest site preference and success in a gregarious, ground-nesting bee *Dieunomia triangulifera*. *Ecol. Entomol.* **24**, 471–479. (doi:10.1046/j.1365-2311.1999.00215.x)
126. Sardiñas HS, Kremen C. 2014 Evaluating nesting microhabitat for ground-nesting bees using emergence traps. *Basic Appl. Ecol.* **15**, 161–168. (doi:10.1016/j.baee.2014.02.004)
127. Potts SG, Vulliamy B, Roberts S, O'Toole C, Dafni A, Ne'eman G, Willmer P. 2005 Role of nesting resources in organising diverse bee communities in a Mediterranean landscape. *Ecol. Entomol.* **30**, 78–85. (doi:10.1111/j.0307-6946.2005.00662.x)
128. Hackwell GA. 1967 The biology and behavior of the alkali bee *Nomia Melanderi* Cockerell (hymenoptera: apoidea). Doctoral dissertation, Oregon State University, Corvallis, OR, USA.
129. Kudo G, Cooper EJ. 2019 When spring ephemerals fail to meet pollinators: mechanism of phenological mismatch and its impact on plant reproduction. *Proc. R. Soc. B* **286**, 20190573. (doi:10.1098/rspb.2019.0573)
130. Azcárate FM, Peco B. 2012 Abandonment of grazing in a mediterranean grassland area: consequences for ant assemblages. *Insect Conserv. Divers.* **5**, 279–288. (doi:10.1111/j.1752-4598.2011.00165.x)
131. Retana J, Cerdá X. 2000 Patterns of diversity and composition of Mediterranean ground ant communities tracking spatial and temporal variability in the thermal environment. *Oecologia* **123**, 436–444. (doi:10.1007/s004420051031)
132. Azcárate FM, Kovacs E, Peco B. 2007 Microclimatic conditions regulate surface activity in harvester ants *Messor barbarus*. *J. Insect Behav.* **20**, 315–329. (doi:10.1007/s10905-007-9074-3)
133. Grundel R, Jean RP, Frohnapple KJ, Glowacki GA, Scott PE, Pavlovic NB. 2010 Floral and nesting resources, habitat structure, and fire influence bee distribution across an open-forest gradient. *Ecol. Appl.* **20**, 1678–1692. (doi:10.1890/08-1792.1)
134. Azcárate FM, Peco B. 2003 Spatial patterns of seed predation by harvester ants (*Messor Forel*) in Mediterranean grassland and scrubland. *Insectes Soc.* **50**, 120–126. (doi:10.1007/s00040-003-0635-y)
135. Mathewson JA. 1968 Nest construction and life history of the eastern cucurbit bee, *Peponapis pruinosa* (Hymenoptera: Apoidea). *J. Kansas Entomol. Soc.* **41**, 255–261.
136. Ullmann KS, Meisner MH, Williams NM. 2016 Impact of tillage on the crop pollinating, ground-nesting bee, *Peponapis pruinosa* in California. *Agric. Ecosyst. Environ.* **232**, 240–246. (doi:10.1016/j.agee.2016.08.002)

137. Roger-Estrade J, Anger C, Bertrand M, Richard G. 2010 Tillage and soil ecology: partners for sustainable agriculture. *Soil Tillage Res.* **111**, 33–40. (doi:10.1016/j.still.2010.08.010)
138. Willis Chan DS, Prosser RS, Rodríguez-Gil JL, Raine NE. 2019 Assessment of risk to hoary squash bees (*Peponapis pruinosa*) and other ground-nesting bees from systemic insecticides in agricultural soil. *Sci. Rep.* **9**, 1–13. (doi:10.1038/s41598-019-47805-1)
139. Maher S, Manco F, Ings TC. 2019 Using citizen science to examine the nesting ecology of ground-nesting bees. *Ecosphere* **10**, e02911. (doi:10.1002/ecs2.2911)
140. Houston TF. 1984 Biological observations of bees in the genus *Ctenocolletes*. (Hymenoptera: Stenotritidae). *Rec. West. Aust. Mus.* **11**, 153–172.
141. Zaitlin B, Hayashi M. 2012 Interactions between soil biota and the effects on geomorphological features. *Geomorphology* **157–158**, 142–152. (doi:10.1016/j.geomorph.2011.07.029)
142. Cane JH. 2003 Annual displacement of soil in nest tumuli of alkali bees (*Nomia melanderi*) (Hymenoptera: Apiformes: Halictidae) across an agricultural landscape. *J. Kansas Entomol. Soc.* **76**, 172–176.
143. Cammeraat LH, Willott SJ, Compton SG, Incoll LD. 2002 The effects of ants' nests on the physical, chemical and hydrological properties of a rangeland soil in semi-arid Spain. *Geoderma* **105**, 1–20. (doi:10.1016/S0016-7061(01)00085-4)
144. Mandel RD, Sorenson CJ. 1982 The role of the western harvester ant (*Pogonomyrmex occidentalis*) in soil formation. *Soil Sci. Soc. Am. J.* **46**, 785–788. (doi:10.2136/sssaj1982.03615995004600040024x)
145. Zhong Y, Yan W, Shanguan Z. 2016 The effects of nitrogen enrichment on soil CO₂ fluxes depending on temperature and soil properties. *Glob. Ecol. Biogeogr.* **25**, 475–488. (doi:10.1111/geb.12430)